

8

Anthropogenic and Natural Radiative Forcing

Coordinating Lead Authors:

Gunnar Myhre (Norway), Drew Shindell (USA)

Lead Authors:

François-Marie Bréon (France), William Collins (UK), Jan Fuglestad (Norway), Jianping Huang (China), Dorothy Koch (USA), Jean-François Lamarque (USA), David Lee (UK), Blanca Mendoza (Mexico), Teruyuki Nakajima (Japan), Alan Robock (USA), Graeme Stephens (USA), Toshihiko Takemura (Japan), Hua Zhang (China)

Contributing Authors:

Borgar Aamaas (Norway), Olivier Boucher (France), Stig B. Dalsøren (Norway), John S. Daniel (USA), Piers Forster (UK), Claire Granier (France), Joanna Haigh (UK), Øivind Hodnebrog (Norway), Jed O. Kaplan (Switzerland/Belgium/USA), George Marston (UK), Claus J. Nielsen (Norway), Brian C. O'Neill (USA), Glen P. Peters (Norway), Julia Pongratz (Germany), Michael Prather (USA), Venkatachalam Ramaswamy (USA), Raphael Roth (Switzerland), Leon Rotstayn (Australia), Steven J. Smith (USA), David Stevenson (UK), Jean-Paul Vernier (USA), Oliver Wild (UK), Paul Young (UK)

Review Editors:

Daniel Jacob (USA), A.R. Ravishankara (USA), Keith Shine (UK)

This chapter should be cited as:

Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestad, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura and H. Zhang, 2013: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Table 8.7 | GWP and GTP with and without inclusion of climate–carbon feedbacks (cc fb) in response to emissions of the indicated non-CO₂ gases (climate-carbon feedbacks in response to the reference gas CO₂ are always included).

	Lifetime (years)		GWP ₂₀	GWP ₁₀₀	GTP ₂₀	GTP ₁₀₀
CH ₄ ^b	12.4 ^a	No cc fb	84	28	67	4
		With cc fb	86	34	70	11
HFC-134a	13.4	No cc fb	3710	1300	3050	201
		With cc fb	3790	1550	3170	530
CFC-11	45.0	No cc fb	6900	4660	6890	2340
		With cc fb	7020	5350	7080	3490
N ₂ O	121.0 ^a	No cc fb	264	265	277	234
		With cc fb	268	298	284	297
CF ₄	50,000.0	No cc fb	4880	6630	5270	8040
		With cc fb	4950	7350	5400	9560

Notes:

Uncertainties related to the climate–carbon feedback are large, comparable in magnitude to the strength of the feedback for a single gas.

^a Perturbation lifetime is used in the calculation of metrics.^b These values do not include CO₂ from methane oxidation. Values for fossil methane are higher by 1 and 2 for the 20 and 100 year metrics, respectively (Table 8.A.1).

and GTP. For the more long-lived gases the GWP₁₀₀ values increase by 10 to 12%, while for GTP₁₀₀ the increase is 20 to 30%. Table 8.A.1 gives metric values including the climate–carbon feedback for CO₂ only, while Supplementary Material Table 8.SM.16 gives values for all halocarbons that include the climate–carbon feedback. Though uncertainties in the carbon cycle are substantial, it is *likely* that including the climate–carbon feedback for non-CO₂ gases as well as for CO₂ provides a better estimate of the metric value than including it only for CO₂.

Emission metrics can be estimated based on a constant or variable background climate and this influences both the adjustment times and the concentration–forcing–temperature relationships. Thus, all metric values will need updating due to changing atmospheric conditions as well as improved input data. In AR5 we define the metric values with respect to a constant present-day condition of concentrations and climate. However, under non-constant background, Joos et al. (2013) found decreasing CO₂ AGWP₁₀₀ for increasing background levels (up to 23% for RCP8.5). This means that GWP for all non-CO₂ gases (except CH₄ and N₂O) would increase by roughly the same magnitude. Reisinger et al. (2011) found a reduction in AGWP for CO₂ of 36% for RCP8.5 from 2000 to 2100 and that the CH₄ radiative efficiency and AGWP also decrease with increasing CH₄ concentration. Accounting for both effects, the GWP₁₀₀ for CH₄ would increase by 10 to 20% under low and mid-range RCPs by 2100, but would decrease by up to 10% by mid-century under the highest RCP. While these studies have focused on the background levels of GHGs, the same issues apply for temperature. Olivé et al. (2012) find different temperature IRFs depending on the background climate (and experimental set up).

User related choices (see Box 8.4) such as the time horizon can greatly affect the numerical values obtained for CO₂ equivalents. For a change in time horizon from 20 to 100 years, the GWP for CH₄ decreases by a factor of approximately 3 and its GTP by more than a factor of 10. Short-lived species are most sensitive to this choice. Some approaches have removed the time horizon from the metrics (e.g., Boucher, 2012), but discounting is usually introduced which means that a discount rate

r (for the weighting function e^{-rt}) must be chosen instead. The choice of discount rate is also value based (see WGIII, Chapter 3).

For NTCFs the metric values also depend on the location and timing of emission and whether regional or global metrics are used for these gases is also a choice for the users. Metrics are usually calculated for pulses, but some studies also give metric values that assume constant emissions over the full time horizon (e.g., Shine et al., 2005a; Jacobson, 2010). It is important to be aware of the idealized assumption about constant future emissions (or change in emissions) of the compound being considered if metrics for sustained emissions are used.

8.7.1.5 New Metric Concepts

New metric concepts have been developed both to modify physical metrics to address shortcomings as well as to replace them with metrics that account for economic dimensions of problems to which metrics are applied. Modifications to physical metrics have been proposed to better represent CO₂ emissions from bioenergy, regional patterns of response, and for peak temperature limits.

Emissions of CO₂ from the combustion of biomass for energy in national emission inventories are currently assumed to have no net RF, based on the assumption that these emissions are compensated by biomass regrowth (IPCC, 1996). However, there is a time lag between combustion and regrowth, and while the CO₂ is resident in the atmosphere it leads to an additional RF. Modifications of the GWP and GTP for bioenergy (GWP_{bio}, GTP_{bio}) have been developed (Cherubini et al., 2011; Cherubini et al., 2012). The GWP_{bio} give values generally between zero (current default for bioenergy) and one (current for fossil fuel emissions) (Cherubini et al., 2011), and negative values are possible for GTP_{bio} due to the fast time scale of atmospheric–ocean CO₂ exchange relative to the growth cycle of biomass (Cherubini et al., 2012). GWP_{bio} and GTP_{bio} have been used in only a few applications, and more research is needed to assess their robustness and applicability. Metrics for biogeophysical effects, such as albedo changes, have been proposed (Betts, 2000; Rotenberg and Yakir, 2010), but as for NTCFs regional variations