

Wildfire Emission Estimates for 2022

[Errata: This report was initially published on August 24, 2023, and later updated on September 23, 2024 to (1) correct a data transcription error in Table 2, (2) correct a canopy mortality rate assumption that had led to minor differences in emissions and fuel consumption, and (3) include an additional fire (the Lost Lake Fire) that has been added to CAL FIRE's database since the initial publication of this report. These updates are described in more detail in Footnotes 1, 2, 3, and 5.]

Introduction

The California Air Resources Board (CARB) annually releases estimates of greenhouse gas (GHG) and particulate matter emissions from wildfires. This document summarizes estimates of statewide wildfire emissions from 2000 to 2022 for carbon dioxide (CO₂), particulate matter that are 10 microns or smaller in size (PM₁₀), and particulate matter that are 2.5 microns or smaller in size (PM_{2.5}). Emissions are estimated using the First Order Fire Effects Model (FOFEM) developed by the U.S. Forest Service (USFS) for fires reported in the California Department of Forestry and Fire Protection's (CAL FIRE) wildfire geodatabase using Geographic Information Systems (GIS) data on wildfire perimeters, vegetation fuels, and fuel moisture.

Fire has served natural functions in California's diverse ecosystems for millennia, such as facilitating germination of seeds for certain tree species, replenishing soil nutrients, clearing dead biomass to make room for living trees to grow, and reducing accumulation of fuel that leads to high-intensity wildfires. However, fire also impacts human health and safety and releases GHG emissions and other air pollutants, including those that contribute to ozone formation. In recent years the magnitude and intensity of wildfires have increased across California.

Overview of the 2022 Fire Season

State data reported 305 wildfires totaling approximately 0.3 million acres for 2022, comparable to total acreage reported for 2019. Total wildfire area in 2022 was approximately one-tenth the amount of wildfire area in 2021. As in previous years, a few wildfires comprised most of the year's reported burned area. Over half of the state total burn acreage was dominated by three fires: Mosquito (El Dorado and Placer counties), McKinney (Siskiyou County), and Campbell (Trinity County). The Mosquito Fire was the 77th largest fire event reported by CAL FIRE (76,739 acres), while the McKinney (60,077 acres) and Campbell (30,120 acres) fires were the 107th and 246th largest fires. Fuel loads in forested lands together with dry conditions fostered extensive fuel consumption and emissions.

Emissions Modeling Results

The estimates in this document are derived using the FOFEM developed by the USFS. Fire is a physical process that is highly variable at all spatial and temporal scales, so estimates of wildfire emissions for the entire state will have inherently high uncertainties. Sources of uncertainties include vegetation fuel types, fuel loading, fuel moisture, burned area, modeled fuel consumption in flaming and smoldering phases, and emission factors. CARB staff used GIS data on wildfire perimeters, vegetation fuels, fuel moisture, and FOFEM to account for some of this uncertainty and variation when they estimate emissions from fires reported in an interagency wildfire geodatabase (CAL FIRE 2023).

A wildfire's total emissions represent the contribution from the mosaic of vegetation types and fuels consumed within the fire footprint. Forest and woodland vegetation typically contain greater fuel loads per unit area (typically dead wood and surface fuels) than vegetation types dominated by shrubs, herbaceous plants, or grasses. Large fires extend across a variety of vegetation types. For example, the Mosquito fire extended across 28 different vegetation types, spanning forest, woodland, shrub and grass-dominated areas. However, 73% of the fire area and 77% of the total pre-burn fuel load was dominated by just three vegetation types: Douglas Fir-Sugar Pine-Tanoak Forest (FCCS 7), Jeffrey Pine-Ponderosa Pine-Douglas Fir-California Oak Woodland (FCCS 16) and California Black Oak Woodland (FCCS 14).

While there is year-to-year variation in the amount of land area affected by wildfire, the variation in total annual fire emissions is properly understood as a function of the amounts of fuel consumed. Total estimated fuel consumption (Table 1) in 2022 was less than a tenth of the fuel consumption in 2021. Total emissions of PM₁₀, PM_{2.5} and CO₂ in 2022 were correspondingly smaller in magnitude compared to 2021. The 2019 and 2022 fire seasons experienced similar total acreage burned (Figure 1), while contrasting fuel consumption accounts for their emissions differences (Figures 2, 3, and 4). In 2022, total fuel consumption was approximately twice the magnitude of total fuel consumption in 2019. In turn, emissions of CO₂ and PM₁₀ in 2022 were nearly double the magnitude of 2019's emissions.

Forest vegetation types dominated pre-fire fuel loads in 2022, with loads ranging from 0 to 138 tons/acre, averaging 19 tons/acre. The model estimated high rates of fuel consumption, with nearly complete consumption of litter, shrubs, and 1-hr, 10-hr and 100-hr dead fuels, as well as high rates for large-diameter dead fuels. Consumption of large-diameter dead fuels and duff (a forest floor dead organic layer between litter and soil layers) is largely associated with combustion in the smoldering phase. Carbon dioxide (CO₂) emissions are associated with fuel consumption in both the flaming and smoldering phases. Table 1 is a summary of 2022 wildfire area, fuel consumption, and emissions.

Table 1. Summary of 2022 wildfire area, fuel consumption, and emissions.¹

Wildfire Area (million acres)	Fuel Consumed (million short tons)	PM ₁₀ (thousand short tons)	PM _{2.5} (thousand short tons)	CO ₂ (million metric tons)
0.31	6.7	101	85	9.1

The top twenty wildfires comprised approximately 94%¹ of total area burned. Their estimated emissions are listed in Table 2. Together, the Mosquito, McKinney and Campbell fires contributed 58%¹ of the PM_{2.5} emissions estimated for 2022.

¹ Errata (September 23, 2024): Minor calculation errors were identified and corrected since the initial publication of this report on August 24, 2023. In the August 2023 version of this report, CARB staff had implemented a new, more efficient data processing pipeline using the same methodology and data sources as in prior years. However, a typo in the code inadvertently set the canopy mortality to zero instead of the default value of 39% which was used in all prior years. This resulted in PM₁₀, PM_{2.5}, and CO₂ emissions and fuel consumption values to be lower than expected. In the July 2024 update of this report, CARB staff updated the data processing code and reanalyzed 2022 wildfire areas and emissions. Besides resetting the canopy mortality rate to the default value, additional minor differences in wildfire area were observed (0.4% or 1,154 more acres across the entire state compared to the July 2023 version of this report) which are likely attributed to the updated spatial analyses required in calculating emissions. In addition, since the August 2023 publication of this report, CAL FIRE has updated their database (FRAP 2023) with an additional fire, the Lost Lake Fire, which CARB staff added to Table 2 for the July 2024 update. All numerical values in this report have been updated to reflect these corrections.

Table 2. Top 20 wildfires of 2022 by area of wildland vegetation burned²

Rank	Fire Name	Date	Area (acres)*	CO2 (million metric tons)	PM10 (thousand short tons)	PM2.5 (thousand short tons)
1	Mosquito	9/7/2022	74,919	2.0	20.1	17.0
2	Mckinney	7/29/2022	56,013	1.6	17.9	15.2
3	Campbell	8/5/2022	29,864	1.5	20.1	17.0
4	Fairview	9/6/2022	27,675	0.6	1.9	1.6
5	Oak	7/22/2022	18,620	0.5	3.0	2.5
6	Mountain	9/2/2022	13,395	0.6	9.6	8.1
7	Ammon	8/5/2022	11,306	0.5	7.3	6.2
8	Red	8/4/2022	7,941	0.4	6.3	5.3
9	Yeti	7/30/2022	7,501	0.3	4.5	3.8
10	Barnes	9/7/2022	5,575	0.1	1.8	1.6
11	Lost Lake ³	5/26/2022	5,239	0.0	0.1	0.1
12	Route	8/31/2022	5,059	0.1	0.3	0.2
13	Washburn	7/7/2022	4,812	0.2	1.8	1.5
14	Electra	7/4/2022	4,323	<0.1	0.3	0.2
15	Border 32	8/31/2022	4,168	0.1	0.3	0.3
16	Mill	9/2/2022	3,549	0.1	0.4	0.3
17	Airport	2/16/2022	3,207	<0.1	0.1	0.1
18	Rodgers	8/8/2022	2,550	0.1	1.7	1.4
19	Thunder	6/22/2022	2,398	<0.1	0.1	0.1
20	Summit	8/3/2022	1,253	0.1	1.0	0.9


*Emission estimates are associated with wildland vegetation and do not include developed areas, croplands, or water bodies, which are not included in the vegetation fuel data layer available to CARB.

² Errata (September 23, 2024): Table 2 has been updated to reflect the corrections described in Footnote 1. Additionally, in the previous version of this report published in 2023, there was a transcription error that led to PM₁₀ being reported as 1/100ths of the calculated value which has since been corrected.

³ Erratum (September 23, 2024): since the August 2023 publication of this report, CAL FIRE has updated their database (FRAP 2023) with an additional fire, the Lost Lake Fire, which CARB staff added to Table 2 for the July 2024 update.

Figures 1 to 4 present annual wildfire acreages and emissions of CO₂, PM₁₀, and PM_{2.5} for 2000 to 2022.^{4,5}

Figure 1. Acreage of Burned Wildland Vegetation Area*



*These acreages do not include areas where wildland vegetation data for model inputs are not available, e.g., developed areas and croplands.

⁴ The wildfire emissions in Figures 1 to 4 include all fire events in the CAL FIRE database (CAL FIRE 2023), including those with the "resource benefit (WFU)" tag. WFU refers to fires that are managed to accomplish specific pre-stated resource management objectives in predefined geographic areas outlined in fire management plans.

⁵ Values for 2022 in Figures 1 through 4 have been updated as per the errata in Footnote 1. Specifically, CO₂, PM₁₀, and PM_{2.5} emissions have been updated and are now slightly higher than in the July 2023 version.

Figure 2. Estimates of Wildfire CO₂ Emissions




Figure 3. Estimates of Wildfire PM₁₀ Emissions





Figure 4. Estimates of Wildfire PM_{2.5} Emissions



Data Sources and Methods

Emissions are estimated using GIS format data on fire perimeters (FRAP 2023), alarm and containment dates, natural vegetation fuel type (fuel component size class), fuel loads (tons/acre), fuel moistures, and burn severity. The geospatial data are used to develop inputs to a wildland fire emission model (FOFEM version 6.7) (FOFEM 2023). Modeled emissions in flaming and smoldering phases (lbs/acre) by fuel type are integrated over the areas of each vegetation fuel type associated with each wildfire. Flaming and smoldering emissions are summed for reporting and include every fire reported and mapped for the calendar year.

The magnitudes of emissions are proportional to the amount of fuel consumed, and various pollutants are generated in the flaming and smoldering phases of combustion. Fuel moisture influences the proportions of fuel consumed in flaming versus smoldering phases. Forest and woodland vegetation types contain greater fuel loads than vegetation types dominated by shrubs, herbaceous plants, or grasses. Large fires often extend across a variety of vegetation types. Vegetation fuel maps based on the Fuel Characteristic Classification System (FCCS) are developed for specific years by the LANDFIRE.GOV consortium (Ottmar 2007, FCCS 2022). For all other years, CARB staff use FCCS-based vegetation fuel maps developed by researchers at the University of California at Berkeley (UCB 2019). Fuel loads for FCCS vegetation types are defined in FOFEM. Fuel moistures (Abatzoglou 2013, gridMET 2022) are obtained from the Climate Engine consortium (CE 2023). Pollutant emissions associated with fuel consumption in the smoldering phase include PM₁₀ and PM_{2.5}. Emissions associated with the flaming phase include CO₂.

Uncertainty

Uncertainties associated with mapped vegetation fuel types, fuel loading (tons/acre by fuel size category) (Collins et al. 2016, McKenzie et al. 2007, Riccardi et al. 2007, Sikkink and Keane 2008), fuel moisture, burned area, modeled fuel consumption in flaming and smoldering

phases, and emission factors (EFs, mass of pollutant species per unit mass fuel consumed) contribute to large uncertainties in emission estimates reported by CARB. EFs are derived from chemical analysis of air samples during biomass burn events. Derived EFs vary with fuel type, fuel component size class, texture, arrangement, moisture content, combustion conditions (wildfire vs. prescribed burn, flaming vs smoldering, wind speed), and methods (laboratory versus field studies). For some pollutants, EF uncertainty approaches a factor of two (Urbanski 2014, Prichard et al. 2020). Fuel loading is an especially large source of uncertainty: across vegetation types and entire landscapes, fuel loading can vary by up to an order of magnitude. A 2011 study (Urbanski et al. 2011) estimated wildfire emissions across the western U.S. for 2003 through 2008 using a geospatially and temporally explicit fire emission model utilizing remotely sensed vegetation fuel, wildfire activity, and weather data. The study found that uncertainties were approximately a factor of two at spatial (kilometers) and temporal scales (daily) relevant to air quality modeling. The CARB wildfire emission estimates are developed using sources and methods that are independent from those used for the statewide Natural and Working Lands (NWL) inventory of ecosystem carbon stocks and stock-change.

References

- Abatzoglou, J. T. (2013) Development of gridded surface meteorological data for ecological applications and modelling. *International Journal of Climatology* (33), 121-131. CARB (2021) California Wildfire Emission Estimates. <https://ww2.arb.ca.gov/wildfire-emissions>
- CALFIRE 2023. Historic Fire Perimeters, 2022. Fire and Resource Assessment Program. <https://www.fire.ca.gov/Home/What-We-Do/Fire-Resource-Assessment-Program/GIS-Mapping-and-Data-Analytics>
- CE 2023. Climate Engine: Cloud Computing of Climate and Remote Sensing Data. <https://climate-engine.appspot.com/climateEngine>
- Collins, B., J. M. Lydersen, D. L. Fry, K. Wilkin, T. Moody, S. L. Stephens (2016) Variability in vegetation and surface fuels across mixed-conifer-dominated landscapes with over 40 years of natural fire. *Forest Ecology and Management* 381, 74-83. <https://doi.org/10.1016/j.foreco.2016.09.010>
- FRAP 2023. Fire perimeters: A multi-agency statewide database of fire history. Fire and Resource Assessment Program, California Department of Forestry and Fire Protection. <https://frap.fire.ca.gov/mapping/gis-data/>
- FOFEM 2023. First Order Fire Effects Model (FOFEM) <https://www.fs.usda.gov/ccrc/tools/fofem>
- FCCS 2023. Fuel Characteristic Classification Fuelbeds (FCCS) <https://www.landfire.gov/fccs.php>
- gridMET 2023. gridMET: A dataset of daily high-spatial resolution surface meteorological data covering the contiguous US from 1979 to yesterday. <http://www.climatologylab.org/gridmet.html>
- McKenzie, D., C.L. Raymond, L.-K.B. Kellogg, R.A. Norheim, A.G. Andreu, A.C. Bayard, K.E. Kopper, and E. Elman (2007) Mapping fuels at multiple scales: landscape application of the Fuel Characteristic Classification System. *Can. J. For. Res.* 37: 2421–2437 doi:10.1139/X07-056 <https://doi.org/10.1139/X07-056>
- Ottmar, R. D., D. V. Sandberg, C. L. Riccardi, and S. J. Prichard (2007) An overview of the Fuel Characteristic Classification System - Quantifying, classifying, and creating fuelbeds for resource planning. *Can. J. For. Res.-Rev. Can. Rech. For.*, 37(12), 2383-2393, doi:10.1139/x07077
- Prichard, S.J., O’Neill, S.M., Eagle, P., Andreu, A.G., Drye, B., Dubowy, J., Urbanski, S. and T.M. Strand (2020) Wildland fire emission factors in North America: synthesis of existing data, measurement needs and management applications. *International Journal of Wildland Fire* 2020, 29, 132–147. <https://doi.org/10.1071/WF19066>
- Riccardi, C., S. J. Prichard, D. V. Sandberg, and R. D. Ottmar (2007) Quantifying physical characteristics of wildland fuels using the Fuel Characteristic Classification System. *Can. J. For. Res.* 37: 2413–2420. doi:10.1139/X07-175

Sikkink, P.G., and R. E. Keane (2008) A comparison of five sampling techniques to estimate surface fuel loading in montane forests. *International Journal of Wildland Fire* 17: 363–379. doi:10.1071/WF07003 1049-8001/08/030363

UCB 2019. University of California, Berkeley (2019) Incorporating disturbance effects on fuels in the emissions estimation system. Final Report, CARB contract 15-AQP007.

Urbanski, S. (2014) Wildland fire emissions, carbon, and climate: Emission factors. *Forest Ecology and Management* 317: 51–60. <http://dx.doi.org/10.1016/j.foreco.2013.05.045>

Urbanski, S.P., Hao, W.M. and B. Nordgren (2011) The wildland fire emission inventory: western United States emission estimates and an evaluation of uncertainty. *Atmos. Chem. Phys.*, 11, 12973–13000. doi:10.5194/acp-11-12973-2011