

PART III: SYNTHESIS

SUMMARY

Wet Deposition

Wet sulfate and nitrate deposition were each no greater than $12 \text{ kg ha}^{-1} \text{ yr}^{-1}$ in all years (1985 through 1994) at all sites and were usually less than $12 \text{ kg ha}^{-1} \text{ yr}^{-1}$. Excess sulfate (i.e., excluding the estimated sea-salt contribution) was no greater than $10 \text{ kg ha}^{-1} \text{ yr}^{-1}$ in all years at all sites and was usually less than $4 \text{ kg ha}^{-1} \text{ yr}^{-1}$. Ammonium deposition was less than $5 \text{ kg ha}^{-1} \text{ yr}^{-1}$ in all years at all sites. For comparison, wet sulfate and nitrate deposition in portions of eastern North America exceed 25 and $15 \text{ kg ha}^{-1} \text{ yr}^{-1}$, respectively (Sisterson, 1991); ammonium deposition is less than about $4 \text{ kg ha}^{-1} \text{ yr}^{-1}$ in almost all parts of eastern North America (Sisterson, 1991).

In some areas where sulfate deposition is highest, such as the northwest coast, much of the sulfate had its origin as sea salt.

In most years, wet nitrate deposition was greater in the SoCAB and the southern Sierra Nevada than in other parts of California.

Deposition uncertainties are less than 20 percent in the SoCAB, which has a large number of monitors; uncertainties can be up to 100 percent in portions of northeastern and southeastern California, where little monitoring has been done.

To facilitate comparison with dry deposition estimates, Table 14 summarizes wet deposition rates for sulfur and nitrogen (as S and N).

Table 14. Multi-year mean annual wet deposition of nitrate, sulfate (not adjusted for sea salt), excess sulfate (adjusted for sea salt), and ammonium by site (kg S or N ha⁻¹ yr⁻¹). Years were included only if CI1 and CI3 were at least 75 percent. CADMP and UCSB sites are capitalized. Missing values are shown with periods.

Site	Network	No. yrs.	Species			
			Excess Sulfur	Sulfur	Nitrogen from NO ₃ ⁻	Nitrogen from NH ₄ ⁺
ALPINE MDWS	UCSB	3	0.60	0.56	0.76	0.71
ANAHEIM	CADMP	9	0.63	0.49	0.43	0.66
ANGORA LK	UCSB	3	0.82	0.76	0.84	0.96
BAKERSFIELD	CADMP	10	0.57	0.55	0.41	0.89
BERKELEY	CADMP	8	1.01	0.71	0.57	0.52
BETHEL ISL	CADMP	10	0.41	0.36	0.42	0.98
Bishop	NADP	1	0.35	0.34	0.16	0.21
Chuchupate	NADP	7	0.41	0.38	0.40	0.24
Davis	NADP	10	0.72	0.64	0.74	1.59
EASTRN BRK LK	UCSB	3	0.36	0.35	0.40	0.39
EL MONTE	CADMP	10	1.09	0.94	0.83	1.22
EMERALD LK	UCSB	3	0.75	0.70	0.75	0.93
ESCONDIDO	CADMP	10	0.77	0.50	0.46	0.51
EUREKA	CADMP	4	1.64	0.68	0.30	0.36
GASQUET	CADMP	9	2.16	0.94	0.49	0.45
Hopland	NADP	14	0.79	0.55	0.49	0.37
Hopland Intercomp	NADP	1	0.52	0.30	0.41	0.27
KAISER PASS	UCSB	2	0.59	0.56	0.58	0.65
LAKE ISABELLA	CADMP	7	0.27	0.25	0.31	0.26
LAKEPORT	CADMP	4	0.56	0.40	0.49	0.74
LINDCOVE	CADMP	3	0.53	0.49	0.88	1.52
LOS ANGELES	CADMP	4	1.34	1.13	0.76	1.45
LYNWOOD	CADMP	10	1.09	0.92	0.54	0.82
MAMMOTH MTN	CADMP	1	0.68	0.65	0.64	0.69
MINERAL KING	UCSB	3	0.56	0.54	0.74	0.92
MMTH MT/UC	UCSB	3	0.75	0.72	0.75	0.82
MONTAGUE	CADMP	9	0.26	0.25	0.31	0.35
MONTAGUE CL	CADMP	4	0.30	0.29	0.35	0.43
MT WILSON	CADMP	10	0.83	0.73	0.88	0.68
Montague	NADP	9	0.24	0.23	0.28	0.27

Table 14, continued

Site	Network	No. yrs.	Species			Nitrogen from NH ₄ ⁺
			Excess Sulfur	Sulfur	Nitrogen from NO ₃ ⁻	
NAPA	CADMP	9	1.18	0.93	0.74	0.86
NIPOMO	CADMP	4	0.57	0.37	0.31	0.39
ONION VLLY	UCSB	3	0.46	0.44	0.53	0.55
Organ Pipe Cactus	NADP	13	0.71	0.64	0.40	0.37
PASADENA	CADMP	10	1.26	1.08	1.15	1.18
Palomar Mountain	NADP	4	1.81	1.43	1.16	0.79
QUINCY	CADMP	7	0.63	0.57	0.66	0.54
RESEDA	CADMP	6	0.77	0.67	0.80	0.72
Re Rock Canyon	NADP	6	0.46	0.45	0.56	0.37
SACRAMENTO	CADMP	10	0.73	0.65	0.79	1.80
SACRAMENTO C	CADMP	4	0.65	0.59	0.71	1.91
SALINAS	CADMP	4	0.45	0.28	0.27	0.51
SAN BERNARD	CADMP	10	0.82	0.68	1.03	1.98
SAN JOSE	CADMP	10	0.55	0.40	0.30	0.59
SAN NICOLAS	CADMP	4	0.65	0.20	0.17	0.12
SAN RAFAEL	CADMP	3	1.67	1.22	0.87	1.07
SANTA BARB	CADMP	10	0.70	0.44	0.57	0.34
SEQUOIA ASH	CADMP	10	0.66	0.60	1.12	1.58
SEQUOIA GF	CADMP	9	0.79	0.73	1.14	1.63
SEQUOIA GF C	CADMP	2	0.53	0.49	1.03	1.52
SL	UCSB	3	0.43	0.42	0.47	0.40
SODA SPRINGS	CADMP	9	0.92	0.83	1.01	0.81
SOUTH L TAH	CADMP	10	0.36	0.33	0.39	0.33
SP	UCSB	3	0.55	0.52	0.53	0.52
Sequoia NP-Giant	NADP	10	1.12	1.05	1.11	1.42
Silver Lake Rgr Stn	NADP	7	0.24	0.22	0.24	0.15
Smith Valley	NADP	7	2.33	0.20	0.13	0.75
TANBARK	CADMP	10	0.95	0.81	1.18	0.99
TANBARK COL	CADMP	4	0.80	0.66	0.92	0.86
TP	UCSB	3	0.71	0.67	0.75	0.77
Tanbark Flat	NADP	11	1.25	1.07	1.30	0.91
VICTORVILLE	CADMP	4	0.36	0.34	0.49	0.51
YOSEMITE	CADMP	8	0.60	0.56	0.83	0.86
Yosemite Ntnl Park	NADP	5	1.14	1.06	1.05	0.96

Dry Deposition

This project has produced estimates of dry-deposition fluxes at 10 sites in California. These estimates will help improve our understanding of the magnitude of dry deposition. However, it is important to recognize that the calculations are limited in numerous important respects and that they could likely be improved over time with additional effort. The dry-deposition flux estimates are subject to uncertainties of approximately 50 percent.

Estimated deposition of HNO_3 at the 10 sites ranges from 1 to 86 $\text{kg ha}^{-1} \text{ yr}^{-1}$. At the urban sites, HNO_3 deposition accounts for about 30 to 80 percent of the deposition of oxidized nitrogen species and 20 to 70 percent of the total nitrogen deposition.

To facilitate comparison with wet deposition fluxes, Figure 25 shows estimates for total sulfur and nitrogen dry deposition expressed as sulfur (S) and nitrogen (N). At most sites, the annual rate of deposition of HNO_3 exceeds the rates of all other species (Fremont, Gasquet, and Long Beach are the exceptions). Expressed in terms of N (or on a molar basis), the rates of particulate ammonium deposition (not shown) are generally comparable to those of particulate nitrate deposition. The annual rates of deposition of oxidized nitrogen species at the three rural sites are about one-tenth to one-half as great as the values reported by Meyers et al. (1991), which ranged from 1.5 to 4.6 kg N ha^{-1} for sites in the eastern United States. However, the deposition rates calculated for the rural CADMP sites are quite uncertain because many of the measurements were below the limits of quantification. The deposition rates at Azusa, Bakersfield, Long Beach, and Los Angeles exceed those reported by Meyers et al. (1991) by factors of 2 to 17.

**Annual Dry Deposition Rates by Species
CADMP Sites, 1988-93**




Figure 25. Annual dry deposition rates for sulfate, nitrate, SO₂, NH₃, NO₂, and HNO₃ at 11 CADMP sites, averaged over the period 1988-93. The rates are expressed as kg N or S ha⁻¹ yr⁻¹.

Comparison of Wet and Dry Deposition

Six of the 10 dry-deposition sites are collocated with wet-deposition monitors. The sites at Azusa, Fremont, downtown Los Angeles, and Long Beach are not collocated with wet-deposition monitors. For purposes of comparison, we paired these four sites with the precipitation sites at El Monte, San Jose, Pasadena, and Lynwood. Of the 10 locations with paired wet- and dry-deposition data, three are nonurban (Gasquet, Yosemite, and Sequoia).

At the three nonurban sites (Gasquet, Yosemite, and Sequoia), wet nitrate and nonseasalt-sulfate deposition approximately equalled (within $0.7 \text{ kg ha}^{-1} \text{ yr}^{-1}$) dry deposition of oxidized nitrogen and sulfur species (see Figure 26). The multi-year average sum of wet and dry deposition at these three sites were $1\text{-}2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (from oxidized species), $1\text{-}2 \text{ kg S ha}^{-1} \text{ yr}^{-1}$, and $1\text{-}2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (from reduced species).

At the seven urban sites, dry sulfur deposition [SO_2 and particulate sulfate (pSO_4^{2-})] was approximately 1 to 3 times the magnitude of wet sulfur deposition; dry deposition of oxidized nitrogen species [HNO_3 , NO_2 , and particulate nitrate (pNO_3^-)] at the urban sites ranged from about 10 to 30 times the magnitude of wet nitrate deposition (see Figure 27). At all sites, dry deposition of reduced nitrogen species [ammonia (NH_3) and particulate ammonium (pNH_4^+)] was about a factor of 2 greater than wet ammonium deposition. The multi-year average sum of wet and dry deposition at these seven sites were $5\text{-}30 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (from oxidized species), $1\text{-}3 \text{ kg S ha}^{-1} \text{ yr}^{-1}$, and $1\text{-}4 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (from reduced species).

Comparison of Deposition and Emissions Estimates

In this section, we compare our dry-deposition flux estimates to emission estimates. Emission densities of NO_x (as $\text{kg ha}^{-1} \text{ yr}^{-1} \text{ N}$) were obtained from California Air Resources Board, Technical Support Division (1990) and converted to units of $\text{kg N ha}^{-1} \text{ yr}^{-1}$. Although several pieces of evidence suggest that emissions of volatile organic compounds (VOCs) may typically be underestimated by a factor of 2 (e.g., Fujita et al., 1992), NO_x emission estimates are generally thought to be more accurate. The emission estimates were available for either counties or air basins. We delineated areas of approximately equal sizes, which ranged in size from one to nine counties. For each site, these areas were:

- Azusa: South Coast Air Basin
- Fremont: San Francisco Bay Area Air Basin
- Los Angeles: South Coast Air Basin
- Long Beach: South Coast Air Basin
- Bakersfield: Kern County
- Sacramento: Sacramento County plus six neighboring counties (Contra Costa, Sacramento Valley portion of Placer, Solano, San Joaquin, Sutter, and Yolo).
- Santa Barbara: South Central Coast Air Basin

Wet and Dry Deposition of Oxidized Sulfur and Nitrogen, Rural CADMP Sites




Figure 26. Wet and dry deposition of oxidized sulfur and nitrogen species at 3 rural CADMP sites. The averaging periods were 1985-94 for wet deposition and 1988-93 for dry deposition. All rates are expressed as $\text{kg N or S ha}^{-1} \text{ yr}^{-1}$.

Wet Deposition, Dry Deposition, and Emissions of Oxidized Nitrogen Species




Figure 27. Wet deposition, dry deposition, and emission rates of oxidized nitrogen species at 6 urban CADMP sites. The averaging periods were 1985-94 for wet deposition and 1988-93 for dry deposition. Emission rates are for 1990. All rates are expressed as $\text{kg N ha}^{-1} \text{ yr}^{-1}$.

As shown in Figure 27, the calculated rates of deposition of oxidized nitrogen species at the South Coast Air Basin (SoCAB) stations range from 16 to 37 percent of rate of emissions of NO_x within the SoCAB. The sum of wet plus dry nitrogen deposition rates at Fremont is about 11 percent of the NO_x emissions rate occurring within the San Francisco Bay area. However, since Fremont does not experience the highest peak ozone concentrations in the Bay area, concentrations of photochemical reaction products (including HNO_3) may be greater at other locations within the Bay area, implying that deposition rates in parts of the Bay area may also be greater than those calculated for Fremont. The estimated nitrogen deposition rates at Bakersfield and Sacramento are about 76 and 32 percent, respectively, of the emissions rates of Kern County and the area around Sacramento County. Transport of NO_x from upwind areas could account in part for the relatively large ratio of deposition to emissions at Bakersfield.

LIMITATIONS

Wet-deposition flux estimates are based on data obtained using a proven monitoring technique and a reasonably dense network of stations. The most significant source of potential bias is underestimation of precipitation amounts in alpine regions. The uncertainties in our regionalized estimates of wet deposition vary spatially and among chemical species; they are typically in the range of 20 to 50 percent for the species and areas of greatest interest. In contrast, both the measurements and the model used to calculate dry deposition are subject to potentially large uncertainties. At present, questions remain regarding the accuracy of the denuder difference HNO_3 concentrations. Moreover, the expected uncertainties in dry deposition flux estimates calculated according to the inferential method are on the order of 50 percent.

RECOMMENDATIONS

We offer the following recommendations for consideration:

1. Particular effort should be devoted to resolving the questions pertaining to accurate measurement of nitric acid. At many locations, it is the largest component of total nitrogen deposition. Therefore, accurate measurement is critical.
2. Comparison of results obtained from application of the inferential method and from micrometeorological studies would be highly desirable. Lacking such a comparison, we cannot evaluate the accuracies of the calculated deposition amounts.
3. If analyses of dry-deposition trends are of interest, they should be carried out for the ambient air concentrations, rather than the calculated dry-deposition fluxes, because many uncertainties are introduced in the process of calculating fluxes. For the urban locations, where dry deposition fluxes are many times the magnitude of wet deposition, trends in dry deposition may serve as a surrogate for trends in total deposition. At rural locations, where wet and dry deposition fluxes are of comparable magnitudes, consideration should be given to co-analyzing the time trends in wet and dry deposition.

REFERENCES

- D. S. Bigelow. Quality Assurance Report NADP/NTN Deposition Monitoring: Field Operations, July 1978 Through December 1983. Ft. Collins CO: National Atmospheric Deposition Program, 1986.
- D. S. Bigelow, D. L. Sisterson, and L. J. Schroder. 1989. An interpretation of differences between field and laboratory pH values reported by the National Atmospheric Deposition Program/National Trends Network monitoring program. Environ. Sci. Technol. 23: 881-887.
- C. L. Blanchard and K. A. Tonnesen. 1993. Precipitation-chemistry measurements from the California Acid Deposition Monitoring Program, 1985-1990. Atmos. Environ. 27A(11): 1755-1763.
- C. L. Blanchard and H. M. Michaels. Regional Estimates of Acid Deposition Fluxes in California. Contract number A132-149. Sacramento, CA: California Air Resources Board, March 1994.
- California Air Resources Board (Technical Support Division). Emission Inventory, 1987. Sacramento CA: California Air Resources Board, March 1990.
- R. M. Cionco. 1972. A wind-profile index for canopy flow. Boundary Layer Meteorology. 3: 255-263.
- R. M. Cionco. 1978. Analysis of canopy index values for various canopy densities. Boundary Layer Meteorology. 15: 81-93.
- J. F. Clarke, E. S. Edgerton, and R. P. Boksleitner. 1992. Routine estimation and reporting of dry deposition for the U.S.A. dry deposition network. U.S. Environmental Protection Agency, Research Triangle Park, NC 27711.
- A. H. El-Shaarawi. 1989. Inferences about the mean from censored water quality data. Water Resources Research 25: 686-690.
- E. Englund and A. Sparks. GEO-EAS 1.2.1: Geostatistical Environmental Assessment Software, User's Guide. EPA 600/8-91/008. Las Vegas, NV: U.S. Environmental Protection Agency, 1991.
- E. M. Fujita, B. E. Croes, C. L. Bennett, D. R. Lawson, F. W. Lurmann, and H. H. Main. 1992. Comparison of emission inventory and ambient concentration ratios of CO, NMOG, and NO_x in California's South Coast Air Basin. J. Air Waste Manag. Assoc. 42: 264-276.

- R. J. Gilliom and D. R. Helsel. 1986. Estimation of distributional parameters for censored trace level water quality data. 1. Estimation techniques. Water Resources Research 22(2): 135-146.
- K. Guertin, J. Villeneuve, and S. Deschenes. 1988. The choice of working variables in the geostatistical estimation of the spatial distribution of ion concentration from acid precipitation. Atmos. Environ. 22(12): 2787-2801.
- T. C. Haas. 1990. Kriging and automated variogram modeling within a moving window. Atmos. Environ. 24A(7): 1759-1769.
- T. C. Haas. 1992. Redesigning continental-scale networks. Atmos. Environ. 26A(18): 3323-3333.
- B. B. Hicks, D. D. Baldocchi, T. P. Meyers, R. P. Hosker, Jr., and D. R. Matt. 1987. A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities. Wat. Air Soil Pollut. 36:311-330.
- B. B. Hicks, R. P. Hosker, Jr., T. P. Meyers, and J. D. Womack. 1991. Dry deposition inferential measurement techniques - I. Design and tests of a prototype meteorological and chemical system for determining dry deposition. Atmos. Environ. 25A(10): 2345-2359.
- J. A. Horrocks and J. D. Kowalski. 1987. A computerized quality control method for the validation of wet deposition laboratory analysis. Paper no. 87-91.2. American Waste Management Association Annual Meeting.
- A. G. Journel and C. Huijbregts. Mining Geostatistics. New York NY: Academic Press, 1978. 600 pp.
- J. M. Lockard. Quality Assurance Report NADP/NTN Deposition Monitoring: Laboratory Operations, July 1978 Through December 1983. Ft. Collins, CO: National Atmospheric Deposition Program, 1987.
- R. T. McMillen. 1990. Estimating the spatial variability of trace gas deposition velocities. National Oceanic and Atmospheric Administration Technical Memorandum ERL ARL-181.
- J. M. Melack, J. O. Sickman, F. Setaro, and D. Dawson. Monitoring of Wet Deposition in Alpine Areas in the Sierra Nevada. California Air Resources Board Contract A932-081. Santa Barbara, CA: Marine Science Institute, University of California, 16 January 1995.
- T. P. Meyers and T. S. Yuen. 1987. An assessment of averaging strategies associated with

day/night sampling of dry-deposition fluxes of SO₂ and O₃. J. Geophys. Res. 92(D6): 6705-6712.

G. P. Meyers, B. B. Hicks, R. P. Hosker, Jr., J. D. Womack, and L. C. Satterfield. 1991. Dry deposition inferential measurement techniques - II. Seasonal and annual deposition rates of sulfur and nitrogen. Atmos. Environ. 25A(10): 2361-2370.

G. W. Oehlert. 1993. Regional trends in sulfate wet deposition. JASA 88(422): 390-399.

J. L. Peden. Quality Assurance Report NADP/NTN Deposition Monitoring: Laboratory Operations, January 1984 Through December 1985. Ft. Collins CO: National Atmospheric Deposition Program, 1988.

R. B. See, L. J. Schroder, and T. C. Willoughby. External Quality-Assurance Results for the National Atmospheric Deposition Program and the National Trends Network During 1987. Water-Resources Investigations report 89-4015. Denver, CO: U.S. Geological Survey, 1989.

S. K. Seilkop and P. L. Finkelstein. 1987. Acid precipitation patterns and trends in eastern North America, 1980 - 84. J. Climate and App. Meteor. 26: 980-994.

J. Seinfeld. Atmospheric Chemistry and Physics of Air Pollution. New York: John Wiley & Sons, 1986. 738 pp.

R. H. Shaw and A. R. Pereira. 1982. Aerodynamic roughness of a plant canopy: a numerical experiment. Agricultural Meteorology 26: 51-65

A. Sirois. 1990. The effects of missing data on the calculation of precipitation-weighted-mean concentrations in wet deposition. Atmospheric Environment 24A(9): 2277-2288.


D. L. Sisterson. Deposition monitoring: methods and results. In: Acidic Deposition: State of Science and Technology, Summary Report of the U.S. National Acid Precipitation Assessment Program. P. M. Irving, ed. Washington DC: National Acid Precipitation Assessment Program, 1991.


P. A. Solomon, T. Fall, L. Salmon, P. Lin, F. Vasquez, and G. R. Cass. Acquisition of Acid Vapor and Aerosol Concentration Data for Use in Dry Deposition Studies in the South Coast Air Basin. ARB Research Contract No. A4-144-32. Sacramento CA: California Air Resources Board, 1988.

W. Stumm and J. J. Morgan. Aquatic Chemistry: An Introduction Emphasizing Chemical Equilibria in Natural Waters, 2nd ed. New York NY: John Wiley & Sons, 1981. 780 pp.

- B. K. Takemoto, B. Cahill, C. Buenviaje, R. Abangan, and T. Houston. California Acid Deposition Data Summary -- July 1987 to June 1990, Hydrological Years 1988, 1989 and 1990. Sacramento CA: California Air Resources Board, February 1995a.
- B. K. Takemoto, B. Cahill, C. Buenviaje, R. Abangan, and T. Houston. California Acid Deposition Data Summary -- July 1990 to June 1993, Hydrological Years 1991, 1992 and 1993. Sacramento CA: California Air Resources Board, July 1995b.
- E. C. Tuazon, C. L. Blanchard, S. J. Hering, D. Lucas, and G. I. Mackay. A Review of Nitric Acid Measurements by Tunable Diode Laser Absorption Spectroscopy (TDLAS). Contract number 93-340. Sacramento CA: California Air Resources Board, December 1995.
- J. G. Watson, J. C. Chow, R. T. Egami, J. L. Bowen, C. A. Frazier, A. W. Gertler, D. H. Lowenthal, and K. Fung. Dry Deposition measurements for the California Acid Deposition Monitoring Program. Contract number A6-076-32. Sacramento, CA: California Air Resources Board, 1991.
- M. L. Wesely. 1989. Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models. Atmos. Environ. 23: 1293-1304.

**APPENDIX A: DEPOSITION AND UNCERTAINTY MAPS FOR
PRECIPITATION AND PRECIPITATION CHEMISTRY**







July 1990 - June 1991

Precipitation C.V.(%)









July 1991 - June 1992

Precipitation C.V.(%)










July 1992 - June 1993

Precipitation C.V.(%)










July 1993 - June 1994


Precipitation C.V.(%)


















July 1992 - June 1993


Acidity C.V.(%)








July 1993 - June 1994

Acidity (g/ha)










July 1990 - June 1991

Ammonium C.V.(%)






July 1991 - June 1992


Ammonium (kg/ha)







July 1991 - June 1992

Ammonium C.V.(%)










July 1992 - June 1993

Ammonium C.V.(%)








July 1993 - June 1994


Ammonium C.V.(%)





July 1990 - June 1991


Nitrate (kg/ha)





July 1990 - June 1991


Nitrate C.V.(%)





July 1991 - June 1992


Nitrate (kg/ha)






July 1991 - June 1992

Nitrate C.V.(%)









July 1992 - June 1993

Nitrate C.V.(%)








July 1993 - June 1994

Nitrate C.V.(%)

0-20
60-80

20-40
80-100


40-60






July 1990 - June 1991

Sulfate C.V.(%)











July 1991 - June 1992

Sulfate C.V.(%)












July 1993 - June 1994


Sulfate C.V.(%)





July 1990 - June 1991


Excess Sulfate (kg/ha)






July 1990 - June 1991

Excess Sulfate C.V.(%)









July 1991 - June 1992

Excess Sulfate C.V.(%)








July 1992 - June 1993


Excess Sulfate C.V.(%)





July 1993 - June 1994


Excess Sulfate (kg/ha)





July 1993 - June 1994


Excess Sulfate C.V.(%)




APPENDIX B: TIME SERIES PLOTS OF MEAN PRECIPITATION AMOUNT AND WET DEPOSITION FLUXES FOR EACH AIR BASIN

The figures shown in this appendix are time series plots of precipitation amount and wet deposition fluxes for each air basin, derived from the maps shown in Appendix A and from corresponding maps from earlier water years (Blanchard and Michaels, 1994). The minimum, mean, and maximum grid-cell averages are shown for each year.


AIRBASIN = Great Basin Valleys




AIRBASIN = Lake County




AIRBASIN = Lake Tahoe




AIRBASIN = Mountain Counties




AIRBASIN=North Central Coast




AIRBASIN = North Coast




AIRBASIN = Northeast Plateau




AIRBASIN = Sacramento




AIRBASIN=San Diego




AIRBASIN = San Francisco




AIRBASIN = San Joaquin




AIRBASIN=South Central Coast




AIRBASIN = South Coast




AIRBASIN - South East Desert




AIRBASIN = Great Basin Valleys




AIRBASIN = Lake County




AIRBASIN = Lake Tahoe




AIRBASIN = Mountain Counties




AIRBASIN=North Central Coast




AIRBASIN=North Coast




AIRBASIN = Northeast Plateau




AIRBASIN = Sacramento




AIRBASIN=San Diego




AIRBASIN = San Francisco




AIRBASIN = San Joaquin




AIRBASIN = South Central Coast




AIRBASIN - South Coast




AIRBASIN = South East Desert




AIRBASIN=Great Basin Valleys




AIRBASIN = Lake County




AIRBASIN = Lake Tahoe




AIRBASIN = Mountain Counties




AIRBASIN = North Central Coast




AIRBASIN = North Coast




AIRBASIN = Northeast Plateau




AIRBASIN = Sacramento




AIRBASIN = San Diego




AIRBASIN = San Francisco




AIRBASIN = San Joaquin




AIRBASIN = South Central Coast




AIRBASIN=South Coast




AIRBASIN=South East Desert




AIRBASIN = Great Basin Valleys




AIRBASIN = Lake County




AIRBASIN = Lake Tahoe




AIRBASIN = Mountain Counties




AIRBASIN = North Central Coast




AIRBASIN = North Coast




AIRBASIN = Northeast Plateau




AIRBASIN = Sacramento




AIRBASIN = San Diego




AIRBASIN = San Francisco




AIRBASIN = San Joaquin




AIRBASIN=South Central Coast




AIRBASIN = South Coast




AIRBASIN=South East Desert




AIRBASIN = Great Basin Valleys




AIRBASIN = Lake County




AIRBASIN = Lake Tahoe




AIRBASIN = Mountain Counties




AIRBASIN=North Central Coast




AIRBASIN = North Coast




AIRBASIN = Northeast Plateau




AIRBASIN = Sacramento




AIRBASIN=San Diego




AIRBASIN = San Francisco




AIRBASIN = San Joaquin




AIRBASIN = South Central Coast




AIRBASIN = South Coast




AIRBASIN = South East Desert




AIRBASIN = Great Basin Valleys




AIRBASIN = Lake County




AIRBASIN = Lake Tahoe




AIRBASIN=Mountain Counties




AIRBASIN = North Central Coast




AIRBASIN = North Coast




AIRBASIN = Northeast Plateau




AIRBASIN = Sacramento




AIRBASIN=San Diego




AIRBASIN = San Francisco




AIRBASIN = San Joaquin



AIRBASIN = South Central Coast



AIRBASIN = South Coast



AIRBASIN = South East Desert

