


LIBRARY - AIR RESOURCES BOARD



Final Report

Calibration of diatom-pH-alkalinity methodology for the interpretation of the sedimentary record in Emerald Lake  
Integrated Watershed Study

by

Robert W. Holmes  
Marine Science Institute  
University of California  
Santa Barbara, California 93106

October 10, 1986

California Air Resources Board Contract A4-118-32

Prepared for California Air Resources Board

## Abstract

The present study was designed to establish quantitative relationships between lake air-equilibrated pH, alkalinity, and diatoms occurring in the surface sediments in high elevation Sierra Nevada lakes. These relationships provided the necessary information to develop predictive equations relating lake pH to the composition of surface sediment diatom assemblages in 27 study lakes. Using the Hustedt diatom pH classification system, Index B of Renberg and Hellberg, and multiple linear regression analysis, two equations were developed which predict lake pH from the relative abundance of sediment diatoms occurring in each of four diatom pH groupings. Both equations predicting pH were statistically significant and yielded values of the square of the correlation coefficient of 0.82 and 0.87, respectively. Similar analyses using alkalinity rather than pH as the dependent variable likewise proved statistically significant but with lower values of the square of the correlation coefficient (0.60 and 0.76, respectively).

## Acknowledgments

The author would like to acknowledge the help provided by Dr. John Stoddard who collected the sediment cores and performed the water chemistry analyses and Dr. Mark Whiting who prepared the diatom mounts and counted and identified the diatoms in the sediment samples. Valuable assistance in diatom identification, and identification verification were obtained from PIRLA diatomists K. Camburn, Dr. D. Charles, Dr. J. Kingston, Dr. R. Kreis, and R. Sweets. Dr. C. Reimer of the Academy of Natural Sciences, Philadelphia was of great assistance in helping identify some problem taxa. Dr. R. Davis kindly provided drawings of the modified Hongve corer used to obtain surface sediment samples.

This report was submitted in fulfillment of ARB contract A4-118-32, Calibration of diatom-pH-alkalinity methodology for the interpretation of the sedimentary record in Emerald Lake, integrated watershed study, by R. W. Holmes, Marine Science Institute, University of California, Santa Barbara, California under the partial sponsorship of the California Air Resources Board. Work was completed as of August 5, 1986.

The statements and conclusions in this report are those of the contractor and not necessarily those of the California Air Resources Board. The mention of commercial products, their source or their use in connection with material reported herein is not to be construed as either actual or implied endorsement of such products.

## Table of Contents

	Page
Abstract . . . . .	1
Acknowledgments . . . . .	1
Summary and Conclusions . . . . .	3
Introduction . . . . .	5
Methods . . . . .	7
Results and Discussion . . . . .	11
References . . . . .	14
Appendix A Diatom taxon list and pH categories . . . . .	23
Appendix B Data for individual lakes . . . . .	31

## Summary and Conclusions

Thirty high elevation Sierra Nevada lakes (Fig. 1) of varying air equilibrated pH (5.82-9.53) were sampled in midsummer of 1985. In each lake a surface sediment core was taken in the deepest part of the lake with a modified Hongve (1972) corer (Fig. 2). The upper centimeter of each core was extruded in the field, placed in a plastic bag and sealed. This sample was used for the diatom analysis. A subsurface water sample was also collected and a pH measurement made immediately. Measurements of pH (air-equilibrated), alkalinity, conductivity, and major anions and cations were carried out later at UCSB or the Sierra Nevada Research Laboratory (Appendix B).

An aliquot of each sediment sample was prepared for diatom analysis by first chemically oxidizing the organic matter in the sample and then mounting a measured volume of the "cleaned" material with a high refractive index mounting medium (Hyrax) on a microscope slide. Approximately 500 diatom valves were counted and identified on each slide. One lake sediment sample contained too few diatoms and could not be used in the analysis.

The frequency of occurrence of each taxon as a function of air-equilibrated pH was examined in all lakes and subjectively placed in one of Hustedt's pH categories providing the taxon occurred in three or more lakes or obtained a relative frequency of about 1% in fewer than three lakes. A total of 408 diatom taxa were identified in the study. 173 (43%) of these are apparently undescribed entities and were given reference numbers. Fifty-five of these unknowns could be assigned to pH categories-- altogether 183 taxa were placed in Hustedt (1939) pH categories and used obtaining the diatom-pH-alkalinity predictive equations.

For published taxa, agreement between the pH assignments used in this study and those which have been assigned to pH categories in the published literature was quite good (e.g. Anderson et al. 1986, Brugam and Lwok 1986, Charles 1985, Merilainen 1967, and Stokes and Yung 1986). Since no lake in the Sierra data set had a pH below 5.5 no taxa were placed in the Acidobiontic category even though 3 taxa encountered (Anomoeoneis serians v. brachysira, Cymbella gaeumannii, and Navicula subtilissima) are generally placed in this category. A small number of taxa generally considered Indifferent were placed in the Alkaliphilic category as their Sierra pH range seemed to justify this transfer. Very few Sierra taxa fell into the Indifferent (sensu Hustedt 1939) category.

The relative abundances and pH categories were then employed lake by lake to calculate log Index B (Renberg and Hellberg 1982):

$$\text{Index B} = \frac{\% \text{ Indifferent}}{\% \text{ Indifferent}} + \frac{(5x\% \text{ Acidophils})}{(3.5x\% \text{ Alkaliphils})} + \frac{(40x\% \text{ Acidobionts})}{(108x\% \text{ Alkalibionts})}$$

A regression equation was then obtained using lake air-equilibrated pH as the dependent variable and log Index B as the independent variable. The following equation was obtained (also see Fig. 3B):

$$\text{diatom inferred pH} = 7.11 - 0.40 \log \text{Index B}$$
$$F = 117 \quad \text{Prob. } >F = 0.0001 \quad r^2 = 0.82 \quad n = 27$$

Using multiple linear regression techniques with the same data the following equation was obtained:

$$\text{diatom inferred pH} = 7.08 - 0.0086\% \text{ ACP} + 0.0012\% \text{ IND} +$$
$$0.0081\% \text{ ALP} + 0.021\% \text{ ACB}$$
$$F = 37 \quad \text{Prob. } >F = 0.0001 \quad r^2 = 0.87 \quad n = 27$$

where ACP is the abbreviation for Acidophilous, IND is Indifferent, ALP is Alkaliphilous, and ALB is Alkalibiotic.

Since the IND regression coefficient was not significant in the above regression equation, a new equation was obtained omitting it. The new equation is (also see Fig. 3A):

$$\text{diatom inferred pH} = 7.18 - 0.0097\% \text{ ACP} + 0.0070\% \text{ ALP} +$$
$$0.020\% \text{ ALB}$$
$$F = 52 \quad \text{Prob. } >F = 0.0001 \quad r^2 = 0.87 \quad n = 27$$

All three equations are significant although the  $r^2$  values are somewhat lower than those reported in the literature (e.g. Arzey 1986, Battarbee 1984, Charles 1984, 1985). The fact that the Indifferents can be dropped from regression equation without altering the  $r^2$  value is most likely due to the wide distribution above and below neutrality used in my definition of Indifferent taxa. Taxa which only occur within a limited range ( $\pm 0.5$  pH unit) around neutrality are uncommon in the data set.

This project was undertaken to provide a tool that will enable scientists to estimate pH and its change through time in Sierra lakes from diatoms preserved in lake sediments. Many of these lakes are poorly buffered and it is important to determine their pH history, particularly over the past 100 years or so. If a significant reduction in pH has taken place then it will be important to determine the cause or causes.

The statistical significance of these equations and the very acceptable  $r^2$  values strongly suggest that reasonable pH estimate can be made down core in Sierra lake sediments. In a few months a diatom pH reconstruction of a 20-cm long core from Emerald Lake, Sequoia National Park, using these equations will have been completed.

## Introduction

Ever since Oden's 1968 report (Battarbee *et al.* 1986) drew attention to the probable relationship between atmosphere acid deposition and the acidification of some Swedish lakes, the need to examine long-term pH records from lakes of different chemical composition and buffering capacity has been recognized. Unfortunately such information does not exist for many lakes. Thus scientists interested in possible relationships between acid deposition and changes in lake acidity have found it necessary to seek alternative means of obtaining pH histories of lake waters. It was quite natural for limnologists to turn to the study of lake sediments since lake sediments contain decipherable chemical and biological information about lake and watershed conditions and their changes through recent and geological time (e.g. Deevey 1942, Bradley 1966, Haworth 1969, Pennington 1943).

One of the methods which has been employed frequently in the past decade to study past pH changes in lakes employs the composition of diatom assemblages preserved in lake sediments. The method relies on the observation that the occurrence of diatom species in aquatic environments reflects, among other things, the pH of their environment. Hustedt (1939) was perhaps the first investigator to recognize such relationships. He presented a pH classification scheme which recognized 5 diatom-pH categories:

- Acidobiontic (ACB) : occurring below pH 7.0 with optimum distribution below pH 5.5
- Acidophilous (ACP) : species occurring at about 7.0 with widest distribution at pH less than 7.0
- Indifferent (IND) : equal occurrences on both sides of pH 7.0
- Alkaliphilous (ALP) : occurring at pH about 7 with widest distribution at pH greater than 7
- Alkalibiontic (ALB) : occurring at pH values greater than 7.0

It remained for Nygaard (1956) and Merilainen (1967) to develop semi-quantitative indices relating the composition of diatom assemblages observed growing attached to aquatic plants, rocks, or on the sediment, to lake pH. The indices developed by these investigators employed the diatom pH categories developed by Hustedt (1939) and the relative abundance of taxa in two or more of these pH categories. In 1982 Renberg and Hellberg developed an additional index and in 1984 Charles (1984, 1985) used multiple linear regression (MLR) methods to relate the occurrence of diatoms placed in Hustedt's pH categories to air-equilibrated lake pH. These indices and MLR have been employed in Scandinavia (Berge 1982, Renberg and Hellberg 1982, Tolonen and Jaakkola 1983), in Germany (Arzet 1986), in the United Kingdom (Flower and Battarbee 1983), in Canada (Dickman *et al.* 1983), and the United States (Brugam and Lusk 1986, Charles 1984, Davis *et al.*, 1983,

Del Prete and Schofield, 1981, Ford 1986, and Tolonen et al. 1986) to reconstruct the past pH histories of lakes.

The present study was designed to establish quantitative relationships between lake pH, and lake alkalinity and diatom occurrence in the surface sediments in high elevation lakes in the Sierra Nevada. Once such relationships have been established they can be employed to infer the pH history from sediment cores from Sierra lakes. Such diatom inferred pH reconstructions are important in the Sierra Nevada for many of these lakes are poorly buffered (Melack et al. 1985) and thus sensitive to acidic inputs, such as acid deposition.


## Methods

Thirty lakes were sampled in this study (Fig. 1). They represent, with two exceptions, a subset of the 73 Sierra Nevada lakes studied by Melack *et al.* (1985) and were selected to cover a wide range of pH (5.84 to 9.53) and alkalinity (5.9-1051  $\mu$  equiv. per l.) values (Table 1). They range in elevation from 2162 to 3609 meters (see Appendix B for additional information).

Water samples for chemical analyses were collected at about 1/2 meter depth following the collection of the sediment core (see below). The pH of the water was measured immediately and is designated pH Field in Appendix B. After the sample was returned to the laboratory usually within 8 hours air equilibrated pH (pH Lab) and alkalinity were measured. Samples for the analyses of nitrate, chloride, sulfate, and cations were prepared, stored, and analyzed using the methods described in Melack *et al.* 1985.

Surface sediment cores were taken in the deepest part of each lake with a modified Hongve (1972) corer (Fig. 2). The upper 1 cm of sediment sample was extruded in the field and placed in a plastic bag and sealed. Upon return to SNARL (The Sierra Nevada Aquatic Research Laboratory) each sample was mixed thoroughly by massaging the plastic bag containing the sediment. The bag was then opened. A plastic spoon was dipped into the homogenized sediment, withdrawn, and excess sediment removed by sliding a metal spatula along the lip of the spoon. Any sediment clinging to the sides or bottom of the spoon was carefully removed with paper tissue. The remaining sediment (1.2 cc) was quantitatively transferred with a jet of water to a 1-liter Pyrex beaker. Approximately 30 ml of 30% hydrogen peroxide was then added, and the beaker covered with parafilm. The beaker was then placed in an oven at 50°C and left for 24-48 hrs. The oxidation of the sample was completed by adding approximately 1 cc of potassium dichromate after placing the beaker in a fume hood. Following the completion of the reaction, the beaker was filled with deionized water, covered with parafilm, and set aside for 48 hrs (or more) to allow the sediment to settle. The clear supernatant was carefully decanted and the remaining liquid and sediment quantitatively transferred to a clean 250-ml beaker which was then filled with deionized water. After 48 hrs the supernatant was clear and was siphoned off. The sediment was next transferred to a 6-dram vial and filled with deionized water. Once in the vial and with a reduced dichromate concentration, the settling time of the sediment increased greatly and centrifugation and resuspension of the sediment with deionized water was employed until the supernatant was colorless.


The preparation of permanent diatom mounts was made following the methods of Battarbee (1973). The cleaned material in the vial was mixed by vigorously shaking the vial for 1 minute. The vial was then placed in a vertical position for 30



**Figure 1.** Approximate locations of lakes from which surface sediment samples we collected for the diatom/pH calibration.

Table 1. 1985 Sierra Nevada lake survey. Lake location, in situ pH, air equilibrated pH, and alkalinity ( u equiv./l.).  
For additional information see Appendix B.

No.	Lake Name	Latitude	Longitude	i.s. pH	a.e. pH	Alkalinity
1	Parker Pass	37 50 08	119 12 35	5.64	5.84	5.9
2	Table Meadow	36 36 33	118 39 05	6.6	6.01	40.8
3	Fairy Shrimp	37 26 40	118 45 40	6.92	6.3	14.6
4	Kuna	37 49 50	119 14 15	7.02	6.25	19.1
5	Dana	37 54 35	119 13 07	6.65	6.32	19.3
6	Mosquito 3	36 25 09	118 37 16	6.52	6.41	51.2
7	Mosquito 5	36 24 53	118 37 35	6.87	6.42	52.8
8	Lower Cathedral	37 50 21	119 24 50	6.45	6.44	24.6
9	Upper Treasure	37 23 13	118 46 00	6.89	6.45	27.2
10	Dade	37 22 47	118 45 42	6.8	6.48	24
11	Le Conte	38 52 14	120 08 11	6.25	6.55	31.1
12	Summit	37 26 00	118 46 11	7.1	6.6	21.4
13	Mosquito 1	36 25 28	118 37 08	7.13	6.79	77.4
14	Gem	37 23 50	118 45 20	7.33	6.82	47.6
15	Granite	36 51 47	118 37 12	6.91	6.91	66.9
16	Heather	38 52 36	120 08 16	6.95	6.93	67.1
17	Ruby	37 24 50	118 46 15	7.05	6.95	47.3
18	Upper Angora	38 51 46	120 03 59	7	6.98	88.5
19	Dorothy	37 32 10	118 52 55	7.46	7.2	138.5
20	Eastern Brook	37 25 51	118 44 28	7.35	7.22	132.2
21	Upper Gaylor	37 55 20	119 16 01	7.37	7.27	102.4
22	Lundy	38 01 55	119 13 10	7.6	7.41	260.5
23	Constance	37 30 55	118 52 02	7.95	7.42	271.5
24	Upper Frog	38 02 38	119 17 20	7.85	7.43	257.6
25	Wit-so-nah-pah	37 31 32	118 52 32	8.15	7.57	228.7
26	East Twin	38 09 40	119 20 00	8.14	7.66	383.6
27	West Twin	38 08 51	119 21 40	7.96	7.8	324.4
28	Convict	37 35 30	118 51 25	8.4	8.35	1051
29	Bright Dot	37 32 40	118 51 40	9.07	9.03	627
30	Barney	37 33 47	118 58 02	8.77	9.53	392



**FIGURE 2:**  
Sketch of the modified  
Hongve gravity corer  
(inside diameter of  
core tube = 2.5 cm.).

seconds to allow heavy material and sand to settle. An aliquot usually 250  $\mu$ l ) was then removed and placed in a flask. Following vigorous agitation of the flask, its contents were poured into a Battarbee chamber containing cover glasses. The slide warmer on which the chambers had been placed originally was then turned on and maintained at 36°-37°C until the water in the Battarbee chamber evaporated. The coverslips containing the sedimented material were removed and placed on a hot plate set at about 80°C to drive off any residual water. Hyrax, a high refractive index mountant, was placed on each of the coverslips on the hot plate. After the mountant solvent evaporated, the cover glasses were transferred to microscope slides. The slides were gently heated with a micro Bunsen burner until the mountant melted and flowed out to the edges of the cover glasses. Frequently one, occasionally two, of the cover glasses exhibited a clumped distribution of particulates following the evaporation of the aqueous suspension in a Battarbee chamber. These were not used in the subsequent quantitative diatom enumeration.

For dry weight and ignition loss determinations another 1.2 cc of sediment was removed from each plastic sample bag as described above. Each sample was placed in a tared-aluminum pan, dried at 80°C to a constant weight, and then combusted at 485°C to obtain an estimate of organic matter content.

Approximately 500 diatom valves were identified and counted in each lake sediment sample using an oil immersion 100x objective on a Leitz Ortholux compound microscope equipped with differential interference contrast optics. Diatoms were counted in one or more transects across the coverglass until the required number of valves had been counted.

Diatom identifications were made using standard accepted reference works (Cleve-Euler, 1951-1955, Germain, Hustedt 1927-66, 1930, Patrick and Reimer 1966, 1975) supplemented by numerous more specialized publications. The unpublished but widely distributed PIRLA (Paleoecological Investigations of Recent Lake Acidification) Diatom Iconograph compiled and prepared by Keith Camburn with contributions from other PIRLA diatomists proved very useful. This series of photographic plates of diatom valves was prepared under an Electric Power Research Institute contract to P. Whitehead and D. Charles, Biology Department, Indiana University.

Verification of our diatom identifications (i.e. quality assurance) was achieved in several ways. First two PIRLA diatom identification workshops were attended by the PI. Photographs and prepared material of Sierra diatoms were examined by several PIRLA diatomists who commented upon our identifications. The PI of this study also spent a week with PIRLA diatomist K. Camburn who examined additional material and photographs offering additional comments on our naming of Sierra taxa. Thus our

assignment of names to diatoms is consistent with PIRLA identifications. Lastly, a two-week visit to the Diatom Herbarium of the Academy of Natural Sciences, Philadelphia permitted an examination of their extensive collections and helped resolve identification problems with some remaining taxa. In spite of these intensive efforts 173 taxa (species and varieties) out of 408 "taxa" identified in the Sierra samples could not be given scientific names. They were of necessity given SN (Sierra Nevada) numbers (Appendix A). Fifty-seven of these SN taxa could be assigned to pH preference categories (see below). An additional 16 were assigned PIRLA numbers.

The pH "preference categories" assignment was based on a modification of the Hustedt (1939) categories. Hustedt (1939) pH categories were defined as follows:

Acidobiontic (ACB)	- optimum distribution at pH below 5.5 (occur only in acidic habitats)
Acidophilic (ACP)	- widest distribution at pH less than 7
Indifferent (IND)	- distributed around pH 7
Alkaliphilic (ALP)	- widest distribution at pH greater than 7
Alkalibiontic (ALB)	- occur only at pH greater than 7

Assignment of the Sierra taxa to these pH categories was based on the abundance weighted means (AWM) and frequency of occurrence of each taxon in every lake in which the taxon occurred. A taxon was only placed in a pH category if it occurred in 3 or more lakes or if its relative abundance exceeded about 1% in 3 or fewer lakes. A total of 183 taxa were placed in pH categories and used in this analysis. A taxon was considered acidophilic if its AWM  $\leq 7.0$  even if it occurred in low abundance at pH's between about pH 7.5 and 7.0. Alkaliphilous species had an AWM between pH 7.0-8.0 and even if they occurred occasionally in low abundance between pH 8.5 and 6.5. Alkalibiontic taxa were those with AWM  $> 8.0$ . In the Sierra material the greatest difficulty was experienced in assignment of taxa to the Indifferent group. Very few taxa had AWM close to pH 7.0 (designated as CN in Appendix A) with distributions limited between pH 6.5 and 7.5. In the present study broader limits were used (pH 6.2-7.8) with allowance for low relative abundance beyond these limits. Such taxa are truly Indifferent to pH. Had more restricted criteria been employed calculation of Index B (see below) would have been impossible since the denominator in this equation would have equalled 0 in about 1/3 of the study lakes. The absence of alkaliphilous species in about 1/2 of the study lakes precluded the calculation of Index Alpha (Nygaard 1956) without artificially setting the denominator equal to 1.0 (see Charles 1985).

The assignment of Sierra taxa to the somewhat modified ACP, ALP, ALB, and IND categories of Hustedt (1939) agrees quite well to similar assignments of the same taxa in other geographic areas (Anderson *et al.* 1986, Brugham and Lusk, 1986, Charles 1985,

Lowe 1974, Merilainen 1967, Stokes and Yung 1986). Differences mainly involve a few species (e.g. Anomoeoneis serians v. brachysira, Cymbella gaeumanii, and Navicula subtilissima) which are considered Acidobiontic by most authors and which were placed in the Acidophilous category in the present study since no lakes in the Sierra data set had a pH as low as 5.5. A few "Indifferent" species of several authors were considered alkaliphilous in Sierra samples because some of them had occurrences near pH 8. The effect of these classification differences had little effect upon Index B or the linear regression analyses.

Regression analyses and associated statistics were obtained using PC SAS Version 6 (SAS Institute, Cary, N.C.).

## Results and Discussion

Two diatom pH predictive equations were derived from the relative abundance of each diatom taxon assigned to the modified Hustedt (1939) pH categories in 29 of the 30 lakes sampled. The sediment of Dana Lake contained so few diatoms that quantitative counts could not be obtained. In each of the remaining lakes the relative abundance of taxa in each of the pH categories were summed and Index B (Renberg and Hellberg 1982) calculated using their equation:

$$\text{Index B} = \frac{\text{IND} + (5.0 \times \% \text{ ACP}) + (40 \times \% \text{ ACB})}{\text{IND} + (3.5 \% \text{ ALP}) + (108 \times \% \text{ ALB})}$$

When the observed pH and Log Index B were compared a good relationship was observed except for two extreme outliers which were the two highest pH lakes (Barney pH Lab 9.53, Bright Dot pH 9.03) sampled.

Multiple linear regression analyses using observed air-equilibrated pH values as the dependent variables and diatom pH categories as independent variables also revealed Bright Dot and Barney Lakes as extreme outliers. In both of these lake sediment samples 3-4 diatom taxa were highly dominant and possessed unusual and uncharacterizable pH categories which clearly degraded their pH signal. Thus data from these lakes was excluded when calculating Index B and regression equations, leaving 27 out of the 30 lakes sampled available for pH and alkalinity predictions.

The following relationship between air equilibrated pH (Lab pH) and log Index B was obtained (also see Fig. 3B):

$$\text{Lab pH} = 7.11 - 0.40 \log \text{Index B}$$
$$F = 117 \quad \text{Prob. } > F = 0.0001 \quad r^2 = 0.82 \quad n = 27 \quad (1)$$

Clearly the relationship is statistically significant and the equation can be used to estimate pH from diatoms preserved in diatom sediments (Fig. 3B).

A somewhat better relationship between the relative abundance of diatoms in pH categories in each lake was obtained using multiple linear regression. The same data yielded the following equation:

$$\text{Lab pH} = 7.08 - 0.0086\% \text{ ACP} + 0.0012\% \text{ IND} + 0.0081\%$$
$$\text{ALP} + 0.021\% \text{ ALB}$$
$$F = 37 \quad P>F = 0.0001 \quad r^2 = 0.87 \quad n = 27 \quad (2)$$

These equations are comparable to those reported in the scientific literature (e.g. Arzet *et al.* 1986, Charles 1985, Dickman *et al.* 1984, Renberg and Hellberg 1982). In equation 2 the coefficient for the IND category was not significant. This

is clearly an indication of the difficulty experienced in establishing satisfactory criteria for the Indifferent group (see Methods). A new equation (3) obtained without using the Indifferent category (3) was calculated and yielded the following (also see Fig. 3A):

$$\text{Lab pH} = 7.18 - 0.0097\% \text{ACP} + 0.0070 \text{ ALP} + 0.20 \text{ ALB}$$

$$F = 52 \quad P>F = 0.0001 \quad r^2 = 0.87 \quad n = 27 \quad (3)$$

The F value increased somewhat while the  $r^2$  remained the same, clearly showing that the Indifferent group as defined herein is of no value in predicting pH. It remains possible that a redefinition of the Indifferents might improve the overall F and  $r^2$  values.

In the Sierra Nevada lake data set significant positive correlations exist between Lab pH and alkalinity ( $F = 46.1$ , Prob.  $P > F = 0.0001$ ,  $r^2 = 0.62$ ,  $n = 30$ ) and between Lab pH and Log alkalinity ( $F = 93.5$ , Prob.  $>F = 0.0001$ ,  $r^2 = 0.77$ ,  $n = 30$  - see Figs. 4A and 4B). Thus surface sediment diatoms can also be employed to estimate lake alkalinity.

The following alkalinity predictive equations were obtained using the same 27 lakes employed in the pH predictive equations.

$$\text{Alkalinity} = 204 - 121 \text{ Log Index B} \quad (4)$$

$$F = 38.1, \text{ Prob. } >F = 0.0001, r^2 = 0.60$$

$$\text{Alkalinity} = -4.25 + 0.393\% \text{ACP} + 5.03\% \text{ALP} + 18.2\% \text{ALB} \quad (5)$$

$$F = 26.0, \text{ Prob. } >F = 0.0001, r^2 = 0.77$$

$$\text{Log alkalinity} = 2.089 - 0.00830\% \text{ACP} + 0.00752\% \text{ALP} + 0.00860\% \text{ALB} \quad (6)$$

$$F = 35.2, \text{ Prob. } >F = 0.0001, r^2 = 0.82$$

The equation predicting Log alkalinity (6) has an  $r^2$  value somewhat lower than that obtained with the pH predictive equation 3 ( $r^2 = 0.87$ ) which employed the same diatom pH categories. Thus diatoms can be used with some confidence to predict both pH and alkalinity in Sierra Nevada lake sediment cores.

The analysis has also revealed that the best predictions will be obtained by developing diatom-pH-alkalinity relationships and equations for each area under study. It is doubtful that pH predictive equations developed for Florida, or Maine, or in the Rocky Mountains can provide very satisfactory down core pH (or alkalinity) estimates. Part of difficulty is that widely separated aquatic habitats often have rather different diatom floras. In poorly studied areas like the Sierra Nevada many diatom taxa are new; thus the published literature provides no





Figure 3. Predicted air equilibrated pH values and observed air equilibrated pH values for twenty seven high elevation Sierra Nevada Lakes. A) Log Index B predictive equation (Equation 1). B) Multiple linear regression model (Equation 3).




Figure 4. The relationship between air equilibrated pH and alkalinity in the thirty lakes used to develop the diatom/pH predictive equations. Fig.4a uses untransformed alkalinity while in Fig.4b the alkalinity data has been log transformed. See text for additional information.

information on possible taxon-pH relationships, potentially weakening down core predictive abilities.

The only investigation comparable to the present study of high elevation mountain lakes has been carried out in Rocky Mountain National Park (Baron et al. 1985 - unpublished manuscript). Twenty-three lakes were included in the calibration set in which field-measured pH varied between 5.89 and 9.01 (only 2 lakes had pH values in excess of 7.0). These authors report observing over 110 taxa in 4 lakes cored to a depth of about 50 cm. Of the 14 common dominant diatom taxa in these lakes, none were dominant in Sierra lakes of comparable pH and several were not encountered at all.

Using methods comparable to those of the present study Baron et al. (1985) obtained pH predictive equations employing Log Index Alpha and Log Index B (Renberg and Hellberg 1982). According to the authors these equations were not statistically significant. They then employed the Wilcoxon Sign Test and t-test for paired observations (observed pH and predicted pH) which yielded no significant differences at the 0.05 level, and proceeded to use the non-significant predictive equations for down core diatom inferred pH prediction. This is certainly a questionable procedure. It should also be noted that the surface sediment diatoms in the four lakes for which down core pH reconstructions were obtained were also employed in calculating the pH predictive equations. The validity of including these 4 lakes in the lake calibration data set used in the down core pH reconstruction of these 4 lakes is likewise open to serious criticism. Because of these difficulties and the incompleteness of the unpublished manuscript a meaningful comparison of the Rocky Mountain National Park data and those obtained in the present Sierra study is not yet possible.

Anderson, D., Davis, R. and F. Berge. 1986. Relationships between diatom assemblages in lake surface-sediments and limnological characteristics in southern Norway. In Smol. et al. ed. Diatoms and Lake Acidity. Developments in Hydrobiology 29. W. Junk Publ. pp. 97-109.

Arzet, K., Krause-Dellin, D. and C. Steinberg. 1986. Acidification of four lakes in the Federal Republic of Germany as reflected by diatom assemblages, cladoceran remains, and sediment chemistry. In Smol et al. ed. Diatoms and Lake Acidity. Developments in Hydrobiology 29. W. Junk Publishers. pp. 227-250.

Baron, J., Norton, S., Beeson, D. and R. Herrmann. 1985. Sediment diatom and metal stratigraphy from Rocky Mountain lakes with special reference to atmospheric deposition. Unpublished manuscript. 32 p.

Battarbee, R. 1973. A new method for estimating absolute microfossil numbers, with special reference to diatoms. Limnol. Oceanogr. 28:647-653.

Battarbee, R. W. 1984. Diatom analysis and the acidification of lakes. Phil. Trans. R. Soc. Lond. B:305, 451-477.

Battarbee, R., Smol, J. and J. Merilainen. 1986. Diatoms as indicators of pH: An historical review. In Smol et al. ed. Diatoms and Lake Acidity. Developments in Hydrobiology 29. W. Junk Publishers. pp 5-14.

Berge, F. 1982. Diatoms as indicators of temporal trends in some lakes and rivers in southern Norway. Beih. Nova Hedwigia. 73:249-265.

Bradley, W. 1966. Paleolimnology. In Limnology in North America, ed. D. G. Frey. Univ. Wisconsin Press. pp. 621-652.

Brugum, R. and M. Lusk. 1986. Diatom evidence for neutralization in acid surface mine lakes. In Smol et al. ed. Diatoms and Lake Acidity. Developments in Hydrobiology 29. W. Junk Publ. pp. 115-129.

Charles, D. 1984. Recent pH history of Big Moose Lake (Adirondack Mountains, New York, U.S.A.) inferred from sediment diatom assemblages. Verh. Internat. Verein. Limnol. 22:559-566.

Charles, D. 1985. Relationships between surface sediment diatom assemblages and lakewater characteristics in Adirondack lakes. Ecology 66(3):994-1011.

Cleve-Euler, A. 1951-1955. Die Diatomeen von Schweden und Finland. Kungliga Svenska Vetenskapsakademiens Handlingar, Serien 4:1-961.

Dam, H. van and H. K. van Blokland. 1978. Man-made changes in some Dutch moorland pools, as reflected by historical and recent data about diatoms and macrophytes. *Int. Revue ges. Hydrobiol.* 63:587-607.

Dam, H. van, Surrmond, G. and C. ter Braak. 1981. Impact of acidification on diatoms and chemistry of Dutch moorland pools. *Hydrobiol.* 83:425-459.

Davis, R., Norton, S., Hess, C. and D. Brakke. 1983. Paleolimnological reconstruction of the effects of atmospheric deposition of acids and heavy metals on the chemistry and biology of lakes in New England and Norway. *Hydrobiologia* 103:113-124.

Del Prete, A. and C. Schofield. 1981. The utility of diatom analyses of lake sediments for evaluating acid precipitation effects on dilute lakes. *Arch. Hydrobiol.* 91:332-340.

Dickman, M., Dixit, S., Fortescue, J., Barlow, B. and J. Terasme. 1984. Diatoms as indicators of the rate of lake acidification. *Water, Air, and Soil Poll.* 21:375-386.

Flower, R. J. and R. Battarbee. 1983. Diatom evidence for recent acidification of two Scottish lochs. *Nature*. 305:130-133.

Ford, J. 1986. The recent history of a naturally acidic lake (Cone Pond, N.H.). In Smol *et al.* *Diatoms and Lake Acidity. Developments in Hydrobiology* 29. W. Junk Publishers. pp.131-148.

Germain, H. 1981. Flore des diatomees eaux douces et saumâtres, des éditions Boubee. Paris: 444 pp.

Haworth, E. 1966. Diatoms in a sediment core from Blea Tarn, Langdale. *J. Ecol.* 57:429-441.

Haworth, E. 1976. Two late-glacial (Late Devensian) diatom assemblage profiles from northern Scotland. *New Phytol.* 77:227-256.

Hongve, D. 1972. En bunnhenter som er lett å lage. *Fauna* 25:281-283.

Hustedt, F. 1930. Bacillariophyta (Diatomeae). In Pascher (ed.) *Die Süsswasserflora Mitteluropas* 10, Fisher, Jena: 466 pp.

Hustedt, F. 1927-1966. Die Kieselalgen Deutschlands, Österreichs, und der Schweiz. In L. Rabenhorst's *Kryptogamenflora*, Leipzig. Band 7(1) (1927-1930): 1-920, Band 7(2) (1931-1959): 1-845, Band 7(3) (1961-1966): 1-816.

Hustedt, F. 1939. Systematische und ökologische Untersuchungen über die Diatomeen-Flora von Java, Bali, und Sumatra nach dem Material der Deutschen Limnologischen Sunda-Expedition III. Die ökologischen Factorin und ihr Einfluss auf die Diatomeen flora. Archiv. für Hydrobiol. 16:274-394.

Huttunen, P. and J. Merilainen. 1986. Applications of multivariate techniques to infer limnological conditions from diatom assemblages. In Smol et al. ed. Diatoms and Lake Acidity. Developments in Hydrobiology 29. W. Junk Publ. pp 201-211.

Jørgensen, E. K. 1948. Diatom communities in Danish lakes and ponds. Det Kongelige Danske Videnskabernes Selskab, Biologiske Skrifter 5:1-140.

Lowe, R. L. 1974. Environmental requirements and pollution tolerance of freshwater diatoms. EPA-670/4-74-005. United States Environmental Protection Agency, Cincinnati, Ohio, USA.

Melack, J., Stoddard, J. and C. Ochs. 1985. Major ion chemistry and sensitivity to acid precipitation of Sierra Nevada lakes. Water Resources Research 21:27-32.

Merilainen, J. 1967. The diatom flora and the hydrogen-ion concentration of the water. Ann. Bot. Fenn. 4:51-58.

Nygaard, G. 1956. Ancient and recent flora of diatoms and chrysophyceae in Lake Gribso. Studies on the humic acid Lake Gribso. Fol. Limnol. Scand. 8:32-94.

Patrick, R. and C. W. Reimer. 1966. The diatoms of the United States. Volume 1. Monographs of the Academy of Natural Sciences Number 13, Philadelphia, Pennsylvania, USA.

Patrick, R. and C. W. Reimer. 1975. The diatoms of the United States. Volume 2, Part 1. Monographs of the Academy of Natural Sciences Number 13, Philadelphia, Pennsylvania, USA.

Pennington, W. 1943. Lake sediments: the bottom deposits of the N. Basin of Windermere with special reference to the diatom succession. New Phytol. 43:1-27.

Renberg, I. 1978. Paleolimnology and varve counts of the annually laminated sediment of Lake Rudetjarn, northern Sweden. Early Norland 11:63-92.

Renberg, I. and T. Hellberg. 1982. The pH history of lakes in southwestern Sweden, as calculated from the subfossil diatom flora of the sediments. Ambio. 11(1):30-33.

Simola, H., Kenttämies, K. and O. Sandman. 1985. The recent pH history of some Finnish headwater and seepage lakes, studied by

means of diatom analysis of Pb<sup>210</sup> dated sediment cores. *Aqua Fennica* 15(2):245-255.

Stokes, P. M. and Y. Yung. 1986. Phytoplankton in selected LaCloche (Ontario) lakes, pH 4.2-7.0, with special reference to algae as indicators of chemical characteristics. In Smol et al. ed. *Diatoms and Lake Acidity. Developments in Hydrobiology* 29. W. Junk Publ. pp. 57-72.

Tolonen, K. Liukkonen, M., Jarjula, R. and A. Patila. 1986. Acidification of small lakes in Finland documentd by sedimentary diatom and chrysophycean remains. In Smol et al. ed. *Diatoms and Lake Acidity. Developments in Hydrobiology* 29. W. Junk Publ. pp 159-199.

October 1, 1986  
R. W. Holmes

Appendix A Diatom Taxon List and pH Categories -  
CARB/Sierra Nevada Lake Survey

(ACP = acidophil, CN = circumneutral, ALPF = alkaliphil,  
ALB = alkalibiotic, IND = indifferent, - = unclassified)

---	2001	Achnanthes affinis Grun. v. affinis
ACP	2002	Achnanthes austriaca Hust. v. austriaca
ACP	2003	Achnanthes austriaca Hust. v. helvetica
---	99001	Achnanthes bicapitata Hust.
ACP	2049	Achnanthes bioreti Germain v. bioreti
---	99002	Achnanthes calcar Cl.
---	2004	Achnanthes clevei Grun. v. clevei
CN?	99003	Achnanthes daui v. alaskaensis Foged
ACP	2042	Achnanthes detha Hohn & Hellerm. v. detha
ALP	2006	Achnanthes didyma Hust. v. didyma
---	2012	Achnanthes haukiana Grun. v. haukiana
ALP	2015	Achnanthes lanceolata (Breb.) Grun. v. lanceolata
---	99004	Achnanthes lanceolata f. capitata O. Mull.
---	2016	Achnanthes lanceolata v. dubia Grun.
ACP	99333	Achnanthes lapponica Hust.
ALP	2048	Achnanthes laterostrata Hust. v. laterostrata
---	2022	Achnanthes levanderi Hust. v. levanderi
---	2023	Achnanthes levanderi v. helvetica Hust.
ALP	2024	Achnanthes linearis (W. Sm.) Grun. v. linearis
CN?	2026	Achnanthes linearis v. pusilla Grun.
ACP	2028	Achnanthes marginulata Grun. v. marginulata
ACP	2029	Achnanthes microcephala (Kutz.) Grun. v. microcephala
IND	2030	Achnanthes minutissima Kutz. v. minutissima
---	2047	Achnanthes peragalli v. fossilis Temp. & Perag.
---	99334	Achnanthes recurvata Hust.
ALP	99006	Achnanthes suchlandi Hust.
---	2889	Achnanthes spp.
---	2051	Achnanthes 3-PIRLA
ACP	99007	Achnanthes 1-SN
---	99008	Achnanthes 2-SN
IND	99009	Achnanthes 3-SN
---	99010	Achnanthes 4-SN
---	99011	Achnanthes 6-SN
IND	99012	Achnanthes 7-SN
---	99013	Achnanthes 8-SN
---	99014	Achnanthes 9-SN
ACP	99015	Achnanthes 10-SN
---	99016	Achnanthes 11-SN
---	99017	Achnanthes 13-SN
---	99018	Achnanthes 16-SN
---	99019	Achnanthes 18-SN
ALB	99021	Achnanthes 21-SN
---	99025	Achnanthes 31-SN
CN	99026	Achnanthes 32-SN
---	99027	Achnanthes 34-SN

---	99029	Achnanthes 37-SN
---	99031	Achnanthes 39-SN
---	7001	Amphora ovalis (Kutz.) Kutz. v. ovalis
---	7002	Amphora ovalis v. affinis (Kutz.) V. H. ex Det.
IND	7003	Amphora ovalis v. pediculus (Kutz.) V. H. ex Det.
IND	7004	Amphora perpusilla (Grun.) Grun. v. perpusilla
ACP	8003	Anomoeoneis serians (Breb. ex Kutz.) Cl. v. serians
ACP	8005	Anomoeoneis serians v. brachysira (Breb.) ex Kutz.) Hustedt
ACP	99035	Anomoeoneis serians v. 1-SN
ACP	8001	Anomoeoneis exilis v. lanceolata A. Mayer
ALP	9001	Asterionella formosa Hust. v. formosa
---	9002	Asterionella ralfsii v. americana Korn.
IND	12001	Caloneis bacillum (Grun.) Cl. v. bacillum
ALP	16001	Coccconeis diminuta Pant. v. diminuta
---	99038	Coccconeis placentula Ehr. v. placentula
---	99039	Coccconeis placentula v. euglypta (Ehr.) Cl.
---	99040	Coccconeis 2-SN
ALB	20002	Cyclotella bodanica Eulenst. v. bodanica
ALP	20005	Cyclotella kuetzingiana Thwaites v. kuetzingiana
ALP	20009	Cyclotella ocellata Pant. v. ocellata
ALP	20012	Cyclotella pseudostelligera Hust. v. pseudostelligera
IND	20010	Cyclotella stelligera (Cl. & Grun. V. H. v. stelligera
---	99042	Cyclotella 1-SN
ACP	23004	Cymbella cesatii (Rabh.) Grun. ex A. S. v. cesatii
ALP	23005	Cymbella cistula (Ehr.) Kirchner v. cistula
---	99342	Cymbella cymbiformis Agardh
ACP	23007	Cymbella gaeumannii Meist. v. gaeumannii
ACP	23008	Cymbella hebridica Grun. ex Cl. v. hebridica
ACP	23009	Cymbella lunata W. Sm. v. lunata
ALP	23010	Cymbella microcephala Grun. v. microcephala
IND	23012	Cymbella minuta Hilse ex Rabh. v. minuta
---	23013	Cymbella minuta f. latens (Krasske) Reim.
---	23014	Cymbella minuta v. pseudogracilis (Choln.) Reim.
IND	23015	Cymbella minuta v. silesiaca (Bleisch ex Rabh.) Reim.
---	23031	Cymbella muelleri Hust. v. muelleri
CN?	23016	Cymbella naviculiformis Auersw. ex Heib. v. naviculiformis
---	99050	Cymbella rainierensis Sov.
---	99051	Cymbella sinuata Greg.
---	23889	Cymbella spp.
ACP	23021	Cymbella 1-PIRLA
---	99052	Cymbella 1-SN
---	99053	Cymbella 2-SN
---	99054	Cymbella 3-SN
---	99059	Cymbella 8-SN
ACP	99060	Cymbella 9-SN
---	99061	Cymbella 10-SN
---	99062	Cymbella 11-SN
CN	99350	Cymbella 13-SN
---	27001	Diatoma anceps (Ehr.) Kirchn. v. anceps
IND	27002	Diatoma hiemale v. mesodon (Ehr.) Grun.
CN?	30001	Diploneis elliptica (Kutz.) Cl. v. elliptica
---	30003	Diploneis marginestriata Hust. v. marginestriata

ALP	99067	<i>Epithemia adnata</i> (Kutz.) Breb.
ALP	99069	<i>Epithemia turgida</i> (Ehr.) Kutz.
ACP	33036	<i>Eunotia naegelii</i> Migula v. <i>naegelii</i>
CN?	33001	<i>Eunotia arcus</i> Ehr. v. <i>arcus</i>
ACP	33008	<i>Eunotia curvata</i> (Kutz.) Langerst. v. <i>curvata</i>
---	33009	<i>Eunotia curvata</i> v. <i>capitata</i> (Grun.) Woodhead & Tweed
ACP	33010	<i>Eunotia curvata</i> v. <i>subarcuata</i> (Naeg.) Woodhead & Tweed
---	99071	<i>Eunotia curvata</i> v. 1-SN
ACP	33011	<i>Eunotia denticulata</i> (Breb.) Rabh. v. <i>denticulata</i>
ACP	33015	<i>Eunotia exigua</i> (Breb. ex Kutz.) Rabh. v. <i>exigua</i>
ACP	33018	<i>Eunotia fallax</i> Cl.-Eul. v. <i>fallax</i>
ACP	99072	<i>Eunotia fallax</i> v. <i>gracillima</i> Krasske
ACP	33019	<i>Eunotia flexuosa</i> Breb. ex Kutz. v. <i>flexuosa</i>
---	33021	<i>Eunotia formica</i> Ehr. v. <i>formica</i>
ACP	33026	<i>Eunotia incisa</i> W. Sm. ex Greg. v. <i>incisa</i>
---	33030	<i>Eunotia maior</i> (W. Sm.) v. <i>maior</i>
---	33039	<i>Eunotia pectinalis</i> (O. F. Mull.?) Rabh. v. <i>pectinalis</i>
ACP	33040	<i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.
---	33046	<i>Eunotia praerupta</i> v. <i>bidens</i> (Ehr.) Grun.
---	33051	<i>Eunotia rhomboidea</i> Hust. v. <i>rhomboidea</i>
---	33054	<i>Eunotia serra</i> Ehr. Hust. v. <i>serra</i>
---	99355	<i>Eunotia</i> sp. (girdle view)
---	33059	<i>Eunotia sudetica</i> O. Mull. v. <i>sudetica</i>
ACP	33060	<i>Eunotia tenella</i> (Grun.) Cl. v. <i>tenella</i>
---	33061	<i>Eunotia trinacria</i> Krasske v. <i>trinacria</i>
ACP	33065	<i>Eunotia vanheurckii</i> Pat. v. <i>vanheurckii</i>
---	33066	<i>Eunotia vanheurckii</i> v. <i>intermedia</i> (Krasske) ex Hust. Patr.
---	99074	<i>Eunotia</i> 1-SN
---	99075	<i>Eunotia</i> 2-SN
---	99081	<i>Eunotia</i> 11-SN
---	99082	<i>Eunotia</i> 12-SN
---	99083	<i>Eunotia</i> 13-SN
---	99084	<i>Eunotia</i> 14-SN
IND	34003	<i>Fragilaria brevistriata</i> Grun. v. <i>brevistriata</i>
---	34006	<i>Fragilaria capucina</i> Desm. v. <i>capucina</i>
---	99086	<i>Fragilaria capucina</i> v. <i>mesolepta</i> Rabh.
IND	34030	<i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v. <i>vaucheriae</i>
ALB	34012	<i>Fragilaria construens</i> (Ehr. Grun. v. <i>construens</i>
ALP	34013	<i>Fragilaria construens</i> v. <i>binodis</i> (Ehr.) Grun.
---	34014	<i>Fragilaria construens</i> v. <i>pumila</i> Grun.
IND	34016	<i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.
ALP	34017	<i>Fragilaria crotensis</i> Kitton v. <i>crotensis</i>
ALP	34022	<i>Fragilaria leptostauron</i> (Ehr.) Hust.
ALP	34023	<i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.
---	34025	<i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>
CN	34038	<i>Fragilaria pinnata</i> v. <i>acuminata</i> A. Mayer
---	34026	<i>Fragilaria pinnata</i> v. <i>intercedens</i> (Grun.) Hust.
ALB	34027	<i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.
IND	34030	<i>Fragilaria vaucheriae</i> (Kutz.) Peters var. <i>vaucheriae</i>
ALP	34032	<i>Fragilaria virescens</i> Rolfs v. <i>virescens</i>
IND	34037	<i>Fragilaria virescens</i> v. <i>exigua</i> Grun.
---	34034	<i>Fragilaria virescens</i> v. <i>oblongella</i> Grun.
---	34889	<i>Fragilaria</i> spp.

ALP 99087 *Fragilaria* 6-PIRLA  
 ALP 99088 *Fragilaria* 2-SN  
 --- 99089 *Fragilaria* 4-SN  
 ALB 99090 *Fragilaria* 7-SN  
 --- 99091 *Fragilaria* 8-SN  
 --- 99092 *Fragilaria* 9-SN  
 --- 99093 *Fragilaria* 10-SN  
 --- 99094 *Fragilaria* 11-SN  
 --- 99095 *Fragilaria* 13-SN  
 ALP 99096 *Fragilaria* 14-SN  
 CN 99098 *Fragilaria* 16-SN  
 ACP 35001 *Frustulia rhomboides* (Ehr.) DeT. v. *rhomboides*  
 --- 35002 *Frustulia rhomboides* v. *amphipleuroides* (Grun.) Pet.  
 --- 35003 *Frustulia rhomboides* v. *capitata* (A. Mayer) Patr.  
 ACP 35005 *Frustulia rhomboides* v. *saxonica* (Rabh.) DeT.  
 ACP 99102 *Frustulia rhomboides* v. 1-SN  
 --- 99103 *Gomphonema affine* v. *insigne* (Greg.) Andrews  
 ACP 37003 *Gomphonema angustatum* (Kutz.) Rabh.  
 ALP 37004 *Gomphonema angustatum* v. *citera* (Hohn & Hum) Kutz.  
 --- 37005 *Gomphonema consector* Hohn & Hell. Patr. v. *consector*  
 --- 37010 *Gomphonema parvulum* (Kutz.) Kutz. v. *parvulum*  
 ACP 99105 *Gomphonema puiggarianum* v. *aequatorialis* (Cl.) Camburn  
 ACP 99345 *Gomphonema quadripunctatum* (Ost.) Wils. v.  
     quadripunctatum  
 --- 99106 *Gomphonema subclavatum* (Grun.) Grun.  
 --- 37012 *Gomphonema subtile* Ehr. v. *subtile*  
 ACP 99107 *Gomphonema tackei* v. *abbreviatum* Camburn  
 --- 37014 *Gomphonema truncatum* v. *capitatum* (Ehr.) Patr.  
 --- 99348 *Gomphonema truncatum* Ehr. v. *truncatum*  
 ACP 99335 *Gomphonema ventricosum*  
 --- 37889 *Gomphonema* spp.  
 --- 99109 *Gomphonema* 3-SN  
 --- 99111 *Gomphonema* 5-SN  
 --- 99115 *Gomphonema* 9-SN  
 --- 99117 *Gomphonema* 11-SN  
 ACP 99119 *Gomphonema* 13-SN  
 --- 99120 *Gomphonema* 14-SN  
 --- 99336 *Gyrosigma obtusatum* (Sullivan & Wormley) Boyer  
 --- 99121 *Gyrosigma spencerii* (Quekett) Griffith & Henfries  
 --- 99123 *Hannea arcus* (Ehr.) Pat.  
 --- 40002 *Hantzschia amphioxus* (Ehr.) Grun. v. *amphioxus*  
 CN? 99124 *Krasskella kriegeriana* (Krasske) Ross & Sims  
 ALP 44001 *Melosira ambigua* (Grun.) O. Mull. v. *ambigua*  
 --- 44002 *Melosira distans* (Ehr.) Kutz. v. *distans*  
 ACP 44030 *Melosira distans* v. *nivalis* (W. Sm.) Kirchn.  
 CN? 44008 *Melosira islandica* O. Mull. v. *islandica*  
 ALP 44010 *Melosira italicica* (Ehr.) Kutz. v. *italicica*  
 --- 44011 *Melosira italicica* ssp. *subarctica* O. Mull.  
 ALP 44040 *Melosira italicica* ssp. *subarctica* f. *tenuissima* (Grun.)  
     Camburn  
 ACP 44013 *Melosira italicica* v. *valida* (Grun.) Hust.  
 CN 44014 *Melosira lirata* (Ehr.) Kutz. v. *lirata*  
 ACP 44022 *Melosira perglabra* Ostr. v. *perglabra*  
 --- 44027 *Melosira* 1-PIRLA  
 ACP 99126 *Melosira* 1-SN

--- 99127 *Melosira* 2-SN  
 CN 99128 *Melosira* 3-SN  
 ACP 99129 *Melosira* 4-SN  
 --- 99130 *Melosira* 5-SN  
 ACP 45001 *Meridion circulare* (Grev.) Agardh v. *circulare*  
 --- 46002 *Navicula angusta* Grun. v. *angusta*  
 --- 99131 *Navicula aurora* Sov.  
 --- 46008 *Navicula bremensis* Hust.  
 --- 99132 *Navicula capitata* v. *hungarica* (Grun.) Ross  
 ACP 99133 *Navicula cari* Ehr.  
 --- 46115 *Navicula confervacea* (Kutz.) Grun. v. *fervaceae*  
 ALB 46014 *Navicula cryptocephala* Kutz. v. *cryptocephala*  
 --- 99135 *Navicula cryptocephala* v. *venata* (Kutz.) Rabh.  
 --- 46021 *Navicula globulifera* Hust. v. *globulifera*  
 ACP 46095 *Navicula heimansii* van Dam & Kooyman v. *heimansii*  
 --- 46030 *Navicula krasskei* Hust. v. *krasskei*  
 IND 46032 *Navicula laevissima* Kutz. v. *laevissima*  
 ACP 46038 *Navicula mediocris* Krasske v. *mediocris*  
 ACP 99137 *Navicula mediopunctata* Hust.  
 --- 46042 *Navicula mutica* Kutz. v. *mutica*  
 --- 46102 *Navicula mutica* v. *cohnii* (Hilse) Grun.  
 --- 99138 *Navicula pseudolanceolata* Lange-Bertalot  
 CN 46050 *Navicula pseudoscutiformis* Hust. v. *pseudoscutiformis*  
 ALP 46051 *Navicula pupula* Kutz. v. *pupula*  
 --- 46101 *Navicula pupula* v. *elliptica* Hust.  
 --- 46054 *Navicula pupula* v. *rectangularis* (Greg.) Grun.  
 CN 46056 *Navicula radiosa* Kutz. v. *radiosa*  
 --- 46057 *Navicula radiosa* v. *parva* Wallace  
 ACP 46113 *Navicula cf. subtilissima* v. 2-PIRLA  
 ACP 46121 *Navicula cf. subtilissima* v. 4-PIRLA  
 --- 99355 *Navicula subtilissima* 5 SN  
 --- 46889 *Navicula* spp.  
 ACP 46133 *Navicula* 14-PIRLA  
 ACP 46123 *Navicula* 23-PIRLA  
 IND 99142 *Navicula* 1-SN  
 ACP 99143 *Navicula* 6-SN  
 ACP 99144 *Navicula* 7-SN  
 ACP 99145 *Navicula* 9-SN  
 IND 99146 *Navicula* 10-SN  
 --- 99147 *Navicula* 11-SN  
 ACP 99148 *Navicula* 13-SN  
 ACP 99149 *Navicula* 14-SN  
 ACP 99150 *Navicula* 16-SN  
 --- 99151 *Navicula* 18-SN  
 --- 99152 *Navicula* 21-SN  
 --- 99153 *Navicula* 22-SN  
 --- 99345 *Navicula* 23-SN  
 --- 99154 *Navicula* 24-SN  
 --- 99155 *Navicula* 25-SN  
 ACP 99156 *Navicula* 26-SN  
 ALP 99157 *Navicula* 27-SN  
 --- 99158 *Navicula* 28-SN  
 --- 99161 *Navicula* 30-SN  
 --- 99162 *Navicula* 31-SN  
 ACP 99163 *Navicula* 32-SN

--- 99164 Navicula 35-SN  
 --- 99165 Navicula 36-SN  
 --- 99166 Navicula 37-SN  
 --- 99167 Navicula 38-SN  
 --- 99168 Navicula 39-SN  
 ACP 99172 Navicula 43-SN  
 ALP 99175 Navicula 46-SN  
 ACP 99176 Navicula 47-SN  
 ACP 99179 Navicula 50-SN  
 --- 99180 Navicula 51-SN  
 --- 99181 Navicula 52-SN  
 --- 99182 Navicula 53-SN  
 --- 99183 Navicula 54-SN  
 --- 99184 Navicula 55-SN  
 ACP 99185 Navicula 56-SN  
 --- 99186 Navicula 57-SN  
 ALP 99187 Navicula 58-SN  
 --- 99188 Navicula 59-SN  
 --- 99189 Navicula 60-SN  
 ALP 99190 Navicula 61-SN  
 --- 99194 Navicula 65-SN  
 --- 99198 Navicula 69-SN  
 --- 99200 Navicula 71-SN  
 ACP 99201 Navicula 72-SN  
 --- 99202 Navicula 73-SN  
 --- 99206 Navicula 77-SN  
 --- 99209 Navicula 80-SN  
 --- 99210 Navicula 81-SN  
 --- 99211 Navicula 82-SN  
 --- 99212 Navicula 83-SN  
 --- 99213 Navicula 84-SN  
 --- 99214 Navicula 85-SN  
 ACP 47001 Neidium affine (Ehr.) Pfitz. v. affine  
 CN 47002 Neidium affine v. amphirhynchus (Ehr.) Cl. ACP  
 ACP 47007 Neidium bisulcatum (Lagerst.) Cl. v. bisulcatum  
 --- 47008 Neidium bisulcatum v. baicalense (Skv. & Meyer) Reim.  
 --- 47014 Neidium iridis (Ehr.) Cl. v. iridis  
 --- 47016 Neidium iridis v. amphigomphus (Ehr.) Temp. & Perag.  
 ACP 47025 Neidium 2-PIRLA  
 --- 99218 Neidium 1-SN  
 --- 99219 Neidium 2-SN  
 --- 99220 Neidium 3-SN  
 --- 99221 Neidium 4-SN  
 --- 99223 Neidium 6-SN  
 --- 99226 Nitzschia acuta Hantzsch  
 --- 99227 Nitzschia admissoides Hust.  
 ALB 99341 Nitzschia amphibiodes Hust.  
 --- 48008 Nitzschia dissipata (Kutz.) Grun. v. dissipata  
 ACP 99228 Nitzschia dissipata v. undulata Sovereign  
 --- 48030 Nitzschia romana Grun. v. romana  
 --- 48889 Nitzschia spp.  
 ALP 48035 Nitzschia 1-PIRLA  
 IND 99229 Nitzschia frustulum 1-SN  
 IND 99230 Nitzschia frustulum 2-SN  
 IND 99231 Nitzschia frustulum 3-SN

--- 99233 *Nitzschia frustulum* 5-SN  
 --- 99234 *Nitzschia frustulum* 6-SN  
 IND 99235 *Nitzschia frustulum* 7-SN  
 --- 99236 *Nitzschia frustulum* 8-SN  
 --- 99237 *Nitzschia* 1-SN  
 ACP 99239 *Nitzschia* 3-SN  
 IND 99240 *Nitzschia* 5-SN  
 IND 99241 *Nitzschia* 6-SN  
 ACP 99242 *Nitzschia* 7-SN  
 --- 99243 *Nitzschia* 8-SN  
 ACP 99244 *Nitzschia* 9-SN  
 --- 99246 *Nitzschia* 11-SN  
 --- 99247 *Nitzschia* 12-SN  
 --- 99249 *Nitzschia* 14-SN  
 --- 99250 *Nitzschia* 15-SN  
 --- 99252 *Nitzschia* 17-SN  
 --- 99257 *Nitzschia* 25-SN  
 --- 99259 *Nitzschia* 27-SN  
 --- 99260 *Nitzschia* 28-SN  
 --- 99261 *Nitzschia* 30-SN  
 ALP 99266 *Nitzschia* 35-SN  
 ALP 99267 *Nitzschia* 36-SN  
 --- 99268 *Nitzschia* 37-SN  
 --- 99269 *Nitzschia* 38-SN  
 --- 99270 *Nitzschia* 39-SN  
 ALP 99351 *Nitzschia* 41-SN  
 --- 99356 *Nitzschia* 42-SN  
 ALP 99271 *Opephora martyi* Heribaud  
 --- 52001 *Pinnularia abaujensis* (Pant.) Ross v. *abaujensis*  
 ACP 52002 *Pinnularia abaujensis* v. *linearis* (Hust.) Patr.  
 ACP 52078 *Pinnularia abaujensis* v. 2-PIRLA  
 --- 52011 *Pinnularia biceps* Greg. v. *biceps*  
 --- 52074 *Pinnularia biceps* v. 1-PIRLA  
 --- 52013 *Pinnularia borealis* Ehr. v. *borealis*  
 ACP 52086 *Pinnularia cf.braunii* v. *amphicephala* f. *subconica*  
     Vankantanaraman-PIRLA  
 --- 52025 *Pinnularia divergens* W. Sm. v. *divergens*  
 ACP 52027 *Pinnularia divergentissima* (Grun.) Cl. v.  

*divergentissima*

  
 ACP 52038 *Pinnularia maior* (Kutz.) Rabh. v. *maior*  
 ACP 52080 *Pinnularia cf. pseudomicrostaurón* Gandhi v.  

*pseudomicrostauron*

  
 IND 99274 *Pinnularia subcapitata* v. 1-SN  
 --- 99275 *Pinnularia substomatophora* v. 2-PIRLA  
 --- 52069 *Pinnularia termitina* (Ehr.) Patr. v. *termitina*  
 --- 52889 *Pinnularia* spp.  
 --- 52079 *Pinnularia* 9-PIRLA  
 ACP 99276 *Pinnularia* 1-SN  
 ACP 99277 *Pinnularia* 2-SN  
 --- 99278 *Pinnularia* 5-SN  
 --- 99280 *Pinnularia* 8-SN  
 ACP 99281 *Pinnularia* 9-SN  
 --- 99282 *Pinnularia* 10-SN  
 ACP 99283 *Pinnularia* 11-SN  
 --- 99284 *Pinnularia* 12-SN

--- 99294 Pinnularia 23-SN  
 --- 99297 Pinnularia 26-SN  
 --- 99298 Pinnularia 27-SN  
 --- 99299 Pinnularia 28-SN  
 --- 99303 Pinnularia 32-SN  
 --- 99305 Pinnularia 34-SN  
 --- 99309 Pinnularia 39-SN  
 --- 99310 Pinnularia 40-SN  
 --- 99311 Pinnularia 41-SN  
 --- 99353 Pinnularia 43-SN  
 --- 58001 Rhopalodia gibba (Ehr.) O. Mull. v. gibba  
 --- 62002 Stauroneis anceps Ehr. v. anceps  
 ALP 62003 Stauroneis anceps f. gracilis Rabh.  
 ACP 62024 Stauroneis anceps v. 1-PIRLA  
 ACP 62022 Stauroneis anceps v. 2-PIRLA  
 --- 62015 Stauroneis phoenicenteron (Nitz.) Ehr. v. phoenicenteron  
 --- 63003 Stenopterobia anceps (Lewis) Breb. ex V. H. v. anceps  
 ACP 63002 Stenopterobia intermedia (Lewis) V. H. v. intermedia  
 ALB 99316 Stephanodiscus parvus Stoermer & Hakansson  
 ALB 99317 Stephanodiscus 1-SN  
 --- 99318 Stephanodiscus 2-SN  
 --- 99319 Stephanodiscus 3-SN  
 --- 99320 Stephanodiscus 4-SN  
 --- 99321 Stephanodiscus 5-SN  
 ACP 65011 Surirella delicatissima Lewis v. delicatissima  
 ACP 65033 Surirella delicatissima f. tenuissima Mang.  
 ACP 65014 Surirella linearis W. Sm. v. linearis  
 --- 65015 Surirella linearis v. constricta Grun.  
 --- 99327 Synedra capitata Ehr.  
 ALB 66014 Synedra parasitica (W. Sm.) Hust. v. parasitica  
 --- 99347 Synedra pulchella Kutz.  
 ALP 66015 Synedra radians (Kutz.) v radians  
 --- 66016 Synedra rumpens Kutz. v. rumpens  
 --- 66023 Synedra tenera W. Sm. v. tenera  
 --- 66024 Synedra ulna (Nitz.) Ehr. v. ulna  
 --- 66026 Synedra ulna v. danica (Kutz.) V. H.  
 ALP 66029 Synedra 1-PIRLA  
 ALP 99330 Synedra 2-SN  
 ALP 99331 Synedra 4-SN  
 --- 67002 Tabellaria fenestrata (Lyngb.) Kutz. v. fenestrata  
 IND 67005 Tabellaria flocculosa Roth (Kutz.) strain III sensu  
       Koppen  
 ACP 67006 Tabellaria flocculosa Roth (Kutz.) strain IV sensu  
       Koppen  
       Unidentified SN pennate diatoms

## APPENDIX B

LAKE: PARKER PASS LAT DEG: 37 MIN: 50 SEC: 8 LONG DEG: 119 MIN: 12 SEC: 35  
 QUADRANGLE: MONO CRATERS AREA: 2.85 SHED AREA: 101 ELEV: 3316  
 VEG CLASS: 2 ROCK TYPE: % COVER: A: 1 B: 4 C: 1 D: 1  
 pH FIELD: 5.64 pH LAB: 5.84 CONDUCTIVITY: 16 uS25:  
 HCO<sub>3</sub>: 5.9 Cl: 4.15 NO<sub>3</sub>: 8.32 SO<sub>4</sub>: 106.2 MICR EQUIV/L:  
 Ca: 86.6 Mg: 17.6 Na : 12.7 K : 5.72  
 DATE SAMPLED: 8-10-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 0.9954 GMS PER CC:  
 SED DRY WT: 0.1419 GMS PER CC: INGN LOSS: 14.2

TAXON NAME	Total	No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. <i>austriaca</i>	517.00	3.00	0.580	
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	517.00	40.00	7.737	
2022 <i>Achnanthes levanderi</i> Hust. v. <i>levanderi</i>	517.00	1.00	0.193	
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	517.00	141.00	27.273	
2049 <i>Achnanthes bioreti</i> Germain v. <i>bioreti</i>	517.00	11.00	2.128	
7003 <i>Amphora ovalis</i> v. <i>pediculus</i> (Kutz.) V. H. ex Det.	517.00	3.00	0.580	
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.)	517.00	2.00	0.387	
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	517.00	9.00	1.741	
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. <i>cesatii</i>	517.00	3.00	0.580	
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	517.00	7.00	1.354	
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	517.00	10.00	1.934	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	517.00	9.00	1.741	
33040 <i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.	517.00	1.00	0.193	
35001 <i>Frustulia rhombooides</i> (Ehr.) DeT. v. <i>rhombooides</i>	517.00	4.00	0.774	
44011 <i>Melosira italica</i> ssp. <i>subarctica</i> O. Mull.	517.00	1.00	0.193	
44027 <i>Melosira</i> 1-PIRLA	517.00	32.00	6.190	
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. <i>bisulcatum</i>	517.00	7.00	1.354	
62003 <i>Stauroneis anceps</i> f. <i>gracilis</i> Rabh.	517.00	1.00	0.193	
62022 <i>Stauroneis anceps</i> v. 2-PIRLA	517.00	5.00	0.967	
63002 <i>Stenopterobia intermedia</i> (Lewis) V. H. v. <i>intermedia</i>	517.00	1.00	0.193	
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	517.00	7.00	1.354	
65014 <i>Surirella linearis</i> W. Sm. v. <i>linearis</i>	517.00	4.00	0.774	
99007 <i>Achnanthes</i> 1-SN	517.00	151.00	29.207	
99015 <i>Achnanthes</i> 10-SN	517.00	4.00	0.774	
99018 <i>Achnanthes</i> 16-SN	517.00	2.00	0.387	
99126 <i>Melosira</i> 1-SN	517.00	28.00	5.416	
99143 <i>Navicula</i> 6-SN	517.00	3.00	0.580	
99148 <i>Navicula</i> 13-SN	517.00	8.00	1.547	
99221 <i>Neidium</i> 4-SN	517.00	1.00	0.193	
99239 <i>Nitzschia</i> 3-SN	517.00	5.00	0.967	
99241 <i>Nitzschia</i> 6-SN	517.00	1.00	0.193	
99244 <i>Nitzschia</i> 9-SN	517.00	3.00	0.580	
99277 <i>Pinnularia</i> 2-SN	517.00	6.00	1.161	
99281 <i>Pinnularia</i> 9-SN	517.00	3.00	0.580	

LAKE: TABLE MEADOW LAT DEG: 36 MIN: 36 SEC: 33 LONG DEG: 118 MIN: 39 SEC: 5  
 QUADRANGLE: TRIPLE DIVIDE AREA: 1.55 SHED AREA: 25.1 ELEV: 3139  
 VEG CLASS: 2 ROCK TYPE: % COVER: A: 1 B: 1 C: 4 D: 1  
 pH FIELD: 6.6 pH LAB: 6.01 CONDUCTIVITY: 5.6 uS@25:  
 HCO<sub>3</sub>: 40.8 Cl: 4.01 NO<sub>3</sub>: 0.78 SO<sub>4</sub>: 3.91 MICR EQUIV/L:  
 Ca: 29.8 Mg: 3.45 Na: 13 K: 3.22  
 DATE SAMPLED: 7-20-1985 INVESTIGATOR: J.S.AND R.W.H.  
 SED WET WT: 0.9328 GMS PER CC:  
 SED DRY WT: 0.1212 GMS PER CC: INGN LOSS: 19.4

TAXON NAME	Total No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	508.00	24.00	4.724
2016 <i>Achnanthes lanceolata</i> v. <i>dubia</i> Grun.	508.00	1.00	0.197
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	508.00	20.00	3.937
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	508.00	6.00	1.181
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.	508.00	3.00	0.591
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	508.00	3.00	0.591
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	508.00	1.00	0.197
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	508.00	3.00	0.591
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	508.00	1.00	0.197
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	508.00	1.00	0.197
27001 <i>Diatoma anceps</i> (Ehr.) Kirchn. v. <i>anceps</i>	508.00	1.00	0.197
33008 <i>Eunotia curvata</i> (Kutz.) Langerst. v. <i>curvata</i>	508.00	8.00	1.575
33015 <i>Eunotia exigua</i> (Breb. ex Kutz.) Rabh. v. <i>exigua</i>	508.00	2.00	0.394
33026 <i>Eunotia incisa</i> W. Sm. ex Greg. v. <i>incisa</i>	508.00	2.00	0.394
33040 <i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.	508.00	2.00	0.394
33060 <i>Eunotia tenella</i> (Grun.) Cl. v. <i>tenella</i>	508.00	3.00	0.591
34003 <i>Fragilaria brevistriata</i> Grun. v. <i>brevistriata</i>	508.00	2.00	0.394
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	508.00	2.00	0.394
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	508.00	5.00	0.984
34026 <i>Fragilaria pinnata</i> v. <i>intercedens</i> (Grun.) Hust.	508.00	2.00	0.394
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	508.00	14.00	2.756
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. <i>rhomboides</i>	508.00	6.00	1.181
37003 <i>Gomphonema angustatum</i> (Kutz.) Rabh.	508.00	3.00	0.591
37010 <i>Gomphonema parvulum</i> (Kutz.) Kutz. v. <i>parvulum</i>	508.00	1.00	0.197
40002 <i>Hantzschia amphioxus</i> (Ehr.) Grun. v. <i>amphioxus</i>	508.00	1.00	0.197
44013 <i>Melosira italica</i> v. <i>valida</i> (Grun.) Hust.	508.00	2.00	0.394
44022 <i>Melosira perglabra</i> Ostr. v. <i>perglabra</i>	508.00	1.00	0.197
44027 <i>Melosira</i> 1-PIRLA	508.00	81.00	15.945
44030 <i>Melosira distans</i> v. <i>nivalis</i> (W. Sm.) Kirchn.	508.00	6.00	1.181
45001 <i>Meridion circulare</i> (Grev.) Agardh v. <i>circulare</i>	508.00	6.00	1.181
46038 <i>Navicula mediocris</i> Krasske v. <i>mediocris</i>	508.00	7.00	1.378
46054 <i>Navicula pupula</i> v. <i>rectangularis</i> (Greg.) Grun.	508.00	2.00	0.394
46113 <i>Navicula</i> cf. <i>subtilissima</i> v. 2-PIRLA	508.00	2.00	0.394
46121 <i>Navicula</i> cf. <i>subtilissima</i> v. 4-PIRLA	508.00	2.00	0.394
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. <i>affine</i>	508.00	1.00	0.197
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. <i>bisulcatum</i>	508.00	2.00	0.394
47016 <i>Neidium iridis</i> v. <i>amphigomphus</i> (Ehr.) Temp. & Perag.	508.00	3.00	0.591
47025 <i>Neidium</i> 2-PIRLA	508.00	1.00	0.197
52889 <i>Pinnularia</i> spp.	508.00	1.00	0.197
62022 <i>Stauroneis anceps</i> v. 2-PIRLA	508.00	8.00	1.575
62024 <i>Stauroneis anceps</i> v. 1-PIRLA	508.00	2.00	0.394
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	508.00	14.00	2.756

TAXON NAME	Total No	This Tax	%REL ABUND
65014 <i>Surirella linearis</i> W. Sm. v. <i>linearis</i>	508.00	3.00	0.591
90990 <i>Nitzschia</i> 42-SN	508.00	2.00	0.394
99007 <i>Achnanthes</i> 1-SN	508.00	84.00	16.535
99012 <i>Achnanthes</i> 7-SN	508.00	1.00	0.197
99050 <i>Cymbella rainierensis</i> Sov.	508.00	4.00	0.787
99054 <i>Cymbella</i> 3-SN	508.00	2.00	0.394
99071 <i>Eunotia curvata</i> v. 1-SN	508.00	1.00	0.197
99109 <i>Gomphonema</i> 3-SN	508.00	2.00	0.394
99126 <i>Melosira</i> 1-SN	508.00	49.00	9.646
99133 <i>Navicula cari</i> Ehr.	508.00	9.00	1.772
99137 <i>Navicula mediopunctata</i> Hust.	508.00	1.00	0.197
99142 <i>Navicula</i> 1-SN	508.00	3.00	0.591
99144 <i>Navicula</i> 7-SN	508.00	23.00	4.528
99148 <i>Navicula</i> 13-SN	508.00	10.00	1.969
99149 <i>Navicula</i> 14-SN	508.00	2.00	0.394
99229 <i>Nitzschia frustulum</i> 1-SN	508.00	2.00	0.394
99231 <i>Nitzschia frustulum</i> 3-SN	508.00	1.00	0.197
99239 <i>Nitzschia</i> 3-SN	508.00	14.00	2.756
99240 <i>Nitzschia</i> 5-SN	508.00	3.00	0.591
99247 <i>Nitzschia</i> 12-SN	508.00	2.00	0.394
99281 <i>Pinnularia</i> 9-SN	508.00	29.00	5.709
99283 <i>Pinnularia</i> 11-SN	508.00	2.00	0.394
99333 <i>Achnanthes lapponica</i> Hust.	508.00	1.00	0.197

LAKE: FAIRY SHRIMP LAT DEG: 37 MIN: 26 SEC: 40 LONG DEG: 118 MIN: 45 SEC: 40  
 QUADRANGLE: MT ABBOT AREA: 1.3 SHED AREA: 15.5 ELEV: 3517  
 VEG CLASS: 2 ROCK TYPE: % COVER: A: 1 B: 1 C: 1 D: 4  
 pH FIELD: 6.92 pH LAB: 6.3 CONDUCTIVITY: 3 uS@25:  
 HCO<sub>3</sub>: 14.6 Cl: 1.8 NO<sub>3</sub>: 3.75 SO<sub>4</sub>: 4.15 MICR EQUIV/L:  
 Ca: 16 Mg: 1.96 Na : 4.25 K : 2.29  
 DATE SAMPLED: 8-11-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.8633 GMS PER CC:  
 SED DRY WT: 1.3601 GMS PER CC: INGN LOSS: 2.31

TAXON NAME	Total	No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. <i>austriaca</i>	529.00	10.00	1.890	
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	529.00	199.00	37.618	
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	529.00	197.00	37.240	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	529.00	12.00	2.268	
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	529.00	1.00	0.189	
33046 <i>Eunotia praerupta</i> v. <i>bidens</i> (Ehr.) Grun.	529.00	2.00	0.378	
33060 <i>Eunotia tenella</i> (Grun.) Cl. v. <i>tenella</i>	529.00	3.00	0.567	
44027 <i>Melosira</i> 1-PIRLA	529.00	18.00	3.403	
46030 <i>Navicula krasskei</i> Hust. v. <i>krasskei</i>	529.00	1.00	0.189	
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. <i>bisulcatum</i>	529.00	1.00	0.189	
47025 <i>Neidium</i> 2-PIRLA	529.00	2.00	0.378	
52013 <i>Pinnularia borealis</i> Ehr. v. <i>borealis</i>	529.00	3.00	0.567	
62002 <i>Stauroneis anceps</i> Ehr. v. <i>anceps</i>	529.00	4.00	0.756	
90990 <i>Nitzschia</i> 42-SN	529.00	1.00	0.189	
99015 <i>Achnanthes</i> 10-SN	529.00	6.00	1.134	
99035 <i>Anomoeoneis serians</i> v. 1-SN	529.00	1.00	0.189	
99072 <i>Eunotia fallax</i> v. <i>gracillima</i> Krasske	529.00	11.00	2.079	
99075 <i>Eunotia</i> 2-SN	529.00	1.00	0.189	
99102 <i>Frustulia rhomboides</i> v. 1-SN	529.00	8.00	1.512	
99144 <i>Navicula</i> 7-SN	529.00	1.00	0.189	
99148 <i>Navicula</i> 13-SN	529.00	10.00	1.890	
99176 <i>Navicula</i> 47-SN	529.00	10.00	1.890	
99221 <i>Neidium</i> 4-SN	529.00	2.00	0.378	
99239 <i>Nitzschia</i> 3-SN	529.00	2.00	0.378	
99244 <i>Nitzschia</i> 9-SN	529.00	1.00	0.189	
99274 <i>Pinnularia subcapitata</i> v. 1-SN	529.00	1.00	0.189	
99276 <i>Pinnularia</i> 1-SN	529.00	8.00	1.512	
99277 <i>Pinnularia</i> 2-SN	529.00	1.00	0.189	
99297 <i>Pinnularia</i> 26-SN	529.00	1.00	0.189	
99298 <i>Pinnularia</i> 27-SN	529.00	1.00	0.189	

LAKE: KUNA 11 LAT DEG: 37 MIN: 49 SEC: 50 LONG DEG: 119 MIN: 14 SEC: 15  
 QUADRANGLE: MONO CRATERS AREA: 4.92 SHED AREA: 23.3 ELEV: 3536  
 VEG CLASS: 1 ROCK TYPE: % COVER: A: 1 B: 1 C: 4 D: 1  
 pH FIELD: 7.02 pH LAB: 6.25 CONDUCTIVITY: 3.6 USA25:  
 HCO<sub>3</sub>: 19.1 Cl: 1.79 NO<sub>3</sub>: 4.19 SO<sub>4</sub>: 4.17 MICR EQUIV/L:  
 Ca: 16.2 Mg: 3.28 Na: 6.16 K: 3.16  
 DATE SAMPLED: 8-14-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.5586 GMS PER CC:  
 SED DRY WT: 0.6634 GMS PER CC: INGN LOSS: 3.14

TAXON NAME	Total No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. <i>austriaca</i>	512.00	5.00	0.977
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	512.00	44.00	8.594
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	512.00	207.00	40.430
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	512.00	33.00	6.445
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	512.00	1.00	0.195
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	512.00	5.00	0.977
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	512.00	1.00	0.195
33011 <i>Eunotia denticulata</i> (Breb.) Rabh. v. <i>denticulata</i>	512.00	3.00	0.586
33059 <i>Eunotia sudetica</i> O. Mull. v. <i>sudetica</i>	512.00	1.00	0.195
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	512.00	1.00	0.195
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. <i>rhomboides</i>	512.00	1.00	0.195
44027 <i>Melosira</i> 1-PIRLA	512.00	8.00	1.563
46030 <i>Navicula krasskei</i> Hust. v. <i>krasskei</i>	512.00	3.00	0.586
46133 <i>Navicula</i> 14-PIRLA	512.00	1.00	0.195
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. <i>affine</i>	512.00	1.00	0.195
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. <i>bisulcatum</i>	512.00	3.00	0.586
52086 <i>Pinnularia cf.braunii</i> v. <i>amphicephala</i> f. <i>subconica</i>	512.00	1.00	0.195
62022 <i>Stauroneis anceps</i> v. 2-PIRLA	512.00	2.00	0.391
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	512.00	1.00	0.195
65014 <i>Surirella linearis</i> W. Sm. v. <i>linearis</i>	512.00	2.00	0.391
99007 <i>Achnanthes</i> 1-SN	512.00	111.00	21.680
99015 <i>Achnanthes</i> 10-SN	512.00	26.00	5.078
99035 <i>Anomoeoneis serians</i> v. 1-SN	512.00	1.00	0.195
99107 <i>Gomphonema tackei</i> v. <i>abbreviatum</i> Camburn	512.00	14.00	2.734
99126 <i>Melosira</i> 1-SN	512.00	6.00	1.172
99137 <i>Navicula mediopunctata</i> Hust.	512.00	2.00	0.391
99144 <i>Navicula</i> 7-SN	512.00	1.00	0.195
99145 <i>Navicula</i> 9-SN	512.00	1.00	0.195
99148 <i>Navicula</i> 13-SN	512.00	7.00	1.367
99150 <i>Navicula</i> 16-SN	512.00	2.00	0.391
99156 <i>Navicula</i> 26-SN	512.00	5.00	0.977
99223 <i>Neidium</i> 6-SN	512.00	1.00	0.195
99244 <i>Nitzschia</i> 9-SN	512.00	10.00	1.953
99280 <i>Pinnularia</i> 8-SN	512.00	1.00	0.195

LAKE: MOSQUITO 3 LAT DEG: 36 MIN: 25 SEC: 9 LONG DEG: 118 MIN: 37 SEC: 16  
 QUADRANGLE: TRIPLE DIVIDE AREA: 0.78 SHED AREA: 132 ELEV: 2999  
 VEG CLASS: 4 ROCK TYPE: % COVER: A: 1 B: 1 C: 4 D: 1  
 pH FIELD: 6.52 pH LAB: 6.41 CONDUCTIVITY: 5.8 uS@25:  
 HCO<sub>3</sub>: 51.2 Cl: 2.19 NO<sub>3</sub>: 1.01 SO<sub>4</sub>: 9.91 MICR EQUIV/L:  
 Ca: 45.2 Mg: 5.26 Na : 10.5 K : 3.39  
 DATE SAMPLED: 7-21-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 0.9118 GMS PER CC:  
 SED DRY WT: 0.0745 GMS PER CC: INGN LOSS: 52.5

TAXON NAME	Total	No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	521.00	1.00	0.192	
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	521.00	16.00	3.071	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	521.00	1.00	0.192	
8001 <i>Anomoeoneis exilis</i> v. <i>lanceolata</i> A. Mayer	521.00	3.00	0.576	
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.	521.00	36.00	6.910	
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. <i>cesatii</i>	521.00	4.00	0.768	
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	521.00	45.00	8.637	
23008 <i>Cymbella hebridica</i> Grun. ex Cl. v. <i>hebridica</i>	521.00	6.00	1.152	
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	521.00	38.00	7.294	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	521.00	3.00	0.576	
23021 <i>Cymbella</i> 1-PIRLA	521.00	7.00	1.344	
33009 <i>Eunotia curvata</i> v. <i>capitata</i> (Grun.) Woodhead & Tweed	521.00	1.00	0.192	
33026 <i>Eunotia incisa</i> W. Sm. ex Greg. v. <i>incisa</i>	521.00	13.00	2.495	
33040 <i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.	521.00	4.00	0.768	
33051 <i>Eunotia rhomboidea</i> Hust. v. <i>rhomboidea</i>	521.00	2.00	0.384	
33054 <i>Eunotia serra</i> Ehr. Hust. v. <i>serra</i>	521.00	1.00	0.192	
33059 <i>Eunotia sudetica</i> O. Mull. v. <i>sudetica</i>	521.00	1.00	0.192	
33060 <i>Eunotia tenella</i> (Grun.) Cl. v. <i>tenella</i>	521.00	1.00	0.192	
34017 <i>Fragilaria crotonensis</i> Kitton v. <i>crotonensis</i>	521.00	3.00	0.576	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	521.00	1.00	0.192	
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	521.00	1.00	0.192	
34034 <i>Fragilaria virescens</i> v. <i>oblongella</i> Grun.	521.00	1.00	0.192	
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	521.00	72.00	13.820	
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. <i>rhomboides</i>	521.00	22.00	4.223	
35005 <i>Frustulia rhomboides</i> v. <i>saxonica</i> (Rabh.) DeT.	521.00	15.00	2.879	
37003 <i>Gomphonema angustatum</i> (Kutz.) Rabh.	521.00	4.00	0.768	
37005 <i>Gomphonema consector</i> Hohn & Hell. Patr. v. <i>consector</i>	521.00	2.00	0.384	
44001 <i>Melosira ambigua</i> (Grun.) O. Mull. v. <i>ambigua</i>	521.00	2.00	0.384	
44022 <i>Melosira perglabra</i> Ostr. v. <i>perglabra</i>	521.00	2.00	0.384	
44027 <i>Melosira</i> 1-PIRLA	521.00	8.00	1.536	
44030 <i>Melosira distans</i> v. <i>nivalis</i> (W. Sm.) Kirchn.	521.00	4.00	0.768	
46002 <i>Navicula angusta</i> Grun. v. <i>angusta</i>	521.00	3.00	0.576	
46038 <i>Navicula mediocris</i> Krasske v. <i>mediocris</i>	521.00	10.00	1.919	
46121 <i>Navicula</i> cf. <i>subtilissima</i> v. 4-PIRLA	521.00	7.00	1.344	
46123 <i>Navicula</i> 23-PIRLA	521.00	15.00	2.879	
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. <i>affine</i>	521.00	5.00	0.960	
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. <i>bisulcatum</i>	521.00	1.00	0.192	
52002 <i>Pinnularia abaujensis</i> v. <i>linearis</i> (Hust.) Patr.	521.00	1.00	0.192	
52027 <i>Pinnularia divergentissima</i> (Grun.) Cl. v.	521.00	8.00	1.536	
52080 <i>Pinnularia</i> cf. <i>pseudomicrostauron</i> Gandhi v.	521.00	7.00	1.344	
67005 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain III sensu	521.00	4.00	0.768	
67006 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain IV sensu	521.00	7.00	1.344	

TAXON NAME	Total No	This Tax	%REL ABUND
90990 <i>Nitzschia</i> 42-SN	521.00	2.00	0.384
99007 <i>Achnanthes</i> 1-SN	521.00	13.00	2.495
99053 <i>Cymbella</i> 2-SN	521.00	5.00	0.960
99074 <i>Eunotia</i> 1-SN	521.00	2.00	0.384
99106 <i>Gomphonema subclavatum</i> (Grun.) Grun.	521.00	2.00	0.384
99126 <i>Melosira</i> 1-SN	521.00	30.00	5.758
99128 <i>Melosira</i> 3-SN	521.00	1.00	0.192
99133 <i>Navicula cari</i> Ehr.	521.00	6.00	1.152
99142 <i>Navicula</i> 1-SN	521.00	3.00	0.576
99231 <i>Nitzschia frustulum</i> 3-SN	521.00	48.00	9.213
99239 <i>Nitzschia</i> 3-SN	521.00	1.00	0.192
99274 <i>Pinnularia subcapitata</i> v. 1-SN	521.00	3.00	0.576
99275 <i>Pinnularia substomatophora</i> v. 2-PIRLA	521.00	1.00	0.192
99281 <i>Pinnularia</i> 9-SN	521.00	7.00	1.344
99284 <i>Pinnularia</i> 12-SN	521.00	1.00	0.192
99334 <i>Achnanthes recurvata</i> Hust.	521.00	1.00	0.192

LAKE: MOSQUITO 5 LAT DEG: 36 MIN: 24 SEC: 53 LONG DEG: 118 MIN: 37 SEC: 35  
 QUADRANGLE: TRIPLE DIVIDE AREA: 3.11 SHED AREA: 76.7 ELEV: 3048  
 VEG CLASS: 3 ROCK TYPE: % COVER: A: 1 B: 1 C: 4 D: 1  
 pH FIELD: 6.87 pH LAB: 6.42 CONDUCTIVITY: 6.7 uS@25:  
 HCO<sub>3</sub>: 52.8 Cl: 1.51 NO<sub>3</sub>: 2.18 SO<sub>4</sub>: 13.25 MICR EQUIV/L:  
 Ca: 51 Mg: 4.49 Na : 11 K : 3.16  
 DATE SAMPLED: 7-21-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.047 GMS PER CC:  
 SED DRY WT: 0.1102 GMS PER CC: INGN LOSS: 15.9

TAXON NAME	Total No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. austriaca	499.00	1.00	0.200
2003 <i>Achnanthes austriaca</i> Hust. v. helvetica	499.00	18.00	3.607
2028 <i>Achnanthes marginulata</i> Grun. v. marginulata	499.00	41.00	8.216
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. detha	499.00	12.00	2.405
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.	499.00	1.00	0.200
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	499.00	8.00	1.603
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	499.00	7.00	1.403
23008 <i>Cymbella hebridica</i> Grun. ex Cl. v. <i>hebridica</i>	499.00	1.00	0.200
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	499.00	22.00	4.409
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	499.00	20.00	4.008
27002 <i>Diatoma hiemale</i> v. <i>mesodon</i> (Ehr.) Grun.	499.00	1.00	0.200
30001 <i>Diploneis elliptica</i> (Kutz.) Cl. v. <i>elliptica</i>	499.00	2.00	0.401
33008 <i>Eunotia curvata</i> (Kutz.) Langerst. v. <i>curvata</i>	499.00	1.00	0.200
33040 <i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.	499.00	1.00	0.200
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	499.00	8.00	1.603
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	499.00	10.00	2.004
35001 <i>Frustulia rhomboidea</i> (Ehr.) DeT. v. <i>rhomboidea</i>	499.00	7.00	1.403
37010 <i>Gomphonema parvulum</i> (Kutz.) Kutz. v. <i>parvulum</i>	499.00	2.00	0.401
44010 <i>Melosira italica</i> (Ehr.) Kutz. v. <i>italica</i>	499.00	2.00	0.401
44011 <i>Melosira italica</i> ssp. <i>subarctica</i> O. Mull.	499.00	5.00	1.002
44013 <i>Melosira italica</i> v. <i>valida</i> (Grun.) Hust.	499.00	1.00	0.200
44027 <i>Melosira</i> 1-PIRLA	499.00	9.00	1.804
44040 <i>Melosira italica</i> ssp. <i>subarctica</i> f. <i>tenuissima</i> (Grun.)	499.00	1.00	0.200
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	499.00	1.00	0.200
46050 <i>Navicula pseudoscutiformis</i> Hust. v. <i>pseudoscutiformis</i>	499.00	2.00	0.401
46101 <i>Navicula pupula</i> v. <i>elliptica</i> Hust.	499.00	3.00	0.601
46133 <i>Navicula</i> 14-PIRLA	499.00	10.00	2.004
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. <i>affine</i>	499.00	1.00	0.200
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. <i>bisulcatum</i>	499.00	4.00	0.802
48030 <i>Nitzschia romana</i> Grun. v. <i>romana</i>	499.00	1.00	0.200
52027 <i>Pinnularia divergentissima</i> (Grun.) Cl. v.	499.00	1.00	0.200
62003 <i>Stauroneis anceps</i> f. <i>gracilis</i> Rabh.	499.00	2.00	0.401
62022 <i>Stauroneis anceps</i> v. 2-PIRLA	499.00	5.00	1.002
63002 <i>Stenopterobia intermedia</i> (Lewis) V. H. v. <i>intermedia</i>	499.00	1.00	0.200
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	499.00	4.00	0.802
65015 <i>Surirella linearis</i> v. <i>constricta</i> Grun.	499.00	1.00	0.200
99007 <i>Achnanthes</i> 1-SN	499.00	144.00	28.858
99015 <i>Achnanthes</i> 10-SN	499.00	7.00	1.403
99105 <i>Gomphonema puiggarianum</i> v. <i>aequatorialis</i> (Cl.) Camburn	499.00	10.00	2.004
99126 <i>Melosira</i> 1-SN	499.00	24.00	4.810
99133 <i>Navicula cari</i> Ehr.	499.00	5.00	1.002
99137 <i>Navicula mediopunctata</i> Hust.	499.00	3.00	0.601

TAXON NAME	Total No	This Tax	%REL ABUND
99142 Navicula 1-SN	499.00	26.00	5.210
99145 Navicula 9-SN	499.00	10.00	2.004
99146 Navicula 10-SN	499.00	6.00	1.202
99148 Navicula 13-SN	499.00	5.00	1.002
99153 Navicula 22-SN	499.00	1.00	0.200
99172 Navicula 43-SN	499.00	2.00	0.401
99228 Nitzschia dissipata v. undulata Sovereign	499.00	1.00	0.200
99230 Nitzschia frustulum 2-SN	499.00	2.00	0.401
99239 Nitzschia 3-SN	499.00	2.00	0.401
99242 Nitzschia 7-SN	499.00	4.00	0.802
99243 Nitzschia 8-SN	499.00	1.00	0.200
99281 Pinnularia 9-SN	499.00	27.00	5.411
99345 Gomphonema quadripunctatum (Ost.) Wils. v.	499.00	2.00	0.401

LAKE: L. CATHEDRAL LAT DEG: 37 MIN: 50 SEC: 21 LONG DEG: 119 MIN: 24 SEC: 50  
 QUADRANGLE: TUOL.MEADOWS AREA: 4.66 SHED AREA: 148 ELEV: 2902  
 VEG CLASS: ROCK TYPE: % COVER: A: 1 B: 1 C: 1 D: 4  
 pH FIELD: 6.45 pH LAB: 6.44 CONDUCTIVITY: 3.2 USA25:  
 HCO<sub>3</sub>: 24.6 Cl: 1.47 NO<sub>3</sub>: 0 SO<sub>4</sub>: 2.6 MICR EQUIV/L:  
 Ca: 16.4 Mg: 3.23 Na: 8.67 K: 1.94  
 DATE SAMPLED: 8-16-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.1618 GMS PER CC:  
 SED DRY WT: 0.0614 GMS PER CC: INGN LOSS: 41.9

TAXON NAME	Total No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	507.00	3.00	0.592
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	507.00	14.00	2.761
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	507.00	1.00	0.197
8003 <i>Anomoeoneis serians</i> (Breb. ex Kutz.) Cl. v. <i>serians</i>	507.00	6.00	1.183
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.)	507.00	36.00	7.101
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	507.00	19.00	3.748
23008 <i>Cymbella hebridica</i> Grun. ex Cl. v. <i>hebridica</i>	507.00	2.00	0.394
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	507.00	3.00	0.592
23021 <i>Cymbella</i> 1-PIRLA	507.00	61.00	12.032
33010 <i>Eunotia curvata</i> v. <i>subarcuata</i> (Naeg.) Woodhead & Tweed	507.00	2.00	0.394
33015 <i>Eunotia exigua</i> (Breb. ex Kutz.) Rabh. v. <i>exigua</i>	507.00	4.00	0.789
33026 <i>Eunotia incisa</i> W. Sm. ex Greg. v. <i>incisa</i>	507.00	3.00	0.592
33036 <i>Eunotia naegelii</i> Migula v. <i>naegelii</i>	507.00	3.00	0.592
33039 <i>Eunotia pectinalis</i> (O. F. Mull.?) Rabh. v. <i>pectinalis</i>	507.00	1.00	0.197
33040 <i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.	507.00	3.00	0.592
33060 <i>Eunotia tenella</i> (Grun.) Cl. v. <i>tenella</i>	507.00	1.00	0.197
33065 <i>Eunotia vanheurckii</i> Pat. v. <i>vanheurckii</i>	507.00	2.00	0.394
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. <i>rhomboides</i>	507.00	8.00	1.578
35005 <i>Frustulia rhomboides</i> v. <i>saxonica</i> (Rabh.) DeT.	507.00	8.00	1.578
37003 <i>Gomphonema angustatum</i> (Kutz.) Rabh.	507.00	4.00	0.789
44022 <i>Melosira perglabra</i> Ostr. v. <i>perglabra</i>	507.00	1.00	0.197
44027 <i>Melosira</i> 1-PIRLA	507.00	16.00	3.156
44030 <i>Melosira distans</i> v. <i>nivalis</i> (W. Sm.) Kirchn.	507.00	15.00	2.959
45001 <i>Meridion circulare</i> (Grev.) Agardh v. <i>circulare</i>	507.00	2.00	0.394
46021 <i>Navicula globulifera</i> Hust. v. <i>globulifera</i>	507.00	1.00	0.197
46038 <i>Navicula mediocris</i> Krasske v. <i>mediocris</i>	507.00	20.00	3.945
46056 <i>Navicula radiosa</i> Kutz. v. <i>radiosa</i>	507.00	1.00	0.197
46095 <i>Navicula heimansii</i> van Dam & Kooyman v. <i>heimansii</i>	507.00	1.00	0.197
46121 <i>Navicula</i> cf. <i>subtilissima</i> v. 4-PIRLA	507.00	6.00	1.183
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. <i>affine</i>	507.00	1.00	0.197
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. <i>bisulcatum</i>	507.00	4.00	0.789
52002 <i>Pinnularia abaujensis</i> v. <i>linearis</i> (Hust.) Patr.	507.00	1.00	0.197
52011 <i>Pinnularia biceps</i> Greg. v. <i>biceps</i>	507.00	1.00	0.197
52038 <i>Pinnularia maior</i> (Kutz.) Rabh. v. <i>maior</i>	507.00	1.00	0.197
52069 <i>Pinnularia termitina</i> (Ehr.) Patr. v. <i>termitina</i>	507.00	1.00	0.197
52079 <i>Pinnularia</i> 9-PIRLA	507.00	1.00	0.197
52080 <i>Pinnularia</i> cf. <i>pseudomicrostauron</i> Gandhi v.	507.00	9.00	1.775
52086 <i>Pinnularia</i> cf. <i>braunii</i> v. <i>amphicephala</i> f. <i>subconica</i>	507.00	5.00	0.986
63002 <i>Stenopterobia intermedia</i> (Lewis) V. H. v. <i>intermedia</i>	507.00	2.00	0.394
65033 <i>Surirella delicatissima</i> f. <i>tenuissima</i> Mang.	507.00	9.00	1.775
67005 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain III sensu	507.00	3.00	0.592
99035 <i>Anomoeoneis serians</i> v. 1-SN	507.00	18.00	3.550

TAXON NAME	Total	No	This Tax	%REL ABUND
99060 Cymbella 9-SN	507.00	11.00	2.170	
99082 Eunotia 12-SN	507.00	1.00	0.197	
99083 Eunotia 13-SN	507.00	2.00	0.394	
99102 Frustulia rhomboides v. l-SN	507.00	7.00	1.381	
99105 Gomphonema puiggarianum v. aequatorialis (Cl.) Camburn	507.00	8.00	1.578	
99126 Melosira 1-SN	507.00	58.00	11.440	
99133 Navicula cari Ehr.	507.00	12.00	2.367	
99144 Navicula 7-SN	507.00	72.00	14.201	
99231 Nitzschia frustulum 3-SN	507.00	6.00	1.183	
99241 Nitzschia 6-SN	507.00	4.00	0.789	
99274 Pinnularia subcapitata v. 1-SN	507.00	3.00	0.592	
99277 Pinnularia 2-SN	507.00	10.00	1.972	
99278 Pinnularia 5-SN	507.00	1.00	0.197	
99281 Pinnularia 9-SN	507.00	7.00	1.381	
99305 Pinnularia 34-SN	507.00	1.00	0.197	
99309 Pinnularia 39-SN	507.00	1.00	0.197	

LAKE: UP. TREASURE LAT DEG: 37 MIN: 23 SEC: 13 LONG DEG: 118 MIN: 46 SEC: 0  
 QUADRANGLE: MT. ABBOT AREA: 2.85 SHED AREA: 184 ELEV: 3389  
 VEG CLASS: 3 ROCK TYPE: % COVER: A: 1 B: 1 C: 4 D: 1  
 pH FIELD: 6.89 pH LAB: 6.45 CONDUCTIVITY: 5.7 uS@25:  
 HCO<sub>3</sub>: 27.2 Cl: 1.05 NO<sub>3</sub>: 0 SO<sub>4</sub>: 4.72 MICR EQUIV/L:  
 Ca: 25.7 Mg: 2.07 Na : 5.97 K : 6.52  
 DATE SAMPLED: 8-7-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.5195 GMS PER CC:  
 SED DRY WT: 0.7577 GMS PER CC: INGN LOSS: 3.12

TAXON NAME	Total	No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. austriaca	508.00	9.00	1.772	
2003 <i>Achnanthes austriaca</i> Hust. v. helvetica	508.00	46.00	9.055	
2028 <i>Achnanthes marginulata</i> Grun. v. marginulata	508.00	91.00	17.913	
2030 <i>Achnanthes minutissima</i> Kutz. v. minutissima	508.00	2.00	0.394	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. detha	508.00	21.00	4.134	
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. perpusilla	508.00	5.00	0.984	
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. cesatii	508.00	4.00	0.787	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. minuta	508.00	16.00	3.150	
23014 <i>Cymbella minuta</i> v. <i>pseudogracilis</i> (Choln.) Reim.	508.00	2.00	0.394	
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	508.00	42.00	8.268	
33018 <i>Eunotia fallax</i> Cl.-Eul. v. fallax	508.00	1.00	0.197	
33060 <i>Eunotia tenella</i> (Grun.) Cl. v. tenella	508.00	1.00	0.197	
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	508.00	1.00	0.197	
35001 <i>Frustulia rhombooides</i> (Ehr.) DeT. v. rhombooides	508.00	7.00	1.378	
40002 <i>Hantzschia amphioxus</i> (Ehr.) Grun. v. amphioxus	508.00	2.00	0.394	
44027 <i>Melosira</i> 1-PIRLA	508.00	3.00	0.591	
44030 <i>Melosira distans</i> v. <i>nivalis</i> (W. Sm.) Kirchn.	508.00	1.00	0.197	
45001 <i>Meridion circulare</i> (Grev.) Agardh v. circulare	508.00	2.00	0.394	
46133 <i>Navicula</i> 14-PIRLA	508.00	15.00	2.953	
47025 <i>Neidium</i> 2-PIRLA	508.00	3.00	0.591	
52002 <i>Pinnularia abaujensis</i> v. <i>linearis</i> (Hust.) Patr.	508.00	1.00	0.197	
52078 <i>Pinnularia abaujensis</i> v. 2-PIRLA	508.00	1.00	0.197	
62003 <i>Stauroneis anceps</i> f. <i>gracilis</i> Rabh.	508.00	3.00	0.591	
62022 <i>Stauroneis anceps</i> v. 2-PIRLA	508.00	4.00	0.787	
65014 <i>Surirella linearis</i> W. Sm. v. linearis	508.00	6.00	1.181	
65033 <i>Surirella delicatissima</i> f. <i>tenuissima</i> Mang.	508.00	1.00	0.197	
67006 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain IV sensu	508.00	1.00	0.197	
90990 <i>Nitzschia</i> 42-SN	508.00	4.00	0.787	
99001 <i>Achnanthes bicapitata</i> Hust.	508.00	4.00	0.787	
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	508.00	8.00	1.575	
99007 <i>Achnanthes</i> 1-SN	508.00	88.00	17.323	
99009 <i>Achnanthes</i> 3-SN	508.00	3.00	0.591	
99015 <i>Achnanthes</i> 10-SN	508.00	28.00	5.512	
99059 <i>Cymbella</i> 8-SN	508.00	2.00	0.394	
99060 <i>Cymbella</i> 9-SN	508.00	1.00	0.197	
99102 <i>Frustulia rhombooides</i> v. 1-SN	508.00	4.00	0.787	
99107 <i>Gomphonema tackei</i> v. <i>abbreviatum</i> Camburn	508.00	16.00	3.150	
99119 <i>Gomphonema</i> 13-SN	508.00	2.00	0.394	
99126 <i>Melosira</i> 1-SN	508.00	10.00	1.969	
99145 <i>Navicula</i> 9-SN	508.00	2.00	0.394	
99148 <i>Navicula</i> 13-SN	508.00	6.00	1.181	
99150 <i>Navicula</i> 16-SN	508.00	14.00	2.756	

TAXON NAME	Total	No	This Tax	%REL ABUND
99172 Navicula 43-SN	508.00	1.00	0.197	
99176 Navicula 47-SN	508.00	1.00	0.197	
99194 Navicula 65-SN	508.00	3.00	0.591	
99206 Navicula 77-SN	508.00	1.00	0.197	
99211 Navicula 82-SN	508.00	2.00	0.394	
99212 Navicula 83-SN	508.00	1.00	0.197	
99213 Navicula 84-SN	508.00	2.00	0.394	
99228 Nitzschia dissipata v. undulata Sovereign	508.00	2.00	0.394	
99242 Nitzschia 7-SN	508.00	2.00	0.394	
99244 Nitzschia 9-SN	508.00	5.00	0.984	
99247 Nitzschia 12-SN	508.00	1.00	0.197	
99274 Pinnularia subcapitata v. 1-SN	508.00	1.00	0.197	
99281 Pinnularia 9-SN	508.00	2.00	0.394	
99311 Pinnularia 41-SN	508.00	1.00	0.197	

LAKE: DADE LAT DEG: 37 MIN: 22 SEC: 47 LONG DEG: 118 MIN: 45 SEC: 42  
 QUADRANGLE: MT.ABBOT AREA: 3.11 SHED AREA: 130 ELEV: 3511  
 VEG CLASS: 1 ROCK TYPE: % COVER: A: 1 B: 1 C: 4 D: 1  
 pH FIELD: 6.8 pH LAB: 6.48 CONDUCTIVITY: 4.9 us@25:  
 HCO<sub>3</sub>: 24 Cl: 1.18 NO<sub>3</sub>: SO<sub>4</sub>: 5.84 MICR EQUIV/L:  
 Ca: 31.7 Mg: 2.45 Na : 5.28 K : 2.18  
 DATE SAMPLED: 8-7-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.8756 GMS PER CC:  
 SED DRY WT: 1.2642 GMS PER CC: INGN LOSS: 1.08

TAXON NAME	Total No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. austriaca	514.00	1.00	0.195
2003 <i>Achnanthes austriaca</i> Hust. v. helvetica	514.00	35.00	6.809
2015 <i>Achnanthes lanceolata</i> (Breb.) Grun. v. lanceolata	514.00	1.00	0.195
2028 <i>Achnanthes marginulata</i> Grun. v. marginulata	514.00	114.00	22.179
2030 <i>Achnanthes minutissima</i> Kutz. v. minutissima	514.00	16.00	3.113
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. detha	514.00	53.00	10.311
7003 <i>Amphora ovalis</i> v. <i>pediculus</i> (Kutz.) V. H. ex Det.	514.00	2.00	0.389
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	514.00	1.00	0.195
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	514.00	41.00	7.977
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	514.00	9.00	1.751
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	514.00	1.00	0.195
35005 <i>Frustulia rhombooides</i> v. <i>saxonica</i> (Rabh.) DeT.	514.00	1.00	0.195
45001 <i>Meridion circulare</i> (Grev.) Agardh v. <i>circulare</i>	514.00	1.00	0.195
46042 <i>Navicula mutica</i> Kutz. v. <i>mutica</i>	514.00	1.00	0.195
46133 <i>Navicula</i> 14-PIRLA	514.00	2.00	0.389
47025 <i>Neidium</i> 2-PIRLA	514.00	2.00	0.389
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	514.00	1.00	0.195
65014 <i>Surirella linearis</i> W. Sm. v. <i>linearis</i>	514.00	1.00	0.195
67006 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain IV sensu	514.00	1.00	0.195
90990 <i>Nitzschia</i> 42-SN	514.00	2.00	0.389
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	514.00	7.00	1.362
99007 <i>Achnanthes</i> 1-SN	514.00	139.00	27.043
99008 <i>Achnanthes</i> 2-SN	514.00	1.00	0.195
99010 <i>Achnanthes</i> 4-SN	514.00	13.00	2.529
99015 <i>Achnanthes</i> 10-SN	514.00	11.00	2.140
99016 <i>Achnanthes</i> 11-SN	514.00	2.00	0.389
99017 <i>Achnanthes</i> 13-SN	514.00	1.00	0.195
99107 <i>Gomphonema tackei</i> v. <i>abbreviatum</i> Camburn	514.00	17.00	3.307
99137 <i>Navicula mediopunctata</i> Hust.	514.00	2.00	0.389
99145 <i>Navicula</i> 9-SN	514.00	2.00	0.389
99148 <i>Navicula</i> 13-SN	514.00	3.00	0.584
99150 <i>Navicula</i> 16-SN	514.00	11.00	2.140
99151 <i>Navicula</i> 18-SN	514.00	4.00	0.778
99219 <i>Neidium</i> 2-SN	514.00	1.00	0.195
99220 <i>Neidium</i> 3-SN	514.00	1.00	0.195
99239 <i>Nitzschia</i> 3-SN	514.00	5.00	0.973
99247 <i>Nitzschia</i> 12-SN	514.00	1.00	0.195
99282 <i>Pinnularia</i> 10-SN	514.00	1.00	0.195
99335 <i>Gomphonema ventricosum</i>	514.00	6.00	1.167

LAKE: LE CONTE LAT DEG: 38 MIN: 52 SEC: 20 LONG DEG: 120 MIN: 8 SEC: 10  
 QUADRANGLE: FALLEN LEAF LK. AREA: SHED AREA: ELEV: 2500  
 VEG CLASS: ROCK TYPE: % COVER: A: B: C: D:  
 pH FIELD: 6.25 pH LAB: 6.55 CONDUCTIVITY: 3.1 USA25:  
 HCO<sub>3</sub>: 31.1 Cl: 2.82 NO<sub>3</sub>: 0 SO<sub>4</sub>: 3.97 MICR EQUIV/L:  
 Ca: 11.2 Mg: 2.84 Na: 5.13 K: 2.18  
 DATE SAMPLED: 8-22-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 0.9048 GMS PER CC:  
 SED DRY WT: 0.0773 GMS PER CC: INGN LOSS: 27.7

TAXON NAME	Total No	This Tax	%REL ABUND
2002 Achnanthes austriaca Hust. v. austriaca	496.00	6.00	1.210
2003 Achnanthes austriaca Hust. v. helvetica	496.00	8.00	1.613
2028 Achnanthes marginulata Grun. v. marginulata	496.00	31.00	6.250
8005 Anomoeoneis serians v. brachysira (Breb.) ex Kutz.	496.00	10.00	2.016
23007 Cymbella gaeumannii Meist. v. gaeumannii	496.00	14.00	2.823
23009 Cymbella lunata W. Sm. v. lunata	496.00	13.00	2.621
23012 Cymbella minuta Hilse ex Rabh. v. minuta	496.00	1.00	0.202
23015 Cymbella minuta v. silesiaca (Bleisch ex Rabh.) Reim.	496.00	2.00	0.403
23021 Cymbella 1-PIRLA	496.00	6.00	1.210
33018 Eunotia fallax Cl.-Eul. v. fallax	496.00	5.00	1.008
35001 Frustulia rhomboides (Ehr.) DeT. v. rhomboides	496.00	17.00	3.427
35005 Frustulia rhomboides v. saxonica (Rabh.) DeT.	496.00	1.00	0.202
37003 Gomphonema angustatum (Kutz.) Rabh.	496.00	1.00	0.202
44027 Melosira 1-PIRLA	496.00	59.00	11.895
44030 Melosira distans v. nivalis (W. Sm.) Kirchn.	496.00	4.00	0.806
46008 Navicula bremensis Hust.	496.00	2.00	0.403
46038 Navicula mediocris Krasske v. mediocris	496.00	65.00	13.105
46095 Navicula heimansii van Dam & Kooyman v. heimansii	496.00	7.00	1.411
46115 Navicula confervacea (Kutz.) Grun. v. fervaceae	496.00	2.00	0.403
46121 Navicula cf. subtilissima v. 4-PIRLA	496.00	6.00	1.210
47007 Neidium bisulcatum (Lagerst.) Cl. v. bisulcatum	496.00	1.00	0.202
52002 Pinnularia abaujensis v. linearis (Hust.) Patr.	496.00	1.00	0.202
52027 Pinnularia divergentissima (Grun.) Cl. v.	496.00	2.00	0.403
52086 Pinnularia cf. braunii v. amphicephala f. subconica	496.00	21.00	4.234
62002 Stauroneis anceps Ehr. v. anceps	496.00	1.00	0.202
62022 Stauroneis anceps v. 2-PIRLA	496.00	2.00	0.403
63002 Stenopterobia intermedia (Lewis) V. H. v. intermedia	496.00	1.00	0.202
65014 Surirella linearis W. Sm. v. linearis	496.00	4.00	0.806
65033 Surirella delicatissima f. tenuissima Mang.	496.00	2.00	0.403
90990 Nitzschia 42-SN	496.00	2.00	0.403
99035 Anomoeoneis serians v. 1-SN	496.00	1.00	0.202
99060 Cymbella 9-SN	496.00	20.00	4.032
99081 Eunotia 11-SN	496.00	1.00	0.202
99102 Frustulia rhomboides v. l-SN	496.00	22.00	4.435
99105 Gomphonema puiggarianum v. aequatorialis (Cl.) Camburn	496.00	1.00	0.202
99126 Melosira 1-SN	496.00	26.00	5.242
99129 Melosira 4-SN	496.00	13.00	2.621
99144 Navicula 7-SN	496.00	102.00	20.565
99148 Navicula 13-SN	496.00	1.00	0.202
99218 Neidium 1-SN	496.00	2.00	0.403
99231 Nitzschia frustulum 3-SN	496.00	1.00	0.202
99241 Nitzschia 6-SN	496.00	5.00	1.008

TAXON NAME	Total	No	This Tax	%REL ABUND
99266 <i>Nitzschia</i> 35-SN	496.00	1.00	0.202	
99277 <i>Pinnularia</i> 2-SN	496.00	2.00	0.403	
99281 <i>Pinnularia</i> 9-SN	496.00	1.00	0.202	

LAKE: SUMMIT LAT DEG: 37 MIN: 26 SEC: 0 LONG DEG: 118 MIN: 46 SEC: 11  
 QUADRANGLE: MT.ABBOT AREA: 1.81 SHED AREA: 82.9 ELEV: 3609  
 VEG CLASS: 1 ROCK TYPE: % COVER: A: 1 B: 1 C: 1 D: 4  
 pH FIELD: 7.1 pH LAB: 6.6 CONDUCTIVITY: 2.8 uS@25:  
 HCO<sub>3</sub>: 21.4 Cl: 1.14 NO<sub>3</sub>: 0 SO<sub>4</sub>: 3.88 MICR EQUIV/L:  
 Ca: 13.8 Mg: 2.18 Na : 5.43 K : 4.21  
 DATE SAMPLED: 8-11-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.46 GMS PER CC:  
 SED DRY WT: 0.8082 GMS PER CC: INGN LOSS: 1.84

TAXON NAME	Total No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	507.00	18.00	3.550
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	507.00	172.00	33.925
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	507.00	17.00	3.353
2051 <i>Achnanthes</i> 3-PIRLA	507.00	1.00	0.197
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	507.00	2.00	0.394
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	507.00	5.00	0.986
23021 <i>Cymbella</i> 1-PIRLA	507.00	5.00	0.986
33011 <i>Eunotia denticulata</i> (Breb.) Rabh. v. <i>denticulata</i>	507.00	2.00	0.394
33051 <i>Eunotia rhomboidea</i> Hust. v. <i>rhomboidea</i>	507.00	1.00	0.197
33061 <i>Eunotia trinacria</i> Krasske v. <i>trinacria</i>	507.00	1.00	0.197
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	507.00	3.00	0.592
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	507.00	7.00	1.381
35005 <i>Frustulia rhombooides</i> v. <i>saxonica</i> (Rabh.) DeT.	507.00	2.00	0.394
44002 <i>Melosira distans</i> (Ehr.) Kutz. v. <i>distans</i>	507.00	1.00	0.197
44027 <i>Melosira</i> 1-PIRLA	507.00	137.00	27.022
44030 <i>Melosira distans</i> v. <i>nivalis</i> (W. Sm.) Kirchn.	507.00	8.00	1.578
45001 <i>Meridion circulare</i> (Grev.) Agardh v. <i>circulare</i>	507.00	4.00	0.789
46121 <i>Navicula</i> cf. <i>subtilissima</i> v. 4-PIRLA	507.00	1.00	0.197
46133 <i>Navicula</i> 14-PIRLA	507.00	6.00	1.183
52074 <i>Pinnularia biceps</i> v. 1-PIRLA	507.00	1.00	0.197
52086 <i>Pinnularia</i> cf. <i>braunii</i> v. <i>amphicephala</i> f. <i>subconica</i>	507.00	2.00	0.394
62022 <i>Stauroneis anceps</i> v. 2-PIRLA	507.00	4.00	0.789
99007 <i>Achnanthes</i> 1-SN	507.00	50.00	9.862
99015 <i>Achnanthes</i> 10-SN	507.00	6.00	1.183
99107 <i>Gomphonema tackei</i> v. <i>abbreviatum</i> Camburn	507.00	2.00	0.394
99137 <i>Navicula mediopunctata</i> Hust.	507.00	38.00	7.495
99144 <i>Navicula</i> 7-SN	507.00	6.00	1.183
99239 <i>Nitzschia</i> 3-SN	507.00	1.00	0.197
99240 <i>Nitzschia</i> 5-SN	507.00	1.00	0.197
99277 <i>Pinnularia</i> 2-SN	507.00	1.00	0.197
99283 <i>Pinnularia</i> 11-SN	507.00	1.00	0.197
99333 <i>Achnanthes lapponica</i> Hust.	507.00	1.00	0.197

LAKE: MOSQUITO 1 LAT DEG: 36 MIN: 25 SEC: 28 LONG DEG: 118 MIN: 37 SEC: 8  
 QUADRANGLE: TRIPLE DIVIDE AREA: 2.07 SHED AREA: 127 ELEV: 2926  
 VEG CLASS: 4 ROCK TYPE: % COVER: A: 2 B: 1 C: 4 D: 1  
 pH FIELD: 7.13 pH LAB: 6.79 CONDUCTIVITY: 7.4 uS@25:  
 HCO<sub>3</sub>: 77.4 Cl: 1.89 NO<sub>3</sub>: 0.4 SO<sub>4</sub>: 7.33 MICR EQUIV/L:  
 Ca: 58.2 Mg: 8.63 Na: 16.8 K: 5.15  
 DATE SAMPLED: 7-21-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 0.6718 GMS PER CC:  
 SED DRY WT: 0.0368 GMS PER CC: INGN LOSS: 41.6

TAXON NAME	Total	No	This Tax	%REL ABUND
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	520.00	10.00	1.923	
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	520.00	3.00	0.577	
8001 <i>Anomoeoneis exilis</i> v. <i>lanceolata</i> A. Mayer	520.00	22.00	4.231	
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.	520.00	28.00	5.385	
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	520.00	1.00	0.192	
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	520.00	6.00	1.154	
23008 <i>Cymbella hebridica</i> Grun. ex Cl. v. <i>hebridica</i>	520.00	5.00	0.962	
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	520.00	76.00	14.615	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	520.00	9.00	1.731	
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	520.00	1.00	0.192	
23016 <i>Cymbella naviculiformis</i> Auersw. ex Heib. v.	520.00	1.00	0.192	
23021 <i>Cymbella</i> 1-PIRLA	520.00	24.00	4.615	
23889 <i>Cymbella</i> spp.	520.00	1.00	0.192	
27001 <i>Diatoma anceps</i> (Ehr.) Kirchn. v. <i>anceps</i>	520.00	3.00	0.577	
27002 <i>Diatoma hemale</i> v. <i>mesodon</i> (Ehr.) Grun.	520.00	1.00	0.192	
33008 <i>Eunotia curvata</i> (Kutz.) Langerst. v. <i>curvata</i>	520.00	1.00	0.192	
33015 <i>Eunotia exigua</i> (Breb. ex Kutz.) Rabh. v. <i>exigua</i>	520.00	2.00	0.385	
33026 <i>Eunotia incisa</i> W. Sm. ex Greg. v. <i>incisa</i>	520.00	3.00	0.577	
33033 <i>Nitzschia</i> 42-SN	520.00	3.00	0.577	
33036 <i>Eunotia naegelii</i> Migula v. <i>naegelii</i>	520.00	1.00	0.192	
33040 <i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.	520.00	7.00	1.346	
33066 <i>Eunotia vanheurckii</i> v. <i>intermedia</i> (Krasske) ex Hust.	520.00	1.00	0.192	
34017 <i>Fragilaria crotensis</i> Kitton v. <i>crotensis</i>	520.00	3.00	0.577	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	520.00	2.00	0.385	
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	520.00	2.00	0.385	
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	520.00	34.00	6.538	
35001 <i>Frustulia rhombooides</i> (Ehr.) DeT. v. <i>rhombooides</i>	520.00	17.00	3.269	
35005 <i>Frustulia rhombooides</i> v. <i>saxonica</i> (Rabh.) DeT.	520.00	6.00	1.154	
37003 <i>Gomphonema angustatum</i> (Kutz.) Rabh.	520.00	9.00	1.731	
44027 <i>Melosira</i> 1-PIRLA	520.00	9.00	1.731	
46038 <i>Navicula mediocris</i> Krasske v. <i>mediocris</i>	520.00	5.00	0.962	
46050 <i>Navicula pseudoscutiformis</i> Hust. v. <i>pseudoscutiformis</i>	520.00	1.00	0.192	
46095 <i>Navicula heimansii</i> van Dam & Kooyman v. <i>heimansii</i>	520.00	1.00	0.192	
46113 <i>Navicula</i> cf. <i>subtilissima</i> v. 2-PIRLA	520.00	19.00	3.654	
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. <i>affine</i>	520.00	1.00	0.192	
47014 <i>Neidium iridis</i> (Ehr.) Cl. v. <i>iridis</i>	520.00	1.00	0.192	
52002 <i>Pinnularia abaujensis</i> v. <i>linearis</i> (Hust.) Patr.	520.00	5.00	0.962	
52011 <i>Pinnularia biceps</i> Greg. v. <i>biceps</i>	520.00	1.00	0.192	
52025 <i>Pinnularia divergens</i> W. Sm. v. <i>divergens</i>	520.00	2.00	0.385	
62015 <i>Stauroneis phoenicenteron</i> (Nitz.) Ehr. v.	520.00	2.00	0.385	
62022 <i>Stauroneis anceps</i> v. 2-PIRLA	520.00	1.00	0.192	
63002 <i>Stenopterobia intermedia</i> (Lewis) V. H. v. <i>intermedia</i>	520.00	4.00	0.769	

TAXON NAME	Total	No	This Tax	%REL ABUND
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	520.00	3.00	0.577	
67002 <i>Tabellaria fenestrata</i> (Lyngb.) Kutz. v. <i>fenestrata</i>	520.00	1.00	0.192	
67005 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain III sensu	520.00	12.00	2.308	
67006 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain IV sensu	520.00	3.00	0.577	
99007 <i>Achnanthes</i> 1-SN	520.00	4.00	0.769	
99008 <i>Achnanthes</i> 2-SN	520.00	1.00	0.192	
99009 <i>Achnanthes</i> 3-SN	520.00	5.00	0.962	
99011 <i>Achnanthes</i> 6-SN	520.00	1.00	0.192	
99050 <i>Cymbella rainierensis</i> Sov.	520.00	1.00	0.192	
99052 <i>Cymbella</i> 1-SN	520.00	4.00	0.769	
99074 <i>Eunotia</i> 1-SN	520.00	2.00	0.385	
99105 <i>Gomphonema puiggarianum</i> v. <i>aequatorialis</i> (Cl.) Camburn	520.00	3.00	0.577	
99124 <i>Krasskella kriegeriana</i> (Krasske) Ross & Sims	520.00	5.00	0.962	
99126 <i>Melosira</i> 1-SN	520.00	65.00	12.500	
99133 <i>Navicula cari</i> Ehr.	520.00	11.00	2.115	
99144 <i>Navicula</i> 7-SN	520.00	2.00	0.385	
99154 <i>Navicula</i> 24-SN	520.00	2.00	0.385	
99231 <i>Nitzschia frustulum</i> 3-SN	520.00	24.00	4.615	
99240 <i>Nitzschia</i> 5-SN	520.00	5.00	0.962	
99281 <i>Pinnularia</i> 9-SN	520.00	30.00	5.769	
99283 <i>Pinnularia</i> 11-SN	520.00	1.00	0.192	
99333 <i>Achnanthes lapponica</i> Hust.	520.00	1.00	0.192	

LAKE: GEM LAT DEG: 37 MIN: 26 SEC: 30 LONG DEG: 118 MIN: 34 SEC: 11  
 QUADRANGLE: MT ABBOT AREA: 2.77 SHED AREA: 1230 ELEV: 3595  
 VEG CLASS: ROCK TYPE: 3 % COVER: A: B: C: D:  
 pH FIELD: 7.33 pH LAB: 6.82 CONDUCTIVITY: 7.4 uS@25:  
 HCO<sub>3</sub>: 77.4 Cl: 1.89 NO<sub>3</sub>: 0.4 SO<sub>4</sub>: 7.33 MICR EQUIV/L:  
 Ca: 58.2 Mg: 8.63 Na : 16.8 K : 4.14  
 DATE SAMPLED: 7-21-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.6628 GMS PER CC:  
 SED DRY WT: 1.0757 GMS PER CC: INGN LOSS: 3

TAXON NAME	Total	No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	512.00	4.00	0.781	
2004 <i>Achnanthes clevei</i> Grun. v. <i>clevei</i>	512.00	2.00	0.391	
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	512.00	2.00	0.391	
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	512.00	18.00	3.516	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	512.00	8.00	1.563	
7002 <i>Amphora ovalis</i> v. <i>affinis</i> (Kutz.) V. H. ex Det.	512.00	1.00	0.195	
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. <i>perpusilla</i>	512.00	26.00	5.078	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	512.00	1.00	0.195	
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	512.00	17.00	3.320	
30001 <i>Diploneis elliptica</i> (Kutz.) Cl. v. <i>elliptica</i>	512.00	2.00	0.391	
30003 <i>Diploneis marginestriata</i> Hust. v. <i>marginestriata</i>	512.00	9.00	1.758	
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	512.00	5.00	0.977	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	512.00	16.00	3.125	
34889 <i>Fragilaria</i> spp.	512.00	2.00	0.391	
37003 <i>Gomphonema angustatum</i> (Kutz.) Rabh.	512.00	1.00	0.195	
46095 <i>Navicula heimansii</i> van Dam & Kooyman v. <i>heimansii</i>	512.00	1.00	0.195	
65014 <i>Surirella linearis</i> W. Sm. v. <i>linearis</i>	512.00	1.00	0.195	
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	512.00	42.00	8.203	
99004 <i>Achnanthes lanceolata</i> f. <i>capitata</i> O. Mull.	512.00	4.00	0.781	
99006 <i>Achnanthes suchlandi</i> Hust.	512.00	16.00	3.125	
99007 <i>Achnanthes</i> 1-SN	512.00	23.00	4.492	
99015 <i>Achnanthes</i> 10-SN	512.00	6.00	1.172	
99026 <i>Achnanthes</i> 32-SN	512.00	1.00	0.195	
99062 <i>Cymbella</i> 11-SN	512.00	1.00	0.195	
99090 <i>Fragilaria</i> 7-SN	512.00	3.00	0.586	
99094 <i>Fragilaria</i> 11-SN	512.00	58.00	11.328	
99107 <i>Gomphonema</i> <i>tackei</i> v. <i>abbreviatum</i> Camburn	512.00	33.00	6.445	
99119 <i>Gomphonema</i> 13-SN	512.00	15.00	2.930	
99123 <i>Hannea arcus</i> (Ehr.) Pat.	512.00	1.00	0.195	
99137 <i>Navicula</i> <i>mediopunctata</i> Hust.	512.00	8.00	1.563	
99138 <i>Navicula</i> <i>pseudolanceolata</i> Lange-Bertalot	512.00	3.00	0.586	
99146 <i>Navicula</i> 10-SN	512.00	1.00	0.195	
99153 <i>Navicula</i> 22-SN	512.00	1.00	0.195	
99154 <i>Navicula</i> 24-SN	512.00	2.00	0.391	
99161 <i>Navicula</i> 30-SN	512.00	34.00	6.641	
99162 <i>Navicula</i> 31-SN	512.00	14.00	2.734	
99166 <i>Navicula</i> 37-SN	512.00	2.00	0.391	
99179 <i>Navicula</i> 50-SN	512.00	57.00	11.133	
99181 <i>Navicula</i> 52-SN	512.00	3.00	0.586	
99185 <i>Navicula</i> 56-SN	512.00	7.00	1.367	
99200 <i>Navicula</i> 71-SN	512.00	2.00	0.391	
99201 <i>Navicula</i> 72-SN	512.00	44.00	8.594	

TAXON NAME	Total	No	This Tax	%REL ABUND
99231 <i>Nitzschia frustulum</i> 3-SN	512.00	1.00		0.195
99233 <i>Nitzschia frustulum</i> 5-SN	512.00	2.00		0.391
99252 <i>Nitzschia</i> 17-SN	512.00	3.00		0.586
99261 <i>Nitzschia</i> 30-SN	512.00	4.00		0.781
99268 <i>Nitzschia</i> 37-SN	512.00	3.00		0.586
99310 <i>Pinnularia</i> 40-SN	512.00	2.00		0.391

LAKE: GRANITE LAT DEG: 36 MIN: 51 SEC: 47 LONG DEG: 118 MIN: 37 SEC: 12  
 QUADRANGLE: TUOL.MEADOWS AREA: 20.7 SHED AREA: 262 ELEV: 3097  
 VEG CLASS: 3 ROCK TYPE: % COVER: A: 1 B: 1 C: 1 D: 4  
 pH FIELD: 6.91 pH LAB: 6.91 CONDUCTIVITY: 6.9 uS@25:  
 HCO<sub>3</sub>: 66.9 Cl: 1.29 NO<sub>3</sub>: 0 SO<sub>4</sub>: 4.65 MICR EQUIV/L:  
 Ca: 49.6 Mg: 6.68 Na : 15.6 K : 4.09  
 DATE SAMPLED: 7-5-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.3143 GMS PER CC:  
 SED DRY WT: 0.0321 GMS PER CC: INGN LOSS: 22.9

TAXON NAME	Total	No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	511.00	2.00	0.391	
2023 <i>Achnanthes levanderi</i> v. <i>helvetica</i> Hust.	511.00	1.00	0.196	
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	511.00	2.00	0.391	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	511.00	22.00	4.305	
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	511.00	1.00	0.196	
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i> )	511.00	85.00	16.634	
20012 <i>Cyclotella pseudostelligera</i> Hust. v. <i>pseudostelligera</i>	511.00	1.00	0.196	
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. <i>cesatii</i>	511.00	3.00	0.587	
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	511.00	2.00	0.391	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	511.00	10.00	1.957	
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	511.00	24.00	4.697	
23031 <i>Cymbella muelleri</i> Hust. v. <i>muelleri</i>	511.00	1.00	0.196	
34003 <i>Fragilaria brevistriata</i> Grun. v. <i>brevistriata</i>	511.00	20.00	3.914	
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	511.00	14.00	2.740	
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	511.00	2.00	0.391	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	511.00	24.00	4.697	
34027 <i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.	511.00	1.00	0.196	
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	511.00	1.00	0.196	
34038 <i>Fragilaria pinnata</i> v. <i>acuminata</i> A. Mayer	511.00	71.00	13.894	
44001 <i>Melosira ambigua</i> (Grun.) O. Mull. v. <i>ambigua</i>	511.00	5.00	0.978	
44010 <i>Melosira italicica</i> (Ehr.) Kutz. v. <i>italicica</i>	511.00	6.00	1.174	
44011 <i>Melosira italicica</i> ssp. <i>subarctica</i> O. Mull.	511.00	4.00	0.783	
44027 <i>Melosira</i> 1-PIRLA	511.00	4.00	0.783	
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	511.00	5.00	0.978	
46050 <i>Navicula pseudoscutiformis</i> Hust. v. <i>pseudoscutiformis</i>	511.00	1.00	0.196	
46133 <i>Navicula</i> 14-PIRLA	511.00	36.00	7.045	
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. <i>affine</i>	511.00	1.00	0.196	
47008 <i>Neidium bisulcatum</i> v. <i>baicalense</i> (Skv. & Meyer)	511.00	3.00	0.587	
52002 <i>Pinnularia abaujensis</i> v. <i>linearis</i> (Hust.) Patr.	511.00	1.00	0.196	
52025 <i>Pinnularia divergens</i> W. Sm. v. <i>divergens</i>	511.00	1.00	0.196	
52078 <i>Pinnularia abaujensis</i> v. 2-PIRLA	511.00	1.00	0.196	
62003 <i>Stauroneis anceps</i> f. <i>gracilis</i> Rabh.	511.00	1.00	0.196	
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	511.00	1.00	0.196	
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	511.00	1.00	0.196	
99007 <i>Achnanthes</i> 1-SN	511.00	65.00	12.720	
99009 <i>Achnanthes</i> 3-SN	511.00	11.00	2.153	
99105 <i>Gomphonema puiggarianum</i> v. <i>aequatorialis</i> (Cl.) Camburn	511.00	8.00	1.566	
99126 <i>Melosira</i> 1-SN	511.00	6.00	1.174	
99133 <i>Navicula cari</i> Ehr.	511.00	1.00	0.196	
99137 <i>Navicula mediopunctata</i> Hust.	511.00	9.00	1.761	
99138 <i>Navicula pseudolanceolata</i> Lange-Bertalot	511.00	5.00	0.978	
99142 <i>Navicula</i> 1-SN	511.00	11.00	2.153	

TAXON NAME	Total	No	This Tax	%REL ABUND
99146 Navicula 10-SN	511.00		1.00	0.196
99147 Navicula 11-SN	511.00		1.00	0.196
99155 Navicula 25-SN	511.00		2.00	0.391
99163 Navicula 32-SN	511.00		7.00	1.370
99228 Nitzschia dissipata v. undulata Sovereign	511.00		1.00	0.196
99230 Nitzschia frustulum 2-SN	511.00		2.00	0.391
99231 Nitzschia frustulum 3-SN	511.00		1.00	0.196
99239 Nitzschia 3-SN	511.00		1.00	0.196
99240 Nitzschia 5-SN	511.00		1.00	0.196
99242 Nitzschia 7-SN	511.00		3.00	0.587
99249 Nitzschia 14-SN	511.00		1.00	0.196
99250 Nitzschia 15-SN	511.00		1.00	0.196
99277 Pinnularia 2-SN	511.00		2.00	0.391
99281 Pinnularia 9-SN	511.00		8.00	1.566
99283 Pinnularia 11-SN	511.00		1.00	0.196
99330 Synedra 2-SN	511.00		3.00	0.587
99335 Gomphonema ventricosum	511.00		1.00	0.196

LAKE: HEATHER/TAH. LAT DEG: 38 MIN: 52 SEC: 40 LONG DEG: 120 MIN: 8 SEC: 10  
 QUADRANGLE: FALLEN LEAF AREA: SHED AREA: ELEV:  
 VEG CLASS: ROCK TYPE: % COVER: A: B: C: D:  
 pH FIELD: 6.95 pH LAB: 6.93 CONDUCTIVITY: 8.1 USA25:  
 HCO<sub>3</sub>: 67.1 Cl: 2.08 NO<sub>3</sub>: 0 SO<sub>4</sub>: 12.1 MICR EQUIV/L:  
 Ca: 50.2 Mg: 5.82 Na : 12.1 K : 5.9  
 DATE SAMPLED: 8-22-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 0.9953 GMS PER CC:  
 SED DRY WT: 0.0576 GMS PER CC: INGN LOSS: 31.8

TAXON NAME	Total No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. austriaca	513.00	2.00	0.390
2003 <i>Achnanthes austriaca</i> Hust. v. helvetica	513.00	2.00	0.390
2026 <i>Achnanthes linearis</i> v. <i>pusilla</i> Grun.	513.00	41.00	7.992
2028 <i>Achnanthes marginulata</i> Grun. v. marginulata	513.00	10.00	1.949
2029 <i>Achnanthes microcephala</i> (Kutz.) Grun. v. microcephala	513.00	47.00	9.162
2030 <i>Achnanthes minutissima</i> Kutz. v. minutissima	513.00	13.00	2.534
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. detha	513.00	5.00	0.975
8001 <i>Anomoeoneis exilis</i> v. lanceolata A. Mayer	513.00	50.00	9.747
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.)	513.00	9.00	1.754
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. stelligera	513.00	76.00	14.815
20012 <i>Cyclotella pseudostelligera</i> Hust. v. pseudostelligera	513.00	1.00	0.195
23007 <i>Cymbella gaeumannii</i> Meist. v. gaeumannii	513.00	4.00	0.780
23009 <i>Cymbella lunata</i> W. Sm. v. lunata	513.00	14.00	2.729
23010 <i>Cymbella microcephala</i> Grun. v. microcephala	513.00	7.00	1.365
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. minuta	513.00	2.00	0.390
23021 <i>Cymbella</i> 1-PIRLA	513.00	1.00	0.195
33008 <i>Eunotia curvata</i> (Kutz.) Langerst. v. curvata	513.00	3.00	0.585
33010 <i>Eunotia curvata</i> v. <i>subarcuata</i> (Naeg.) Woodhead & Tweed	513.00	2.00	0.390
33015 <i>Eunotia exigua</i> (Breb. ex Kutz.) Rabh. v. exigua	513.00	2.00	0.390
33019 <i>Eunotia flexuosa</i> Breb. ex Kutz. v. flexuosa	513.00	18.00	3.509
33036 <i>Eunotia naegelii</i> Migula v. naegelii	513.00	9.00	1.754
33065 <i>Eunotia vanheurckii</i> Pat. v. vanheurckii	513.00	15.00	2.924
34017 <i>Fragilaria crotonensis</i> Kitton v. crotonensis	513.00	6.00	1.170
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	513.00	1.00	0.195
34037 <i>Fragilaria virescens</i> v. exigua Grun.	513.00	2.00	0.390
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. rhomboides	513.00	6.00	1.170
35003 <i>Frustulia rhomboides</i> v. <i>capitata</i> (A. Mayer) Patr.	513.00	3.00	0.585
35005 <i>Frustulia rhomboides</i> v. <i>saxonica</i> (Rabh.) DeT.	513.00	6.00	1.170
37003 <i>Gomphonema angustatum</i> (Kutz.) Rabh.	513.00	5.00	0.975
37014 <i>Gomphonema truncatum</i> v. <i>capitatum</i> (Ehr.) Patr.	513.00	1.00	0.195
37889 <i>Gomphonema</i> spp.	513.00	2.00	0.390
44011 <i>Melosira italica</i> ssp. <i>subarctica</i> O. Mull.	513.00	2.00	0.390
44013 <i>Melosira italica</i> v. <i>valida</i> (Grun.) Hust.	513.00	1.00	0.195
44027 <i>Melosira</i> 1-PIRLA	513.00	7.00	1.365
46095 <i>Navicula heimansii</i> van Dam & Kooyman v. heimansii	513.00	2.00	0.390
46113 <i>Navicula</i> cf. <i>subtilissima</i> v. 2-PIRLA	513.00	2.00	0.390
46123 <i>Navicula</i> 23-PIRLA	513.00	4.00	0.780
46133 <i>Navicula</i> 14-PIRLA	513.00	2.00	0.390
46889 <i>Navicula</i> spp.	513.00	4.00	0.780
47001 <i>Neidium affine</i> (Ehr.) Pfitz. v. affine	513.00	1.00	0.195
47007 <i>Neidium bisulcatum</i> (Lagerst.) Cl. v. bisulcatum	513.00	2.00	0.390
48008 <i>Nitzschia dissipata</i> (Kutz.) Grun. v. dissipata	513.00	1.00	0.195

TAXON NAME	Total No	This Tax	%REL ABUND
52001 <i>Pinnularia abaujensis</i> (Pant.) Ross v. <i>abaujensis</i>	513.00	1.00	0.195
52025 <i>Pinnularia divergens</i> W. Sm. v. <i>divergens</i>	513.00	2.00	0.390
52038 <i>Pinnularia maior</i> (Kutz.) Rabh. v. <i>maior</i>	513.00	1.00	0.195
52074 <i>Pinnularia biceps</i> v. 1-PIRLA	513.00	3.00	0.585
63002 <i>Stenopterobia intermedia</i> (Lewis) V. H. v. <i>intermedia</i>	513.00	5.00	0.975
65011 <i>Surirella delicatissima</i> Lewis v. <i>delicatissima</i>	513.00	3.00	0.585
66016 <i>Synedra rumpens</i> Kutz. v. <i>rumpens</i>	513.00	5.00	0.975
67005 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain III sensu	513.00	4.00	0.780
67006 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain IV sensu	513.00	2.00	0.390
99007 <i>Achnanthes</i> 1-SN	513.00	1.00	0.195
99009 <i>Achnanthes</i> 3-SN	513.00	19.00	3.704
99013 <i>Achnanthes</i> 8-SN	513.00	2.00	0.390
99025 <i>Achnanthes</i> 31-SN	513.00	1.00	0.195
99035 <i>Anomoeoneis serians</i> v. 1-SN	513.00	1.00	0.195
99094 <i>Fragilaria</i> 11-SN	513.00	3.00	0.585
99102 <i>Frustulia rhomboides</i> v. 1-SN	513.00	2.00	0.390
99119 <i>Gomphonema</i> 13-SN	513.00	3.00	0.585
99120 <i>Gomphonema</i> 14-SN	513.00	2.00	0.390
99123 <i>Hannea arcus</i> (Ehr.) Pat.	513.00	1.00	0.195
99124 <i>Krasskella kriegeriana</i> (Krasske) Ross & Sims	513.00	4.00	0.780
99133 <i>Navicula cari</i> Ehr.	513.00	2.00	0.390
99142 <i>Navicula</i> 1-SN	513.00	1.00	0.195
99143 <i>Navicula</i> 6-SN	513.00	2.00	0.390
99175 <i>Navicula</i> 46-SN	513.00	1.00	0.195
99194 <i>Navicula</i> 65-SN	513.00	1.00	0.195
99211 <i>Navicula</i> 82-SN	513.00	2.00	0.390
99214 <i>Navicula</i> 85-SN	513.00	2.00	0.390
99228 <i>Nitzschia dissipata</i> v. <i>undulata</i> Sovereign	513.00	1.00	0.195
99229 <i>Nitzschia frustulum</i> 1-SN	513.00	4.00	0.780
99237 <i>Nitzschia</i> 1-SN	513.00	2.00	0.390
99240 <i>Nitzschia</i> 5-SN	513.00	2.00	0.390
99241 <i>Nitzschia</i> 6-SN	513.00	1.00	0.195
99242 <i>Nitzschia</i> 7-SN	513.00	3.00	0.585
99269 <i>Nitzschia</i> 38-SN	513.00	3.00	0.585
99270 <i>Nitzschia</i> 39-SN	513.00	1.00	0.195
99277 <i>Pinnularia</i> 2-SN	513.00	1.00	0.195
99281 <i>Pinnularia</i> 9-SN	513.00	11.00	2.144
99303 <i>Pinnularia</i> 32-SN	513.00	1.00	0.195
99330 <i>Synedra</i> 2-SN	513.00	7.00	1.365
99350 <i>Cymbella</i> 13-SN	513.00	7.00	1.365

LAKE: RUBY LAT DEG: 37 MIN: 24 SEC: 50 LONG DEG: 118 MIN: 46 SEC: 15  
 QUADRANGLE: MT.ABBOT AREA: 14.5 SHED AREA: 202 ELEV: 3365  
 VEG CLASS: 3 ROCK TYPE: % COVER: A: 1 B: 1 C: 1 D: 4  
 pH FIELD: 7.05 pH LAB: 6.95 CONDUCTIVITY: 5.5 usa25:  
 HCO<sub>3</sub>: 47.3 Cl: 1.65 NO<sub>3</sub>: 2.88 SO<sub>4</sub>: 7.04 MICR EQUIV/L:  
 Ca: 43.2 Mg: 3.72 Na : 9.01 K : 3.45  
 DATE SAMPLED: 8-11-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.2904 GMS PER CC:  
 SED DRY WT: 0.0835 GMS PER CC: INGN LOSS: 14.3

TAXON NAME	Total No	This Tax	%REL ABUND
2006 <i>Achnanthes didyma</i> Hust. v. <i>didyma</i>	511.00	2.00	0.391
2026 <i>Achnanthes linearis</i> v. <i>pusilla</i> Grun.	511.00	2.00	0.391
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	511.00	29.00	5.675
2029 <i>Achnanthes microcephala</i> (Kutz.) Grun. v. <i>microcephala</i>	511.00	1.00	0.196
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	511.00	8.00	1.566
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	511.00	40.00	7.828
2048 <i>Achnanthes laterostrata</i> Hust. v. <i>laterostrata</i>	511.00	1.00	0.196
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	511.00	34.00	6.654
9002 <i>Asterionella ralfsii</i> v. <i>americana</i> Korn.	511.00	1.00	0.196
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	511.00	140.00	27.397
20012 <i>Cyclotella pseudostelligera</i> Hust. v. <i>pseudostelligera</i>	511.00	46.00	9.002
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. <i>cesatii</i>	511.00	2.00	0.391
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	511.00	1.00	0.196
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	511.00	11.00	2.153
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	511.00	2.00	0.391
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	511.00	3.00	0.587
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. <i>rhomboides</i>	511.00	1.00	0.196
35005 <i>Frustulia rhomboides</i> v. <i>saxonica</i> (Rabh.) DeT.	511.00	2.00	0.391
44011 <i>Melosira italica</i> ssp. <i>subarctica</i> O. Mull.	511.00	2.00	0.391
44013 <i>Melosira italica</i> v. <i>valida</i> (Grun.) Hust.	511.00	6.00	1.174
44014 <i>Melosira lirata</i> (Ehr.) Kutz. v. <i>lirata</i>	511.00	8.00	1.566
44027 <i>Melosira</i> 1-PIRLA	511.00	14.00	2.740
44030 <i>Melosira distans</i> v. <i>nivalis</i> (W. Sm.) Kirchn.	511.00	4.00	0.783
45001 <i>Meridion circulare</i> (Grev.) Agardh v. <i>circulare</i>	511.00	1.00	0.196
46002 <i>Navicula angusta</i> Grun. v. <i>angusta</i>	511.00	1.00	0.196
46050 <i>Navicula pseudoscutiformis</i> Hust. v. <i>pseudoscutiformis</i>	511.00	2.00	0.391
46102 <i>Navicula mutica</i> v. <i>cohnii</i> (Hilse) Grun.	511.00	1.00	0.196
46133 <i>Navicula</i> 14-PIRLA	511.00	4.00	0.783
52027 <i>Pinnularia divergentissima</i> (Grun.) Cl. v.	511.00	1.00	0.196
52078 <i>Pinnularia abaujensis</i> v. 2-PIRLA	511.00	1.00	0.196
63002 <i>Stenopterobia intermedia</i> (Lewis) V. H. v. <i>intermedia</i>	511.00	1.00	0.196
99001 <i>Achnanthes bicapitata</i> Hust.	511.00	1.00	0.196
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	511.00	19.00	3.718
99007 <i>Achnanthes</i> 1-SN	511.00	43.00	8.415
99009 <i>Achnanthes</i> 3-SN	511.00	1.00	0.196
99012 <i>Achnanthes</i> 7-SN	511.00	1.00	0.196
99015 <i>Achnanthes</i> 10-SN	511.00	1.00	0.196
99016 <i>Achnanthes</i> 11-SN	511.00	1.00	0.196
99090 <i>Fragilaria</i> 7-SN	511.00	9.00	1.761
99105 <i>Gomphonema puiggarianum</i> v. <i>aequatorialis</i> (Cl.) Camburn	511.00	5.00	0.978
99107 <i>Gomphonema tackei</i> v. <i>abbreviatum</i> Camburn	511.00	7.00	1.370
99137 <i>Navicula mediopunctata</i> Hust.	511.00	23.00	4.501

TAXON NAME	Total	No	This Tax	%REL ABUND
99142 Navicula 1-SN	511.00	4.00	0.783	
99146 Navicula 10-SN	511.00	1.00	0.196	
99148 Navicula 13-SN	511.00	3.00	0.587	
99150 Navicula 16-SN	511.00	2.00	0.391	
99151 Navicula 18-SN	511.00	1.00	0.196	
99152 Navicula 21-SN	511.00	1.00	0.196	
99172 Navicula 43-SN	511.00	1.00	0.196	
99228 Nitzschia dissipata v. undulata Sovereign	511.00	1.00	0.196	
99239 Nitzschia 3-SN	511.00	2.00	0.391	
99283 Pinnularia 11-SN	511.00	3.00	0.587	
99284 Pinnularia 12-SN	511.00	1.00	0.196	
99330 Synedra 2-SN	511.00	8.00	1.566	

LAKE: UPPER ANGORA LAT DEG: 38 MIN: 51 SEC: 46 LONG DEG: 120 MIN: 3 SEC: 59  
 QUADRANGLE: FALLEN LEAF AREA: 5.44 SHED AREA: 65.3 ELEV: 2292  
 VEG CLASS: 4 ROCK TYPE: % COVER: A: 1 B: 1 C: 4 D: 1  
 pH FIELD: 7 pH LAB: 6.98 CONDUCTIVITY: 9.9 USA25:  
 HCO<sub>3</sub>: 88.5 Cl: 4.95 NO<sub>3</sub>: 0 SO<sub>4</sub>: 1.7 MICR EQUIV/L:  
 Ca: 42.2 Mg: 12.3 Na: 34.3 K: 7.53  
 DATE SAMPLED: 8-22-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.1692 GMS PER CC:  
 SED DRY WT: 0.0603 GMS PER CC: INGN LOSS: 16.1

TAXON NAME	Total No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	507.00	4.00	0.789
2026 <i>Achnanthes linearis</i> v. <i>pusilla</i> Grun.	507.00	1.00	0.197
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	507.00	2.00	0.394
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	507.00	16.00	3.156
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	507.00	1.00	0.197
8001 <i>Anomoeoneis exilis</i> v. <i>lanceolata</i> A. Mayer	507.00	29.00	5.720
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.	507.00	29.00	5.720
16001 <i>Coccinea diminuta</i> Pant. v. <i>diminuta</i>	507.00	1.00	0.197
20012 <i>Cyclotella pseudostelligera</i> Hust. v. <i>pseudostelligera</i>	507.00	42.00	8.284
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	507.00	8.00	1.578
23008 <i>Cymbella hebridica</i> Grun. ex Cl. v. <i>hebridica</i>	507.00	1.00	0.197
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	507.00	40.00	7.890
23016 <i>Cymbella naviculiformis</i> Auersw. ex Heib. v.	507.00	2.00	0.394
33008 <i>Eunotia curvata</i> (Kutz.) Langerst. v. <i>curvata</i>	507.00	8.00	1.578
33010 <i>Eunotia curvata</i> v. <i>subarcuata</i> (Naeg.) Woodhead & Tweed	507.00	7.00	1.381
33011 <i>Eunotia denticulata</i> (Breb.) Rabh. v. <i>denticulata</i>	507.00	1.00	0.197
33015 <i>Eunotia exigua</i> (Breb. ex Kutz.) Rabh. v. <i>exigua</i>	507.00	2.00	0.394
33018 <i>Eunotia fallax</i> Cl.-Eul. v. <i>fallax</i>	507.00	1.00	0.197
33019 <i>Eunotia flexuosa</i> Breb. ex Kutz. v. <i>flexuosa</i>	507.00	1.00	0.197
33021 <i>Eunotia formica</i> Ehr. v. <i>formica</i>	507.00	1.00	0.197
33026 <i>Eunotia incisa</i> W. Sm. ex Greg. v. <i>incisa</i>	507.00	1.00	0.197
33036 <i>Eunotia naegelii</i> Migula v. <i>naegelii</i>	507.00	3.00	0.592
33039 <i>Eunotia pectinalis</i> (O. F. Mull.?) Rabh. v. <i>pectinalis</i>	507.00	2.00	0.394
33065 <i>Eunotia vanheurckii</i> Pat. v. <i>vanheurckii</i>	507.00	4.00	0.789
33066 <i>Eunotia vanheurckii</i> v. <i>intermedia</i> (Krasske) ex Hust.	507.00	2.00	0.394
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	507.00	5.00	0.986
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	507.00	3.00	0.592
35001 <i>Frustulia rhombooides</i> (Ehr.) DeT. v. <i>rhombooides</i>	507.00	11.00	2.170
35002 <i>Frustulia rhombooides</i> v. <i>amphipleurooides</i> (Grun.) Pet.	507.00	1.00	0.197
35005 <i>Frustulia rhombooides</i> v. <i>saxonica</i> (Rabh.) DeT.	507.00	3.00	0.592
37003 <i>Gomphonema angustatum</i> (Kutz.) Rabh.	507.00	12.00	2.367
37889 <i>Gomphonema</i> spp.	507.00	2.00	0.394
44008 <i>Melosira islandica</i> O. Mull. v. <i>islandica</i>	507.00	6.00	1.183
44011 <i>Melosira italica</i> ssp. <i>subarctica</i> O. Mull.	507.00	3.00	0.592
44027 <i>Melosira</i> 1-PIRLA	507.00	18.00	3.550
46008 <i>Navicula bremensis</i> Hust.	507.00	1.00	0.197
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	507.00	3.00	0.592
46038 <i>Navicula mediocris</i> Krasske v. <i>mediocris</i>	507.00	10.00	1.972
46051 <i>Navicula pupula</i> Kutz. v. <i>pupula</i>	507.00	2.00	0.394
46056 <i>Navicula radiosa</i> Kutz. v. <i>radiosa</i>	507.00	3.00	0.592
46057 <i>Navicula radiosa</i> v. <i>parva</i> Wallace	507.00	10.00	1.972
46121 <i>Navicula</i> cf. <i>subtilissima</i> v. 4-PIRLA	507.00	1.00	0.197

TAXON NAME	Total No	This Tax	%REL ABUND
47001 Neidium affine (Ehr.) Pfitz. v. affine	507.00	12.00	2.367
47002 Neidium affine v. amphirhynchus (Ehr.) Cl. ACP	507.00	1.00	0.197
47007 Neidium bisulcatum (Lagerst.) Cl. v. bisulcatum	507.00	1.00	0.197
47014 Neidium iridis (Ehr.) Cl. v. iridis	507.00	4.00	0.789
47025 Neidium 2-PIRLA	507.00	3.00	0.592
48008 Nitzschia dissipata (Kutz.) Grun. v. dissipata	507.00	2.00	0.394
52002 Pinnularia abaujensis v. linearis (Hust.) Patr.	507.00	6.00	1.183
52038 Pinnularia maior (Kutz.) Rabh. v. maior	507.00	3.00	0.592
52080 Pinnularia cf. pseudomicrostauron Gandhi v.	507.00	5.00	0.986
62015 Stauroeis phoenicenteron (Nitz.) Ehr. v.	507.00	4.00	0.789
62022 Stauroeis anceps v. 2-PIRLA	507.00	7.00	1.381
63002 Stenopterobia intermedia (Lewis) V. H. v. intermedia	507.00	5.00	0.986
63003 Stenopterobia anceps (Lewis) Breb. ex V. H. v. anceps	507.00	1.00	0.197
65011 Surirella delicatissima Lewis v. delicatissima	507.00	3.00	0.592
65014 Surirella linearis W. Sm. v. linearis	507.00	1.00	0.197
66016 Synedra rumpens Kutz. v. rumpens	507.00	3.00	0.592
66029 Synedra 1-PIRLA	507.00	3.00	0.592
67005 Tabellaria flocculosa Roth (Kutz.) strain III sensu	507.00	4.00	0.789
90990 Nitzschia 42-SN	507.00	4.00	0.789
99007 Achnanthes 1-SN	507.00	1.00	0.197
99009 Achnanthes 3-SN	507.00	1.00	0.197
99013 Achnanthes 8-SN	507.00	2.00	0.394
99084 Eunotia 14-SN	507.00	1.00	0.197
99098 Fragilaria 16-SN	507.00	11.00	2.170
99102 Frustulia rhomboides v. l-SN	507.00	3.00	0.592
99103 Gomphonema affine v. insigne (Greg.) Andrews	507.00	2.00	0.394
99105 Gomphonema puiggarianum v. aequatorialis (Cl.) Camburn	507.00	19.00	3.748
99107 Gomphonema tackei v. abbreviatum Camburn	507.00	1.00	0.197
99124 Krasskella kriegeriana (Krasske) Ross & Sims	507.00	4.00	0.789
99126 Melosira 1-SN	507.00	12.00	2.367
99128 Melosira 3-SN	507.00	2.00	0.394
99130 Melosira 5-SN	507.00	1.00	0.197
99143 Navicula 6-SN	507.00	1.00	0.197
99144 Navicula 7-SN	507.00	2.00	0.394
99226 Nitzschia acuta Hantzsch	507.00	1.00	0.197
99228 Nitzschia dissipata v. undulata Sovereign	507.00	2.00	0.394
99231 Nitzschia frustulum 3-SN	507.00	1.00	0.197
99236 Nitzschia frustulum 8-SN	507.00	8.00	1.578
99240 Nitzschia 5-SN	507.00	9.00	1.775
99268 Nitzschia 37-SN	507.00	3.00	0.592
99281 Pinnularia 9-SN	507.00	38.00	7.495
99299 Pinnularia 28-SN	507.00	3.00	0.592
99330 Synedra 2-SN	507.00	1.00	0.197
99334 Achnanthes recurvata Hust.	507.00	1.00	0.197
99348 Gomphonema truncatum Ehr. v. truncatum	507.00	2.00	0.394

LAKE: DOROTHY LAT DEG: 37 MIN: 32 SEC: 10 LONG DEG: 118 MIN: 52 SEC: 55  
 QUADRANGLE: MT MORRISON AREA: 61.6 SHED AREA: 396 ELEV: 3121  
 VEG CLASS: 3 ROCK TYPE: % COVER: A: 1 B: 3 C: 3 D: 1  
 pH FIELD: 7.46 pH LAB: 7.2 CONDUCTIVITY: 22.9 USA25:  
 HCO<sub>3</sub>: 138.5 Cl: 2.66 NO<sub>3</sub>: 0.72 SO<sub>4</sub>: 83.2 MICR EQUIV/L:  
 Ca: 189.6 Mg: 21.3 Na: 18 K: 7.64  
 DATE SAMPLED: 8-20-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.3673 GMS PER CC:  
 SED DRY WT: 0.2802 GMS PER CC: INGN LOSS: 6.24

TAXON NAME	Total No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. austriaca	519.00	1.00	0.193
2006 <i>Achnanthes didyma</i> Hust. v. didyma	519.00	4.00	0.771
2023 <i>Achnanthes levanderi</i> v. <i>helvetica</i> Hust.	519.00	5.00	0.963
2024 <i>Achnanthes linearis</i> (W. Sm.) Grun. v. linearis	519.00	22.00	4.239
2028 <i>Achnanthes marginulata</i> Grun. v. marginulata	519.00	17.00	3.276
2029 <i>Achnanthes microcephala</i> (Kutz.) Grun. v. microcephala	519.00	1.00	0.193
2048 <i>Achnanthes laterostrata</i> Hust. v. laterostrata	519.00	6.00	1.156
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. perpusilla	519.00	1.00	0.193
8001 <i>Anomoeoneis exilis</i> v. lanceolata A. Mayer	519.00	1.00	0.193
20009 <i>Cyclotella ocellata</i> Pant. v. ocellata	519.00	14.00	2.697
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. stelligera	519.00	296.00	57.033
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. cesatii	519.00	1.00	0.193
23010 <i>Cymbella microcephala</i> Grun. v. microcephala	519.00	1.00	0.193
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. minuta	519.00	1.00	0.193
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	519.00	2.00	0.385
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	519.00	4.00	0.771
34025 <i>Fragilaria pinnata</i> Ehr. v. pinnata	519.00	21.00	4.046
34027 <i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.	519.00	4.00	0.771
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. <i>rhomboides</i>	519.00	1.00	0.193
44013 <i>Melosira italica</i> v. <i>valida</i> (Grun.) Hust.	519.00	2.00	0.385
44027 <i>Melosira</i> 1-PIRLA	519.00	2.00	0.385
47002 <i>Neidium affine</i> v. <i>amphirhynchus</i> (Ehr.) Cl. ACP	519.00	2.00	0.385
48008 <i>Nitzschia dissipata</i> (Kutz.) Grun. v. <i>dissipata</i>	519.00	2.00	0.385
65014 <i>Surirella linearis</i> W. Sm. v. <i>linearis</i>	519.00	2.00	0.385
66014 <i>Synedra parasitica</i> (W. Sm.) Hust. v. <i>parasitica</i>	519.00	1.00	0.193
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	519.00	7.00	1.349
99006 <i>Achnanthes suchlandi</i> Hust.	519.00	23.00	4.432
99007 <i>Achnanthes</i> 1-SN	519.00	11.00	2.119
99013 <i>Achnanthes</i> 8-SN	519.00	1.00	0.193
99015 <i>Achnanthes</i> 10-SN	519.00	2.00	0.385
99021 <i>Achnanthes</i> 21-SN	519.00	3.00	0.578
99026 <i>Achnanthes</i> 32-SN	519.00	10.00	1.927
99027 <i>Achnanthes</i> 34-SN	519.00	2.00	0.770
99035 <i>Anomoeoneis serians</i> v. 1-SN	519.00	3.00	0.578
99042 <i>Cyclotella</i> 1-SN	519.00	1.00	0.193
99051 <i>Cymbella sinuata</i> Greg.	519.00	2.00	0.385
99061 <i>Cymbella</i> 10-SN	519.00	3.00	0.578
99088 <i>Fragilaria</i> 2-SN	519.00	16.00	3.083
99094 <i>Fragilaria</i> 11-SN	519.00	7.00	1.349
99095 <i>Fragilaria</i> 13-SN	519.00	1.00	0.193
99127 <i>Melosira</i> 2-SN	519.00	1.00	0.193
99137 <i>Navicula mediopunctata</i> Hust.	519.00	1.00	0.193

TAXON NAME	Total No	This Tax	%REL ABUND
99138 Navicula pseudolanceolata Lange-Bertalot	519.00	3.00	0.578
99172 Navicula 43-SN	519.00	1.00	0.193
99186 Navicula 57-SN	519.00	2.00	0.385
99229 Nitzschia frustulum 1-SN	519.00	1.00	0.193
99231 Nitzschia frustulum 3-SN	519.00	2.00	0.385

LAKE: EASTERN BROOK LAT DEG: MIN: SEC: LONG DEG: MIN: SEC:  
 QUADRANGLE: MT TOM AREA: SHED AREA: ELEV:  
 VEG CLASS: ROCK TYPE: % COVER: A: B: C: D:  
 pH FIELD: 7.35 pH LAB: 7.22 CONDUCTIVITY: 12 uS/cm:  
 HCO<sub>3</sub>: 132.2 Cl: 4.7 NO<sub>3</sub>: 0 SO<sub>4</sub>: 6.21 MICR EQUIV/L:  
 Ca: 98.9 Mg: 14.7 Na : 28.5 K : 9.97  
 DATE SAMPLED: 7-3-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.2456 GMS PER CC:  
 SED DRY WT: 0.0715 GMS PER CC: INGN LOSS: 23.1

TAXON NAME	Total	No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	522.00	2.00	0.383	
2006 <i>Achnanthes didyma</i> Hust. v. <i>didyma</i>	522.00	1.00	0.192	
2024 <i>Achnanthes linearis</i> (W. Sm.) Grun. v. <i>linearis</i>	522.00	5.00	0.958	
2026 <i>Achnanthes linearis</i> v. <i>pusilla</i> Grun.	522.00	5.00	0.958	
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	522.00	13.00	2.490	
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	522.00	25.00	4.789	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	522.00	1.00	0.192	
8001 <i>Anomoeoneis exilis</i> v. <i>lanceolata</i> A. Mayer	522.00	12.00	2.299	
8005 <i>Anomoeoneis serians</i> v. <i>brachysira</i> (Breb.) ex Kutz.	522.00	3.00	0.575	
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	522.00	1.00	0.192	
16001 <i>Coccconeis diminuta</i> Pant. v. <i>diminuta</i>	522.00	1.00	0.192	
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	522.00	34.00	6.513	
20012 <i>Cyclotella pseudostelligera</i> Hust. v. <i>pseudostelligera</i>	522.00	27.00	5.172	
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. <i>cesatii</i>	522.00	1.00	0.192	
23007 <i>Cymbella gaeumannii</i> Meist. v. <i>gaeumannii</i>	522.00	2.00	0.383	
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	522.00	14.00	2.682	
23010 <i>Cymbella microcephala</i> Grun. v. <i>microcephala</i>	522.00	14.00	2.682	
33026 <i>Eunotia incisa</i> W. Sm. ex Greg. v. <i>incisa</i>	522.00	1.00	0.192	
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	522.00	12.00	2.299	
34017 <i>Fragilaria crotonensis</i> Kitton v. <i>crotonensis</i>	522.00	1.00	0.192	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	522.00	3.00	0.575	
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	522.00	7.00	1.341	
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	522.00	57.00	10.920	
35001 <i>Frustulia rhomboides</i> (Ehr.) DeT. v. <i>rhomboides</i>	522.00	1.00	0.192	
35005 <i>Frustulia rhomboides</i> v. <i>saxonica</i> (Rabh.) DeT.	522.00	3.00	0.575	
44027 <i>Melosira</i> 1-PIRLA	522.00	46.00	8.812	
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	522.00	2.00	0.383	
46050 <i>Navicula pseudoscutiformis</i> Hust. v. <i>pseudoscutiformis</i>	522.00	2.00	0.383	
46051 <i>Navicula pupula</i> Kutz. v. <i>pupula</i>	522.00	4.00	0.766	
46056 <i>Navicula radios</i> Kutz. v. <i>radiosa</i>	522.00	2.00	0.383	
46095 <i>Navicula heimansii</i> van Dam & Kooyman v. <i>heimansii</i>	522.00	1.00	0.192	
46889 <i>Navicula</i> spp.	522.00	1.00	0.192	
48035 <i>Nitzschia</i> 1-PIRLA	522.00	20.00	3.831	
52889 <i>Pinnularia</i> spp.	522.00	2.00	0.383	
62002 <i>Stauroneis anceps</i> Ehr. v. <i>anceps</i>	522.00	1.00	0.192	
63002 <i>Stenopterobia intermedia</i> (Lewis) V. H. v. <i>intermedia</i>	522.00	4.00	0.766	
66016 <i>Synedra rumpens</i> Kutz. v. <i>rumpens</i>	522.00	1.00	0.192	
90990 <i>Nitzschia</i> 42-SN	522.00	9.00	1.724	
99088 <i>Fragilaria</i> 2-SN	522.00	63.00	12.069	
99096 <i>Fragilaria</i> 14-SN	522.00	12.00	2.299	
99124 <i>Krasskella kriegeriana</i> (Krasske) Ross & Sims	522.00	3.00	0.575	
99126 <i>Melosira</i> 1-SN	522.00	17.00	3.257	

TAXON NAME	Total	No	This Tax	%REL ABUND
99133 <i>Navicula cari</i> Ehr.	522.00	3.00	0.575	
99231 <i>Nitzschia frustulum</i> 3-SN	522.00	3.00	0.575	
99241 <i>Nitzschia</i> 6-SN	522.00	2.00	0.383	
99250 <i>Nitzschia</i> 15-SN	522.00	13.00	2.490	
99260 <i>Nitzschia</i> 28-SN	522.00	1.00	0.192	
99267 <i>Nitzschia</i> 36-SN	522.00	3.00	0.575	
99270 <i>Nitzschia</i> 39-SN	522.00	1.00	0.192	
99281 <i>Pinnularia</i> 9-SN	522.00	5.00	0.958	
99283 <i>Pinnularia</i> 11-SN	522.00	3.00	0.575	
99294 <i>Pinnularia</i> 23-SN	522.00	1.00	0.192	
99330 <i>Synedra</i> 2-SN	522.00	2.00	0.383	
99347 <i>Synedra pulchella</i> Kutz.	522.00	1.00	0.192	
99350 <i>Cymbella</i> 13-SN	522.00	12.00	2.299	
99351 <i>Nitzschia</i> 41-SN	522.00	32.00	6.130	
99353 <i>Pinnularia</i> 43-SN	522.00	1.00	0.192	
99355 <i>Eunotia</i> sp. (girdle view)	522.00	1.00	0.192	
99356 <i>Nitzschia</i> 42-SN	522.00	2.00	0.383	

LAKE: UPPER GAYLOR LAT DEG: 37 MIN: 55 SEC: 20 LONG DEG: 119 MIN: 16 SEC: 1  
 QUADRANGLE: TUOL. MEADOW. AREA: 4.66 SHED AREA: 41.2 ELEV: 3121  
 VEG CLASS: 2 ROCK TYPE: % COVER: A: 1 B: 4 C: 1 D: 1  
 pH FIELD: 7.37 pH LAB: 7.27 CONDUCTIVITY: 11.9 usa25:  
 HCO<sub>3</sub>: 102.4 Cl: 1.17 NO<sub>3</sub>: 0 SO<sub>4</sub>: 20.2 MICR EQUIV/L:  
 Ca: 111 Mg: 5.59 Na : 8.77 K : 1.59  
 DATE SAMPLED: 8-16-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.14 GMS PER CC:  
 SED DRY WT: 0.0886 GMS PER CC: INGN LOSS: 19.4

TAXON NAME	Total No	This Tax	%REL ABUND
2006 <i>Achnanthes didyma</i> Hust. v. <i>didyma</i>	511.00	1.00	0.196
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	511.00	3.00	0.587
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	511.00	3.00	0.587
2010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	511.00	38.00	7.436
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	511.00	3.00	0.587
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	511.00	6.00	1.174
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	511.00	3.00	0.587
23016 <i>Cymbella naviculiformis</i> Auersw. ex Heib. v.	511.00	2.00	0.391
33066 <i>Eunotia vanheurckii</i> v. <i>intermedia</i> (Krasske) ex Hust.	511.00	1.00	0.196
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	511.00	57.00	11.155
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	511.00	23.00	4.501
34027 <i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.	511.00	1.00	0.196
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	511.00	1.00	0.196
44010 <i>Melosira italica</i> (Ehr.) Kutz. v. <i>italica</i>	511.00	4.00	0.783
44027 <i>Melosira</i> 1-PIRLA	511.00	1.00	0.196
46014 <i>Navicula cryptocephala</i> Kutz. v. <i>cryptocephala</i>	511.00	5.00	0.978
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	511.00	3.00	0.587
46050 <i>Navicula pseudoscutiformis</i> Hust. v. <i>pseudoscutiformis</i>	511.00	2.00	0.391
46057 <i>Navicula radiosha</i> v. <i>parva</i> Wallace	511.00	1.00	0.196
46133 <i>Navicula</i> 14-PIRLA	511.00	1.00	0.196
46889 <i>Navicula</i> spp.	511.00	1.00	0.196
48008 <i>Nitzschia dissipata</i> (Kutz.) Grun. v. <i>dissipata</i>	511.00	1.00	0.196
48889 <i>Nitzschia</i> spp.	511.00	1.00	0.196
62003 <i>Stauroneis anceps</i> f. <i>gracilis</i> Rabh.	511.00	2.00	0.391
62024 <i>Stauroneis anceps</i> v. 1-PIRLA	511.00	1.00	0.196
90990 <i>Nitzschia</i> 42-SN	511.00	1.00	0.196
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	511.00	1.00	0.196
99006 <i>Achnanthes suchlandi</i> Hust.	511.00	1.00	0.196
99009 <i>Achnanthes</i> 3-SN	511.00	18.00	3.523
99015 <i>Achnanthes</i> 10-SN	511.00	1.00	0.196
99088 <i>Fragilaria</i> 2-SN	511.00	5.00	0.978
99090 <i>Fragilaria</i> 7-SN	511.00	76.00	14.873
99094 <i>Fragilaria</i> 11-SN	511.00	31.00	6.067
99105 <i>Gomphonema puiggarianum</i> v. <i>aequatorialis</i> (Cl.) Camburn	511.00	2.00	0.391
99109 <i>Gomphonema</i> 3-SN	511.00	1.00	0.196
99137 <i>Navicula mediopunctata</i> Hust.	511.00	2.00	0.391
99142 <i>Navicula</i> 1-SN	511.00	117.00	22.896
99146 <i>Navicula</i> 10-SN	511.00	1.00	0.196
99187 <i>Navicula</i> 58-SN	511.00	73.00	14.286
99189 <i>Navicula</i> 60-SN	511.00	1.00	0.196
99227 <i>Nitzschia admissoides</i> Hust.	511.00	1.00	0.196
99229 <i>Nitzschia frustulum</i> 1-SN	511.00	1.00	0.196

TAXON NAME	Total	No	This Tax	%REL ABUND
99230 <i>Nitzschia frustulum</i> 2-SN	511.00		1.00	0.196
99241 <i>Nitzschia</i> 6-SN	511.00		1.00	0.196
99281 <i>Pinnularia</i> 9-SN	511.00		6.00	1.174
99330 <i>Synedra</i> 2-SN	511.00		5.00	0.978

LAKE: LUNDY LAT DEG: 38 MIN: 1 SEC: 44 LONG DEG: 119 MIN: 16 SEC: 13  
 QUADRANGLE: BODIE AREA: 42.7 SHED AREA: 4507 ELEV: 2390  
 VEG CLASS: 5 ROCK TYPE: % COVER: A: 2 B: 3 C: 3 D: 1  
 PH FIELD: 7.6 pH LAB: 7.41 CONDUCTIVITY: 49.9 us@25:  
 HCO<sub>3</sub>: 260.5 Cl: 3.67 NO<sub>3</sub>: 0 SO<sub>4</sub>: 175 MICR EQUIV/L:  
 Ca: 381.2 Mg: 27.5 Na: 44.2 K: 14.6  
 DATE SAMPLED: 8-5-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.2958 GMS PER CC:  
 SED DRY WT: 0.0688 GMS PER CC: INGN LOSS: 17.3

TAXON NAME	Total No	This Tax	%REL ABUND
2015 <i>Achnanthes lanceolata</i> (Breb.) Grun. v. <i>lanceolata</i>	506.00	2.00	0.395
2024 <i>Achnanthes linearis</i> (W. Sm.) Grun. v. <i>linearis</i>	506.00	2.00	0.395
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	506.00	4.00	0.791
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	506.00	49.00	9.684
2048 <i>Achnanthes laterostrata</i> Hust. v. <i>laterostrata</i>	506.00	2.00	0.395
2889 <i>Achnanthes</i> spp.	506.00	1.00	0.198
7001 <i>Amphora ovalis</i> (Kutz.) Kutz. v. <i>ovalis</i>	506.00	1.00	0.198
8001 <i>Anomoeoneis exilis</i> v. <i>lanceolata</i> A. Mayer	506.00	1.00	0.198
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	506.00	87.00	17.194
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	506.00	2.00	0.395
16001 <i>Cocconeis diminuta</i> Pant. v. <i>diminuta</i>	506.00	1.00	0.198
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	506.00	4.00	0.791
23005 <i>Cymbella cistula</i> (Ehr.) Kirchner v. <i>cistula</i>	506.00	5.00	0.988
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	506.00	4.00	0.791
23010 <i>Cymbella microcephala</i> Grun. v. <i>microcephala</i>	506.00	7.00	1.383
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	506.00	3.00	0.593
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	506.00	3.00	0.593
23031 <i>Cymbella muelleri</i> Hust. v. <i>muelleri</i>	506.00	1.00	0.198
30001 <i>Diploneis elliptica</i> (Kutz.) Cl. v. <i>elliptica</i>	506.00	3.00	0.593
34006 <i>Fragilaria capucina</i> Desm. v. <i>capucina</i>	506.00	2.00	0.395
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	506.00	18.00	3.557
34017 <i>Fragilaria crotonensis</i> Kitton v. <i>crotonensis</i>	506.00	33.00	6.522
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	506.00	2.00	0.395
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	506.00	87.00	17.194
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	506.00	6.00	1.186
34032 <i>Fragilaria virescens</i> Rolfs v. <i>virescens</i>	506.00	1.00	0.198
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	506.00	1.00	0.198
44010 <i>Melosira italica</i> (Ehr.) Kutz. v. <i>italica</i>	506.00	1.00	0.198
44027 <i>Melosira</i> 1-PIRLA	506.00	3.00	0.593
46014 <i>Navicula cryptocephala</i> Kutz. v. <i>cryptocephala</i>	506.00	1.00	0.198
46021 <i>Navicula globulifera</i> Hust. v. <i>globulifera</i>	506.00	1.00	0.198
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	506.00	1.00	0.198
46057 <i>Navicula radiosha</i> v. <i>parva</i> Wallace	506.00	2.00	0.395
46889 <i>Navicula</i> spp.	506.00	1.00	0.198
48008 <i>Nitzschia dissipata</i> (Kutz.) Grun. v. <i>dissipata</i>	506.00	2.00	0.395
52889 <i>Pinnularia</i> spp.	506.00	2.00	0.395
66016 <i>Synedra rumpens</i> Kutz. v. <i>rumpens</i>	506.00	1.00	0.198
66029 <i>Synedra</i> 1-PIRLA	506.00	12.00	2.372
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	506.00	1.00	0.198
99006 <i>Achnanthes suchlandi</i> Hust.	506.00	1.00	0.198
99009 <i>Achnanthes</i> 3-SN	506.00	1.00	0.198
99029 <i>Achnanthes</i> 37-SN	506.00	1.00	0.198

TAXON NAME	Total	No	This Tax	%REL ABUND
99031 Achnanthes 39-SN	506.00	1.00	0.198	
99038 Cocconeis placentula Ehr. v. placentula	506.00	1.00	0.198	
99067 Epithemia adnata (Kutz.) Breb.	506.00	1.00	0.198	
99088 Fragilaria 2-SN	506.00	12.00	2.372	
99090 Fragilaria 7-SN	506.00	35.00	6.917	
99094 Fragilaria 11-SN	506.00	20.00	3.953	
99095 Fragilaria 13-SN	506.00	1.00	0.198	
99096 Fragilaria 14-SN	506.00	4.00	0.791	
99115 Gomphonema 9-SN	506.00	1.00	0.198	
99121 Gyrosigma spencerii (Quekett) Griffith & Henfries	506.00	1.00	0.198	
99128 Melosira 3-SN	506.00	1.00	0.198	
99133 Navicula cari Ehr.	506.00	5.00	0.988	
99142 Navicula 1-SN	506.00	3.00	0.593	
99147 Navicula 11-SN	506.00	1.00	0.198	
99188 Navicula 59-SN	506.00	2.00	0.395	
99190 Navicula 61-SN	506.00	9.00	1.779	
99214 Navicula 85-SN	506.00	5.00	0.988	
99229 Nitzschia frustulum 1-SN	506.00	3.00	0.593	
99231 Nitzschia frustulum 3-SN	506.00	8.00	1.581	
99235 Nitzschia frustulum 7-SN	506.00	6.00	1.186	
99239 Nitzschia 3-SN	506.00	3.00	0.593	
99241 Nitzschia 6-SN	506.00	5.00	0.988	
99327 Synedra capitata Ehr.	506.00	1.00	0.198	
99330 Synedra 2-SN	506.00	14.00	2.767	

LAKE: CONSTANCE LAT DEG: 37 MIN: 30 SEC: 55 LONG DEG: 118 MIN: 52 SEC: 2  
 QUADRANGLE: MT. MORRISON AREA: 14 SHED AREA: 207 ELEV: 3292  
 VEG CLASS: 1 ROCK TYPE: % COVER: A: 1 B: 4 C: 1 D: 1  
 pH FIELD: 7.95 pH LAB: 7.42 CONDUCTIVITY: us@25:  
 HCO<sub>3</sub>: 271.5 Cl: 1.94 NO<sub>3</sub>: 2.1 SO<sub>4</sub>: 97.8 MICR EQUIV/L:  
 Ca: 345 Mg: 12.4 Na: 8.27 K: 7.58  
 DATE SAMPLED: 8-20-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.1284 GMS PER CC:  
 SED DRY WT: 0.1789 GMS PER CC: INGN LOSS: 8.24

TAXON NAME	Total	No	This Tax	%REL ABUND
2002 <i>Achnanthes austriaca</i> Hust. v. austriaca	521.00	2.00	0.384	
2006 <i>Achnanthes didyma</i> Hust. v. didyma	521.00	1.00	0.192	
2026 <i>Achnanthes linearis</i> v. <i>pusilla</i> Grun.	521.00	5.00	0.960	
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	521.00	5.00	0.960	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	521.00	2.00	0.384	
2048 <i>Achnanthes laterostrata</i> Hust. v. <i>laterostrata</i>	521.00	2.00	0.384	
2889 <i>Achnanthes</i> spp.	521.00	5.00	0.960	
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. <i>perpusilla</i>	521.00	4.00	0.768	
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	521.00	45.00	8.637	
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	521.00	1.00	0.192	
20009 <i>Cyclotella ocellata</i> Pant. v. <i>ocellata</i>	521.00	4.00	0.768	
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	521.00	51.00	9.789	
20012 <i>Cyclotella pseudostelligera</i> Hust. v. <i>pseudostelligera</i>	521.00	16.00	3.071	
23009 <i>Cymbella lunata</i> W. Sm. v. <i>lunata</i>	521.00	1.00	0.192	
23010 <i>Cymbella microcephala</i> Grun. v. <i>microcephala</i>	521.00	3.00	0.576	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	521.00	4.00	0.768	
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	521.00	3.00	0.576	
30003 <i>Diploneis marginestriata</i> Hust. v. <i>marginestriata</i>	521.00	1.00	0.192	
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	521.00	9.00	1.727	
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	521.00	15.00	2.879	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	521.00	24.00	4.607	
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	521.00	6.00	1.152	
46889 <i>Navicula</i> spp.	521.00	1.00	0.192	
48008 <i>Nitzschia dissipata</i> (Kutz.) Grun. v. <i>dissipata</i>	521.00	1.00	0.192	
66014 <i>Synedra parasitica</i> (W. Sm.) Hust. v. <i>parasitica</i>	521.00	1.00	0.192	
66015 <i>Synedra radians</i> (Kutz.) v <i>radians</i>	521.00	22.00	4.223	
66016 <i>Synedra rumpens</i> Kutz. v. <i>rumpens</i>	521.00	1.00	0.192	
90990 <i>Nitzschia</i> 42-SN	521.00	4.00	0.768	
99001 <i>Achnanthes bicapitata</i> Hust.	521.00	1.00	0.192	
99002 <i>Achnanthes calcar</i> Cl.	521.00	1.00	0.192	
99006 <i>Achnanthes suchlandi</i> Hust.	521.00	13.00	2.495	
99009 <i>Achnanthes</i> 3-SN	521.00	7.00	1.344	
99013 <i>Achnanthes</i> 8-SN	521.00	1.00	0.192	
99039 <i>Cocconeis placentula</i> v. <i>euglypta</i> (Ehr.) Cl.	521.00	2.00	0.384	
99040 <i>Cocconeis</i> 2-SN	521.00	1.00	0.192	
99088 <i>Fragilaria</i> 2-SN	521.00	51.00	9.789	
99090 <i>Fragilaria</i> 7-SN	521.00	13.00	2.495	
99094 <i>Fragilaria</i> 11-SN	521.00	46.00	8.829	
99096 <i>Fragilaria</i> 14-SN	521.00	1.00	0.192	
99119 <i>Gomphonema</i> 13-SN	521.00	1.00	0.192	
99132 <i>Navicula capitata</i> v. <i>hungarica</i> (Grun.) Ross	521.00	4.00	0.768	
99137 <i>Navicula mediopunctata</i> Hust.	521.00	3.00	0.576	

TAXON NAME	Total	No	This Tax	%REL ABUND
99138 <i>Navicula pseudolanceolata</i> Lange-Bertalot	521.00	1.00	0.192	
99146 <i>Navicula</i> 10-SN	521.00	1.00	0.192	
99157 <i>Navicula</i> 27-SN	521.00	2.00	0.384	
99166 <i>Navicula</i> 37-SN	521.00	3.00	0.576	
99168 <i>Navicula</i> 39-SN	521.00	4.00	0.768	
99175 <i>Navicula</i> 46-SN	521.00	5.00	0.960	
99183 <i>Navicula</i> 54-SN	521.00	2.00	0.384	
99194 <i>Navicula</i> 65-SN	521.00	5.00	0.960	
99198 <i>Navicula</i> 69-SN	521.00	1.00	0.192	
99201 <i>Navicula</i> 72-SN	521.00	2.00	0.384	
99209 <i>Navicula</i> 80-SN	521.00	1.00	0.192	
99210 <i>Navicula</i> 81-SN	521.00	2.00	0.384	
99231 <i>Nitzschia frustulum</i> 3-SN	521.00	3.00	0.576	
99233 <i>Nitzschia frustulum</i> 5-SN	521.00	1.00	0.192	
99240 <i>Nitzschia</i> 5-SN	521.00	1.00	0.192	
99271 <i>Opephora martyi</i> Heribaud	521.00	2.00	0.384	
99330 <i>Synedra</i> 2-SN	521.00	94.00	18.042	
99331 <i>Synedra</i> 4-SN	521.00	5.00	0.960	
99343 <i>Nitzschia</i> 42-SN	521.00	1.00	0.192	
99350 <i>Cymbella</i> 13-SN	521.00	1.00	0.192	

LAKE: UPPER FROG LAT DEG: 38 MIN: 2 SEC: 38 LONG DEG: 119 MIN: 17 SEC: 20  
 QUADRANGLE: MATTERHRN. PK. AREA: 0.78 SHED AREA: 57.2 ELEV: 3231  
 VEG CLASS: 2 ROCK TYPE: % COVER: A: 1 B: 4 C: 1 D: 1  
 pH FIELD: 7.85 pH LAB: 7.43 CONDUCTIVITY: 55.5 uS@25:  
 HCO<sub>3</sub>: 257.6 Cl: 2.86 NO<sub>3</sub>: 3.29 SO<sub>4</sub>: 247.4 MICR EQUIV/L:  
 Ca: 432 Mg: 21.2 Na: 35.8 K : 7.18  
 DATE SAMPLED: 3-8-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.4103 GMS PER CC:  
 SED DRY WT: 0.1238 GMS PER CC: INGN LOSS: 13.6

TAXON NAME	Total No	This Tax	%REL ABUND
2015 <i>Achnanthes lanceolata</i> (Breb.) Grun. v. <i>lanceolata</i>	532.00	2.00	0.376
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. <i>perpusilla</i>	532.00	3.00	0.564
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	532.00	5.00	0.940
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	532.00	7.00	1.316
30003 <i>Diploneis marginestriata</i> Hust. v. <i>marginestriata</i>	532.00	1.00	0.188
34013 <i>Fragilaria construens</i> v. <i>binodis</i> (Ehr.) Grun.	532.00	3.00	0.564
34014 <i>Fragilaria construens</i> v. <i>pumila</i> Grun.	532.00	3.00	0.564
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	532.00	406.00	76.316
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	532.00	31.00	5.827
34027 <i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.	532.00	1.00	0.188
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	532.00	1.00	0.188
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	532.00	1.00	0.188
34038 <i>Fragilaria pinnata</i> v. <i>acuminata</i> A. Mayer	532.00	17.00	3.195
99012 <i>Achnanthes</i> 7-SN	532.00	1.00	0.188
99087 <i>Fragilaria</i> 6-PIRLA	532.00	30.00	5.639
99088 <i>Fragilaria</i> 2-SN	532.00	2.00	0.376
99089 <i>Fragilaria</i> 4-SN	532.00	4.00	0.752
99092 <i>Fragilaria</i> 9-SN	532.00	3.00	0.564
99094 <i>Fragilaria</i> 11-SN	532.00	3.00	0.564
99233 <i>Nitzschia frustulum</i> 5-SN	532.00	7.00	1.316

LAKE: WIT-SO-NAH-PAH LAT DEG: MIN: SEC: LONG DEG: MIN: SEC:  
 QUADRANGLE: MT.MORRISON AREA: SHED AREA: ELEV:  
 VEG CLASS: ROCK TYPE: % COVER: A: B: C: D:  
 pH FIELD: 8.15 pH LAB: 7.57 CONDUCTIVITY: 37.4 usA25:  
 HCO<sub>3</sub>: 228.7 Cl: 2.15 NO<sub>3</sub>: 3.11 SO<sub>4</sub>: 132.9 MICR EQUIV/L:  
 Ca: 385 Mg: 11.4 Na : 10.1 K : 6.94  
 DATE SAMPLED: 8-20-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.5037 GMS PER CC:  
 SED DRY WT: 0.7457 GMS PER CC: INGN LOSS: 5.11

TAXON NAME	Total	No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	516.00	2.00	0.388	
2016 <i>Achnanthes lanceolata</i> v. <i>dubia</i> Grun.	516.00	5.00	0.969	
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	516.00	3.00	0.581	
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	516.00	54.00	10.465	
2042 <i>Achnanthes detha</i> Hohn & Hellerm. v. <i>detha</i>	516.00	1.00	0.194	
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. <i>perpusilla</i>	516.00	1.00	0.194	
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	516.00	1.00	0.194	
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	516.00	2.00	0.388	
20005 <i>Cyclotella kuetzingiana</i> Thwaites v. <i>kuetzingiana</i>	516.00	8.00	1.550	
20009 <i>Cyclotella ocellata</i> Pant. v. <i>ocellata</i>	516.00	1.00	0.194	
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	516.00	67.00	12.984	
20012 <i>Cyclotella pseudostelligera</i> Hust. v. <i>pseudostelligera</i>	516.00	2.00	0.388	
23004 <i>Cymbella cesatii</i> (Rabh.) Grun. ex A. S. v. <i>cesatii</i>	516.00	1.00	0.194	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	516.00	53.00	10.271	
23013 <i>Cymbella minuta</i> f. <i>latens</i> (Krasske) Reim.	516.00	2.00	0.388	
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	516.00	97.00	18.798	
27002 <i>Diatoma hiemale</i> v. <i>mesodon</i> (Ehr.) Grun.	516.00	1.00	0.194	
33040 <i>Eunotia pectinalis</i> v. <i>minor</i> (Kutz.) Rabh.	516.00	1.00	0.194	
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	516.00	1.00	0.194	
34022 <i>Fragilaria leptostauron</i> (Ehr.) Hust.	516.00	20.00	3.876	
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	516.00	54.00	10.465	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	516.00	11.00	2.132	
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	516.00	4.00	0.775	
37004 <i>Gomphonema angustatum</i> v. <i>citera</i> (Hohn & Hum) Kutz.	516.00	6.00	1.163	
46014 <i>Navicula cryptocephala</i> Kutz. v. <i>cryptocephala</i>	516.00	2.00	0.580	
46050 <i>Navicula pseudoscutiformis</i> Hust. v. <i>pseudoscutiformis</i>	516.00	1.00	0.194	
62002 <i>Stauroneis anceps</i> Ehr. v. <i>anceps</i>	516.00	1.00	0.194	
66016 <i>Synedra rumpens</i> Kutz. v. <i>rumpens</i>	516.00	2.00	0.388	
66024 <i>Synedra ulna</i> (Nitz.) Ehr. v. <i>ulna</i>	516.00	1.00	0.194	
90990 <i>Nitzschia</i> 42-SN	516.00	2.00	0.388	
99025 <i>Achnanthes</i> 31-SN	516.00	1.00	0.194	
99094 <i>Fragilaria</i> 11-SN	516.00	3.00	0.581	
99096 <i>Fragilaria</i> 14-SN	516.00	1.00	0.194	
99117 <i>Gomphonema</i> 11-SN	516.00	1.00	0.194	
99123 <i>Hannea arcus</i> (Ehr.) Pat.	516.00	5.00	0.969	
99151 <i>Navicula</i> 18-SN	516.00	1.00	0.194	
99153 <i>Navicula</i> 22-SN	516.00	3.00	0.581	
99184 <i>Navicula</i> 55-SN	516.00	2.00	0.388	
99202 <i>Navicula</i> 73-SN	516.00	2.00	0.388	
99231 <i>Nitzschia frustulum</i> 3-SN	516.00	6.00	1.163	
99236 <i>Nitzschia frustulum</i> 8-SN	516.00	1.00	0.194	
99241 <i>Nitzschia</i> 6-SN	516.00	2.00	0.388	

TAXON NAME	Total	No	This Tax	%REL ABUND
99266 <i>Nitzschia</i> 35-SN	516.00	40.00	7.752	
99267 <i>Nitzschia</i> 36-SN	516.00	33.00	6.395	
99276 <i>Pinnularia</i> 1-SN	516.00	1.00	0.194	
99330 <i>Synedra</i> 2-SN	516.00	3.00	0.581	
99342 <i>Cymbella cymbiformis</i> Agardh	516.00	1.00	0.194	
99347 <i>Synedra pulchella</i> Kutz.	516.00	2.00	0.388	

LAKE: EASTERN TWIN LAT DEG: 38 MIN: 9 SEC: 40 LONG DEG: 119 MIN: 20 SEC: 0  
 QUADRANGLE: MATTERHORN PK AREA: 161 SHED AREA: 6786 ELEV: 2158  
 VEG CLASS: 5 ROCK TYPE: % COVER: A: 2 B: 2 C: 1 D: 4  
 pH FIELD: 8.14 pH LAB: 7.66 CONDUCTIVITY: 49.5 USA25:  
 HCO<sub>3</sub>: 383.6 Cl: 5.72 NO<sub>3</sub>: 0 SO<sub>4</sub>: 78.1 MICR EQUIV/L:  
 Ca: 356 Mg: 60 Na: 71.3 K: 13.9  
 DATE SAMPLED: 8-3-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 0.96 GMS PER CC:  
 SED DRY WT: 0.1192 GMS PER CC: INGN LOSS: 12.4

TAXON NAME	Total No	This Tax	%REL ABUND
2003 <i>Achnanthes austriaca</i> Hust. v. <i>helvetica</i>	501.00	2.00	0.399
2015 <i>Achnanthes lanceolata</i> (Breb.) Grun. v. <i>lanceolata</i>	501.00	6.00	1.198
2026 <i>Achnanthes linearis</i> v. <i>pusilla</i> Grun.	501.00	1.00	0.200
2028 <i>Achnanthes marginulata</i> Grun. v. <i>marginulata</i>	501.00	1.00	0.200
2047 <i>Achnanthes peragalli</i> v. <i>fossilis</i> Temp. & Perag.	501.00	1.00	0.200
2048 <i>Achnanthes laterostrata</i> Hust. v. <i>laterostrata</i>	501.00	1.00	0.200
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	501.00	23.00	4.591
23005 <i>Cymbella cistula</i> (Ehr.) Kirchner v. <i>cistula</i>	501.00	1.00	0.200
23010 <i>Cymbella microcephala</i> Grun. v. <i>microcephala</i>	501.00	4.00	0.798
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	501.00	2.00	0.399
34003 <i>Fragilaria brevistriata</i> Grun. v. <i>brevistriata</i>	501.00	32.00	6.387
34012 <i>Fragilaria construens</i> (Ehr. Grun. v. <i>construens</i>	501.00	1.00	0.200
34013 <i>Fragilaria construens</i> v. <i>binodis</i> (Ehr.) Grun.	501.00	32.00	6.387
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	501.00	7.00	1.397
34017 <i>Fragilaria crotensis</i> Kitton v. <i>crotensis</i>	501.00	168.00	33.533
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	501.00	8.00	1.597
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	501.00	16.00	3.194
37012 <i>Gomphonema subtile</i> Ehr. v. <i>subtile</i>	501.00	2.00	0.399
44001 <i>Melosira ambigua</i> (Grun.) O. Mull. v. <i>ambigua</i>	501.00	9.00	1.796
44010 <i>Melosira italica</i> (Ehr.) Kutz. v. <i>italica</i>	501.00	26.00	5.190
44011 <i>Melosira italica</i> ssp. <i>subarctica</i> O. Mull.	501.00	80.00	15.968
44040 <i>Melosira italica</i> ssp. <i>subarctica</i> f. <i>tenuissima</i> (Grun.)	501.00	19.00	3.792
46133 <i>Navicula</i> 14-PIRLA	501.00	1.00	0.200
67005 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain III sensu	501.00	15.00	2.994
99019 <i>Achnanthes</i> 18-SN	501.00	1.00	0.200
99086 <i>Fragilaria capucina</i> v. <i>mesolepta</i> Rabh.	501.00	2.00	0.399
99087 <i>Fragilaria</i> 6-PIRLA	501.00	5.00	0.998
99105 <i>Gomphonema puiggarianum</i> v. <i>aequatorialis</i> (Cl.) Camburn	501.00	2.00	0.399
99157 <i>Navicula</i> 27-SN	501.00	1.00	0.200
99158 <i>Navicula</i> 28-SN	501.00	1.00	0.200
99229 <i>Nitzschia frustulum</i> 1-SN	501.00	1.00	0.200
99230 <i>Nitzschia frustulum</i> 2-SN	501.00	1.00	0.200
99246 <i>Nitzschia</i> 11-SN	501.00	2.00	0.399
99271 <i>Opephora martyi</i> Heribaud	501.00	6.00	1.198
99281 <i>Pinnularia</i> 9-SN	501.00	1.00	0.200
99316 <i>Stephanodiscus parvus</i> Stoermer & Hakansson	501.00	17.00	3.393
99317 <i>Stephanodiscus</i> 1-SN	501.00	21.00	4.192
99318 <i>Stephanodiscus</i> 2-SN	501.00	1.00	0.200

LAKE: WESTERN TWIN LAT DEG: 38 MIN: 8 SEC: 51 LONG DEG: 119 MIN: 21 SEC: 40  
 QUADRANGLE: MATTERHORN PK AREA: 120 SHED AREA: 5284 ELEV: 2162  
 VEG CLASS: 5 ROCK TYPE: % COVER: A: 2 B: 2 C: 1 D: 4  
 pH FIELD: 7.96 pH LAB: 7.8 CONDUCTIVITY: 40.2 uS/cm:  
 HCO<sub>3</sub>: 324.4 Cl: 3.73 NO<sub>3</sub>: 0 SO<sub>4</sub>: 52.1 MICR EQUIV/L:  
 Ca: 322 Mg: 42.1 Na: 52.1 K: 9.45  
 DATE SAMPLED: 38-3-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.0922 GMS PER CC:  
 SED DRY WT: 0.1195 GMS PER CC: INGN LOSS: 13.5

TAXON NAME	Total	No	This Tax	%REL ABUND
2012 <i>Achnanthes haukiana</i> Grun. v. <i>haukiana</i>	513.00	1.00	0.195	
2015 <i>Achnanthes lanceolata</i> (Breb.) Grun. v. <i>lanceolata</i>	513.00	5.00	0.975	
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	513.00	7.00	1.365	
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	513.00	28.00	5.458	
20012 <i>Cyclotella pseudostelligera</i> Hust. v. <i>pseudostelligera</i>	513.00	1.00	0.195	
23005 <i>Cymbella cistula</i> (Ehr.) Kirchner v. <i>cistula</i>	513.00	1.00	0.195	
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	513.00	6.00	1.170	
23013 <i>Cymbella minuta</i> f. <i>latens</i> (Krasske) Reim.	513.00	3.00	0.585	
27002 <i>Diatoma hiemale</i> v. <i>mesodon</i> (Ehr.) Grun.	513.00	2.00	0.390	
30003 <i>Diploneis marginestriata</i> Hust. v. <i>marginestriata</i>	513.00	1.00	0.195	
34003 <i>Fragilaria brevistriata</i> Grun. v. <i>brevistriata</i>	513.00	1.00	0.195	
34006 <i>Fragilaria capucina</i> Desm. v. <i>capucina</i>	513.00	1.00	0.195	
34012 <i>Fragilaria construens</i> (Ehr.) Grun. v. <i>construens</i>	513.00	1.00	0.195	
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	513.00	1.00	0.195	
34017 <i>Fragilaria crotonensis</i> Kitton v. <i>crotonensis</i>	513.00	50.00	9.747	
34022 <i>Fragilaria leptostauron</i> (Ehr.) Hust.	513.00	1.00	0.195	
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	513.00	10.00	1.949	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	513.00	21.00	4.094	
34027 <i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.	513.00	1.00	0.195	
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	513.00	5.00	0.975	
34032 <i>Fragilaria virescens</i> Rolfs v. <i>virescens</i>	513.00	10.00	1.949	
34038 <i>Fragilaria pinnata</i> v. <i>acuminata</i> A. Mayer	513.00	24.00	4.678	
37889 <i>Gomphonema</i> spp.	513.00	3.00	0.585	
44001 <i>Melosira ambigua</i> (Grun.) O. Mull. v. <i>ambigua</i>	513.00	39.00	7.602	
44010 <i>Melosira italicica</i> (Ehr.) Kutz. v. <i>italicica</i>	513.00	46.00	8.967	
44011 <i>Melosira italicica</i> ssp. <i>subarctica</i> O. Mull.	513.00	91.00	17.739	
44040 <i>Melosira italicica</i> ssp. <i>subarctica</i> f. <i>tenuissima</i> (Grun.)	513.00	9.00	1.754	
45001 <i>Meridion circulare</i> (Grev.) Agardh v. <i>circulare</i>	513.00	1.00	0.195	
46056 <i>Navicula radios</i> Kutz. v. <i>radiosa</i>	513.00	2.00	0.390	
52013 <i>Pinnularia borealis</i> Ehr. v. <i>borealis</i>	513.00	1.00	0.195	
58001 <i>Rhopalodia gibba</i> (Ehr.) O. Mull. v. <i>gibba</i>	513.00	1.00	0.195	
66014 <i>Synedra parasitica</i> (W. Sm.) Hust. v. <i>parasitica</i>	513.00	2.00	0.390	
66023 <i>Synedra tenera</i> W. Sm. v. <i>tenera</i>	513.00	2.00	0.390	
67005 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain III sensu	513.00	22.00	4.288	
99003 <i>Achnanthes daui</i> v. <i>alaskaensis</i> Foged	513.00	1.00	0.195	
99006 <i>Achnanthes suchlandi</i> Hust.	513.00	3.00	0.585	
99007 <i>Achnanthes</i> 1-SN	513.00	2.00	0.390	
99010 <i>Achnanthes</i> 4-SN	513.00	1.00	0.195	
99013 <i>Achnanthes</i> 8-SN	513.00	4.00	0.780	
99051 <i>Cymbella sinuata</i> Greg.	513.00	2.00	0.390	
99067 <i>Epithemia adnata</i> (Kutz.) Breb.	513.00	3.00	0.585	
99069 <i>Epithemia turgida</i> (Ehr.) Kutz.	513.00	5.00	0.975	

TAXON NAME	Total	No	This Tax	%REL ABUND
99087 <i>Fragilaria</i> 6-PIRLA	513.00	4.00	0.780	
99090 <i>Fragilaria</i> 7-SN	513.00	9.00	1.754	
99091 <i>Fragilaria</i> 8-SN	513.00	2.00	0.390	
99092 <i>Fragilaria</i> 9-SN	513.00	1.00	0.195	
99094 <i>Fragilaria</i> 11-SN	513.00	1.00	0.195	
99115 <i>Gomphonema</i> 9-SN	513.00	1.00	0.195	
99123 <i>Hannea arcus</i> (Ehr.) Pat.	513.00	1.00	0.195	
99126 <i>Melosira</i> 1-SN	513.00	5.00	0.975	
99157 <i>Navicula</i> 27-SN	513.00	6.00	1.170	
99162 <i>Navicula</i> 31-SN	513.00	2.00	0.390	
99175 <i>Navicula</i> 46-SN	513.00	1.00	0.195	
99198 <i>Navicula</i> 69-SN	513.00	1.00	0.195	
99229 <i>Nitzschia frustulum</i> 1-SN	513.00	1.00	0.195	
99234 <i>Nitzschia frustulum</i> 6-SN	513.00	2.00	0.390	
99316 <i>Stephanodiscus parvus</i> Stoermer & Hakansson	513.00	8.00	1.559	
99317 <i>Stephanodiscus</i> 1-SN	513.00	44.00	8.577	
99321 <i>Stephanodiscus</i> 5-SN	513.00	1.00	0.195	
99341 <i>Nitzschia amphibiodes</i> Hust.	513.00	2.00	0.390	

LAKE: CONVICT LAT DEG: 37 MIN: 35 SEC: 30 LONG DEG: 118 MIN: 51 SEC: 25  
 QUADRANGLE: MT.MORRISON AREA: 74.6 SHED AREA: 4999 ELEV: 2310  
 VEG CLASS: 5 ROCK TYPE: % COVER: A: 3 B: 3 C: 3 D: 1  
 pH FIELD: 8.4 pH LAB: 8.35 CONDUCTIVITY: 119 uS@25:  
 HCO<sub>3</sub>: 1051 Cl: 1.86 NO<sub>3</sub>: 0 SO<sub>4</sub>: 231.8 MICR EQUIV/L:  
 Ca: 1234 Mg: 35.1 Na: 51.3 K: 15.6  
 DATE SAMPLED: 7-3-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 0.9156 GMS PER CC:  
 SED DRY WT: 0.222 GMS PER CC: INGN LOSS: 6.56

TAXON NAME	Total	No	This Tax	%REL ABUND
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. <i>perpusilla</i>	514.00	4.00	0.778	
9001 <i>Asterionella formosa</i> Hust. v. <i>formosa</i>	514.00	84.00	16.342	
12001 <i>Caloneis bacillum</i> (Grun.) Cl. v. <i>bacillum</i>	514.00	3.00	0.584	
20002 <i>Cyclotella bodanica</i> Eulenst. v. <i>bodanica</i>	514.00	37.00	7.198	
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. <i>stelligera</i>	514.00	1.00	0.195	
23010 <i>Cymbella microcephala</i> Grun. v. <i>microcephala</i>	514.00	5.00	0.973	
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	514.00	2.00	0.389	
27002 <i>Diatoma hiemale</i> v. <i>mesodon</i> (Ehr.) Grun.	514.00	1.00	0.195	
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	514.00	2.00	0.389	
34017 <i>Fragilaria crotonensis</i> Kitton v. <i>crotonensis</i>	514.00	230.00	44.747	
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	514.00	1.00	0.195	
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	514.00	3.00	0.584	
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	514.00	1.00	0.195	
34038 <i>Fragilaria pinnata</i> v. <i>acuminata</i> A. Mayer	514.00	10.00	1.946	
37010 <i>Gomphonema parvulum</i> (Kutz.) Kutz. v. <i>parvulum</i>	514.00	1.00	0.195	
37889 <i>Gomphonema</i> spp.	514.00	2.00	0.389	
44010 <i>Melosira italicica</i> (Ehr.) Kutz. v. <i>italicica</i>	514.00	3.00	0.584	
44011 <i>Melosira italicica</i> ssp. <i>subarctica</i> O. Mull.	514.00	1.00	0.195	
52002 <i>Pinnularia abaujensis</i> v. <i>linearis</i> (Hust.) Patr.	514.00	1.00	0.195	
66014 <i>Synedra parasitica</i> (W. Sm.) Hust. v. <i>parasitica</i>	514.00	2.00	0.389	
66023 <i>Synedra tenera</i> W. Sm. v. <i>tenuer</i>	514.00	2.00	0.389	
67005 <i>Tabellaria flocculosa</i> Roth (Kutz.) strain III sensu	514.00	6.00	1.167	
67009 <i>Nitzschia</i> 42-SN	514.00	1.00	0.195	
99009 <i>Achnanthes</i> 3-SN	514.00	1.00	0.195	
99021 <i>Achnanthes</i> 21-SN	514.00	9.00	1.751	
99038 <i>Cocconeis placentula</i> Ehr. v. <i>placentula</i>	514.00	1.00	0.195	
99067 <i>Epithemia adnata</i> (Kutz.) Breb.	514.00	1.00	0.195	
99087 <i>Fragilaria</i> 6-PIRLA	514.00	3.00	0.584	
99094 <i>Fragilaria</i> 11-SN	514.00	2.00	0.389	
99111 <i>Gomphonema</i> 5-SN	514.00	2.00	0.389	
99133 <i>Navicula cari</i> Ehr.	514.00	1.00	0.195	
99161 <i>Navicula</i> 30-SN	514.00	2.00	0.389	
99164 <i>Navicula</i> 35-SN	514.00	1.00	0.195	
99165 <i>Navicula</i> 36-SN	514.00	1.00	0.195	
99229 <i>Nitzschia frustulum</i> 1-SN	514.00	2.00	0.389	
99230 <i>Nitzschia frustulum</i> 2-SN	514.00	2.00	0.389	
99231 <i>Nitzschia frustulum</i> 3-SN	514.00	1.00	0.195	
99239 <i>Nitzschia</i> 3-SN	514.00	1.00	0.195	
99316 <i>Stephanodiscus parvus</i> Stoermer & Hakansson	514.00	34.00	6.615	
99317 <i>Stephanodiscus</i> 1-SN	514.00	43.00	8.366	
99318 <i>Stephanodiscus</i> 2-SN	514.00	2.00	0.389	
99320 <i>Stephanodiscus</i> 4-SN	514.00	1.00	0.195	
99336 <i>Gyrosigma obtusatum</i> (Sullivan & Wormley) Boyer	514.00	1.00	0.195	

LAKE: BRIGHT DOT LAT DEG: 37 MIN: 32 SEC: 40 LONG DEG: 118 MIN: 51 SEC: 40  
 QUADRANGLE: MT.MORRISON AREA: 11.4 SHED AREA: 140 ELEV: 3194  
 VEG CLASS: 3 ROCK TYPE: % COVER: A: 3 B: 3 C: 1 D: 1  
 pH FIELD: 9.07 pH LAB: 9.03 CONDUCTIVITY: 67 uS@25:  
 HCO<sub>3</sub>: 627 Cl: 2.44 NO<sub>3</sub>: 0 SO<sub>4</sub>: 67.2 MICR EQUIV/L:  
 Ca: 699 Mg: 21.8 Na: 10.8 K: 5.95  
 DATE SAMPLED: 8-18-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.1164 GMS PER CC:  
 SED DRY WT: 0.051 GMS PER CC: INGN LOSS: 28.1

TAXON NAME	Total No	This Tax	%REL ABUND
2004 <i>Achnanthes clevei</i> Grun. v. clevei	514.00	1.00	0.195
2030 <i>Achnanthes minutissima</i> Kutz. v. minutissima	514.00	2.00	0.389
7001 <i>Amphora ovalis</i> (Kutz.) Kutz. v. ovalis	514.00	1.00	0.195
7004 <i>Amphora perpusilla</i> (Grun.) Grun. v. perpusilla	514.00	4.00	0.778
9001 <i>Asterionella formosa</i> Hust. v. formosa	514.00	6.00	1.167
16001 <i>Cocconeis diminuta</i> Pant. v. diminuta	514.00	1.00	0.195
20010 <i>Cyclotella stelligera</i> (Cl. & Grun. V. H. v. stelligera	514.00	47.00	9.144
20012 <i>Cyclotella pseudostelligera</i> Hust. v. pseudostelligera	514.00	12.00	2.335
30003 <i>Diploneis marginestriata</i> Hust. v. marginestriata	514.00	1.00	0.195
34003 <i>Fragilaria brevistriata</i> Grun. v. brevistriata	514.00	2.00	0.389
34016 <i>Fragilaria construens</i> v. venter (Ehr.) Grun.	514.00	74.00	14.397
34022 <i>Fragilaria leptostauron</i> (Ehr.) Hust.	514.00	1.00	0.195
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	514.00	2.00	0.389
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	514.00	101.00	19.650
34027 <i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.	514.00	3.00	0.584
34037 <i>Fragilaria virescens</i> v. <i>exigua</i> Grun.	514.00	1.00	0.195
46014 <i>Navicula cryptocephala</i> Kutz. v. <i>cryptocephala</i>	514.00	19.00	3.696
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	514.00	2.00	0.389
46057 <i>Navicula radiosha</i> v. <i>parva</i> Wallace	514.00	2.00	0.389
48008 <i>Nitzschia dissipata</i> (Kutz.) Grun. v. <i>dissipata</i>	514.00	3.00	0.584
52025 <i>Pinnularia divergens</i> W. Sm. v. <i>divergens</i>	514.00	1.00	0.195
66015 <i>Synedra radians</i> (Kutz.) v. <i>radians</i>	514.00	2.00	0.389
66016 <i>Synedra rumpens</i> Kutz. v. <i>rumpens</i>	514.00	3.00	0.584
66026 <i>Synedra ulna</i> v. <i>danica</i> (Kutz.) V. H.	514.00	2.00	0.389
90990 <i>Nitzschia</i> 42-SN	514.00	2.00	0.389
99007 <i>Achnanthes</i> 1-SN	514.00	3.00	0.584
99009 <i>Achnanthes</i> 3-SN	514.00	1.00	0.195
99012 <i>Achnanthes</i> 7-SN	514.00	1.00	0.195
99013 <i>Achnanthes</i> 8-SN	514.00	6.00	1.167
99025 <i>Achnanthes</i> 31-SN	514.00	1.00	0.195
99087 <i>Fragilaria</i> 6-PIRLA	514.00	4.00	0.778
99088 <i>Fragilaria</i> 2-SN	514.00	20.00	3.891
99090 <i>Fragilaria</i> 7-SN	514.00	46.00	8.949
99094 <i>Fragilaria</i> 11-SN	514.00	73.00	14.202
99107 <i>Gomphonema</i> <i>tackei</i> v. <i>abbreviatum</i> Camburn	514.00	2.00	0.389
99131 <i>Navicula aurora</i> Sov.	514.00	1.00	0.195
99135 <i>Navicula cryptocephala</i> v. <i>venata</i> (Kutz.) Rabh.	514.00	1.00	0.195
99138 <i>Navicula pseudolanceolata</i> Lange-Bertalot	514.00	1.00	0.195
99142 <i>Navicula</i> 1-SN	514.00	10.00	1.946
99154 <i>Navicula</i> 24-SN	514.00	1.00	0.195
99155 <i>Navicula</i> 25-SN	514.00	9.00	1.751
99175 <i>Navicula</i> 46-SN	514.00	1.00	0.195

TAXON NAME	Total No	This Tax	%REL ABUND
99180 Navicula 51-SN	514.00	3.00	0.584
99182 Navicula 53-SN	514.00	3.00	0.584
99183 Navicula 54-SN	514.00	3.00	0.584
99184 Navicula 55-SN	514.00	1.00	0.195
99214 Navicula 85-SN	514.00	1.00	0.195
99229 Nitzschia frustulum 1-SN	514.00	1.00	0.195
99237 Nitzschia 1-SN	514.00	1.00	0.195
99240 Nitzschia 5-SN	514.00	1.00	0.195
99241 Nitzschia 6-SN	514.00	6.00	1.167
99246 Nitzschia 11-SN	514.00	3.00	0.584
99250 Nitzschia 15-SN	514.00	6.00	1.167
99257 Nitzschia 25-SN	514.00	1.00	0.195
99259 Nitzschia 27-SN	514.00	1.00	0.195
99274 Pinnularia subcapitata v. 1-SN	514.00	1.00	0.195
99317 Stephanodiscus 1-SN	514.00	2.00	0.389
99341 Nitzschia amphibioides Hust.	514.00	4.00	0.778

LAKE: BARNEY LAT DEG: 37 MIN: 33 SEC: 47 LONG DEG: 118 MIN: 58 SEC: 2  
 QUADRANGLE: MT.MORRISON AREA: 3.11 SHED AREA: 98.4 ELEV: 3097  
 VEG CLASS: 3 ROCK TYPE: % COVER: A: 1 B: 4 C: 1 D: 1  
 pH FIELD: 8.77 pH LAB: 9.53 CONDUCTIVITY: 56 uS@25:  
 HCO<sub>3</sub>: 392 Cl: 6.68 NO<sub>3</sub>: 0 SO<sub>4</sub>: 157 MICR EQUIV/L:  
 Ca: 497 Mg: 8.68 Na : 80.8 K : 4.97  
 DATE SAMPLED: 8-9-1985 INVESTIGATOR: J.S. AND R.W.H.  
 SED WET WT: 1.436 GMS PER CC:  
 SED DRY WT: 0.0979 GMS PER CC: INGN LOSS: 13.3

TAXON NAME	Total No	This Tax	%REL ABUND
2001 <i>Achnanthes affinis</i> Grun. v. <i>affinis</i>	515.00	1.00	0.194
2016 <i>Achnanthes lanceolata</i> v. <i>dubia</i> Grun.	515.00	1.00	0.194
2030 <i>Achnanthes minutissima</i> Kutz. v. <i>minutissima</i>	515.00	3.00	0.583
7003 <i>Amphora ovalis</i> v. <i>pediculus</i> (Kutz.) V. H. ex Det.	515.00	2.00	0.388
23012 <i>Cymbella minuta</i> Hilse ex Rabh. v. <i>minuta</i>	515.00	2.00	0.388
23015 <i>Cymbella minuta</i> v. <i>silesiaca</i> (Bleisch ex Rabh.) Reim.	515.00	2.00	0.388
34012 <i>Fragilaria construens</i> (Ehr. Grun. v. <i>construens</i>	515.00	19.00	3.689
34016 <i>Fragilaria construens</i> v. <i>venter</i> (Ehr.) Grun.	515.00	111.00	21.553
34023 <i>Fragilaria leptostauron</i> v. <i>dubia</i> (Grun.) Hust.	515.00	1.00	0.194
34025 <i>Fragilaria pinnata</i> Ehr. v. <i>pinnata</i>	515.00	183.00	35.534
34027 <i>Fragilaria pinnata</i> v. <i>lancettula</i> (Schum.) Hust.	515.00	22.00	4.272
34030 <i>Fragilaria vaucheriae</i> (Kutz.) Lange-Bertelot v.	515.00	1.00	0.194
34038 <i>Fragilaria pinnata</i> v. <i>acuminata</i> A. Mayer	515.00	3.00	0.583
46032 <i>Navicula laevissima</i> Kutz. v. <i>laevissima</i>	515.00	2.00	0.388
46051 <i>Navicula pupula</i> Kutz. v. <i>pupula</i>	515.00	1.00	0.194
66014 <i>Synedra parasitica</i> (W. Sm.) Hust. v. <i>parasitica</i>	515.00	1.00	0.194
90990 <i>Nitzschia</i> 42-SN	515.00	1.00	0.194
99007 <i>Achnanthes</i> 1-SN	515.00	1.00	0.194
99014 <i>Achnanthes</i> 9-SN	515.00	1.00	0.194
99088 <i>Fragilaria</i> 2-SN	515.00	17.00	3.301
99090 <i>Fragilaria</i> 7-SN	515.00	96.00	18.641
99093 <i>Fragilaria</i> 10-SN	515.00	1.00	0.194
99094 <i>Fragilaria</i> 11-SN	515.00	9.00	1.748
99131 <i>Navicula aurora</i> Sov.	515.00	2.00	0.388
99157 <i>Navicula</i> 27-SN	515.00	4.00	0.777
99162 <i>Navicula</i> 31-SN	515.00	7.00	1.359
99167 <i>Navicula</i> 38-SN	515.00	1.00	0.194
99229 <i>Nitzschia frustulum</i> 1-SN	515.00	2.00	0.388
99230 <i>Nitzschia frustulum</i> 2-SN	515.00	1.00	0.194
99233 <i>Nitzschia frustulum</i> 5-SN	515.00	5.00	0.971
99240 <i>Nitzschia</i> 5-SN	515.00	1.00	0.194
99257 <i>Nitzschia</i> 25-SN	515.00	1.00	0.194
99316 <i>Stephanodiscus parvus</i> Stoermer & Hakansson	515.00	3.00	0.583
99317 <i>Stephanodiscus</i> 1-SN	515.00	2.00	0.388
99319 <i>Stephanodiscus</i> 3-SN	515.00	4.00	0.777
99341 <i>Nitzschia amphibioides</i> Hust.	515.00	1.00	0.194

LAKE: DANA LAT DEG: 37 MIN: 54 SEC: 35 LONG DEG: 119 MIN: 13 SEC: 7  
QUADRANGLE: MONO CRATERS AREA: 5.18 SHED AREA: 189 ELEV: 3365  
VEG CLASS: 1 ROCK TYPE: % COVER: A: 2 B: 3 C: 1 D: 3  
pH FIELD: 6.65 pH LAB: 6.32 CONDUCTIVITY: 9.9 us@25:  
HCO<sub>3</sub>: 19.3 Cl: 1.28 NO<sub>3</sub>: 9.08 SO<sub>4</sub>: 52.2 MICR EQUIV/L:  
Ca: 63.8 Mg: 8.46 Na : 7.44 K : 2  
DATE SAMPLED: 8-15-1985 INVESTIGATOR: J.S. AND R.W.H.  
SED WET WT: 2.7352 GMS PER CC:  
SED DRY WT: 1.9404 GMS PER CC: INGN LOSS: 1.49