

U.S. GEOLOGICAL SURVEY



Prepared in cooperation with the

CALIFORNIA AIR RESOURCES BOARD



WATER-RESOURCES INVESTIGATIONS REPORT 90-4000

QD 169 H66 1990

ASSESSING THE RESPONSE OF EMERALD LAKE, AN ALPINE WATERSHED IN SEQUOIA NATIONAL PARK, CALIFORNIA, TO ACIDIFICATION DURING SNOWMELT BY USING A SIMPLE HYDROCHEMICAL MODEL

By Richard P. Hooper, Christopher T. West, and Norman E. Peters

U.S. GEOLOGICAL SURVEY

Water-Resources Investigations Report 90-4000

Prepared in cooperation with the

CALIFORNIA AIR RESOURCES BOARD



Doraville, Georgia

1990

DEPARTMENT OF THE INTERIOR MANUEL LUJAN, JR., Secretary

U.S. GEOLOGICAL SURVEY Dallas L. Peck, Director

For additional information write to:

District Chief, WRD U.S. Geological Survey 6481 Peachtree Industrial Blvd. Suite B Doraville, GA 30360 Copies of this report can be purchased from

Open-File Services Section Western Distribution Branch U.S. Geological Survey Box 25425, Federal Center Denver, CO 80225

CONTENTS

Page
Abstract1
Introduction 1
Purpose and scope 2
Site description
Background 5
Acknowledgments6
Model structure
Hydrologic model 6
Chemical model10
Alkalinity generation10
Nitrogen cycle12
Other chemical species
Solution technique13
Model Inputs
Model Results16
Nonreactive solutes
Weathering products
Deposition scenarios
Summary
Conclusions
References cited
Appendix A. Sample input/output from computer model28
Appendix B. Model listing

ILLUSTRATIONS

Figure	1. Map showing location of study site.	3
	2. Map showing subbasins in Emerald Lake watershed	4
	3. Schematic diagram of hydrologic model	7
	4. Volume of ice, epilimnion, and hypolimnion in 1986 and 1987	9
	5. Sum of base cations and silica concentration in stream-water samples collected from all sampled inlets to Emerald Lake in 1986	11
	6. Ratio of weathering products to hydrogen ion concentration and 10-day average discharge	12
	7. Laboratory and modeled solute release from snowpack	15
	8. Observed and predicted concentrations of solutes at inflows 1 and 2	19
	9. Observed and predicted concentrations of solutes in the lake outflow .	20
	10. The effect of halving and doubling current nitrate and sulfate deposition under two different elution scenarios on alkalinity of the lake outflow	23

TABLES

Page

Table	1. Characteristics of terrestrial subbasins
	2. Model inputs
	3. Input-output budget for snowmelt season
	4. Multiples of current deposition loadings of ammonium nitrate and sulfuric acid required to exhaust the alkalinity of Emerald Lake

<u>Page</u>

CONVERSION FACTORS

For those readers who may prefer to use inch-pound units rather than metric (International System) units, conversion factors for the terms used in this report are listed below:

Multiply metric unit	<u>by</u>	<u>To obtain inch-pound units</u>
	Length	
millimeter (mm)	0.03937	inch (in.)
meter (m)	3.281	foot (ft)
kilometer (km)	0.6214	mile (mi)
	Area	
hectare (ha)	2.471	acre
square meter (m ²)	10.76	square foot (ft ²)
	Volume	
liter (L)	0.2642	gallon (gal)
cubic meter (m ³)	35.31	cubic foot (ft ³)
	Flow	
cubic meter per day (m ³ /d)	264.2	gallon per day(gal/d)
liters per second (L/s)	0.0353	cubic feet per second
	Sea Level	

In this report, "sea level" refers to the National Geodetic Vertical Datum of 1929 (NGVD of 1929)—a geodetic datum derived from a general adjustment of the first-order level nets of both the United States and Canada, formerly called Sea Level Datum of 1929.

ASSESSING THE RESPONSE OF EMERALD LAKE, AN ALPINE WATERSHED IN SEQUOIA NATIONAL PARK, CALIFORNIA, TO ACIDIFICATION DURING SNOWMELT BY USING A SIMPLE HYDROCHEMICAL MODEL

By Richard P. Hooper, Christopher T. West, and Norman E. Peters

ABSTRACT

A sparsely parameterized hydrochemical model has been developed by using data from Emerald Lake watershed, which is a 120-hectare alpine catchment in Sequoia National Park, California. Greater than 90 percent of the precipitation to this watershed is snow; hence, snowmelt is the dominant hydrologic event. A model which uses a single alkalinity-generating mechanism, primary mineral weathering, was able to capture the pattern of solute concentrations in surface waters during snowmelt. An empirical representation of the weathering reaction, which is based on rock weathering stoichiometry and which uses discharge as a measure of residence time, was included in the model. Results of the model indicate that current deposition levels would have to be increased between three- and eight-fold to exhaust the alkalinity of the lake during snowmelt if there is a mild acidic pulse in the stream at the beginning of snowmelt as was observed during the study period. The acidic pulse in the inflow stream at the onset of snowmelt was less pronounced than acidic pulses observed in the meltwater draining the snowpack at a point using snow lysimeters or in the laboratory. Sulfate concentrations in the streamwater were the most constant; chloride and nitrate concentrations increased slightly at the beginning of snowmelt. Additional field work is required to resolve whether the an acidic meltwater pulse occurs over a large area as well as at a point (implying sulfate-regulating mechanisms in the soil) or whether, due to physical and chemical processes within the snowpack, the acidic meltwater pulse is attenuated at the catchment scale. The modest data requirements of the model permit its applications to other alpine watersheds that are much less intensively studied than Emerald Lake watershed.

INTRODUCTION

The Emerald Lake watershed, an alpine catchment in Sequoia National Park in the Sierra Nevada of California, has been the subject of an intensive 5-year research effort under the Integrated Watershed Study (IWS) of the California Air Resources Board. All aspects of the watershed—its hydrology, geology, soils, vegetation and aquatic biota—were studied to gain an understanding of the sensitivity of the ecosystem to acidification (California Air Resources Board, 1986). Because over 90 percent of the precipitation in this watershed is in the form of snow, the emphasis of the field program was on the response of the catchment to snowmelt. Field data were collected starting in October, 1985, but data from 1986 and 1987 are more complete than from the first year and they reflect the refinement of snow-measurement techniques from information gained during the first year of sampling.

Purpose and Scope

One phase of the IWS was the development of simulation models to predict surface-water chemistry. These models were to serve as research tools to integrate information gained from the field program and to identify gaps in understanding the chemical mechanisms controlling surface-water chemistry, and as predictive tools to provide a means for forecasting the chemical response of the watershed to different depositional loadings of atmospheric acidity. This paper presents a lumped, sparsely parameterized simulation model, called the Alpine Lake Forecaster, or ALF, that was developed from the IWS data. ALF is a model of intermediate complexity when compared with a fully distributed model (Sorooshian and others, 1989) and a regional lake acidification model (Nishida and others, 1989), which also were developed from this data set.

ALF was designed and calibrated from data collected during water year 1986, which extended from October 1, 1985, to September 30, 1986. This year was an exceptionally heavy snow year, with twice the usual snow accumulation, and a maximum snow depth of more than 5 meters (m). Water-quality samples of surface water were taken on a biweekly basis from the onset of snowmelt throughout summer and fall (Dozier and others, 1989). Intensive sampling of infrequent summer rainstorms was performed only opportunistically (Melack and others, 1989). Snow samples were taken on an event basis during winter, and a whole-basin survey was performed at maximum snow accumulation to estimate the water volume and ionic loadings to the catchment. Data from 1987, a light snow year with one-quarter the amount of snow as 1986, were used for validation of the model. The sampling regime was for surface water samples was similar to that in 1986. Based upon the experience gained from the previous year, only a snow survey at maximum accumulation was performed; no event snow samples were collected. Data considered in construction of the model included stream- and lake-water quality and quantity, snowpack quality and quantity, soil column studies, and geochemical characterizations of the bedrock.

Site Description

The following site description, except where noted, is excerpted from Dracup and others (1989) to which the reader is referred for more details. The Emerald Lake watershed (36° 35' 49" N, 118° 40' 30"W) is within Sequoia National Park on the western slope of the Sierra Nevada in the Marble Fork of the Kaweah River drainage, approximately 5 kilometers (km) east-southeast of Lodgepole (fig. 1). The area of the watershed is about 120 hectares (ha) and altitudes range from 2,800 m at the lake outlet to 3,416 m at the top of Alta Peak. The watershed is an alpine cirque that has the typical bowl shape and more than 180 degrees of aspect. The long axis of the watershed (approximately 1.7 km in length) is oriented from southeast to northwest, and more than half of the watershed has a north to northeast aspect (fig. 2).

Bedrock underlying the Emerald Lake watershed consists of granite, granodiorite, and aplite (Clow, 1987). The granite consists of 2 to 7 percent biotite and hornblende by volume; however, the percentage of these more readily weathered minerals in the granodiorite ranges between 12 and 17 percent by volume. More than one-third of the basin consists of exposed bedrock, and most of the remainder is covered by a thin mantle of talus or colluvium. Only 20 percent of the basin is covered by soils (Lund and others, 1987). Bedrock comprising the basin floor and cliffs has been exposed recently by glacial erosion and frost action and, thus, is weathered only slightly. Both subalpine and alpine vegetation are present; western white pine (*Pinus monticola*) is the predominate tree species and willows (*Salix sp.*) are the dominant shrub.



Figure 1.— Map showing location of study site.



Figure 2.—Map showing subbasins within Emerald Lake watershed.

4

Precipitation is seasonal in the alpine regions of Sequoia National Park. Stephenson (1988) estimated that 18 percent of the annual precipitation falls from September through November, 50 percent from December through February, 30 percent from March through May, and only 2 percent from June through August. Snowmelt begins in March or April and continues through June, except in the heaviest snow years when melt continues through August. Consequently, few rain-on-snow events occur. More commonly during late spring and early summer, snowstorms occur after the melt has begun.

Emerald Lake is a tarn fed by three major and five minor stream channels; only inflows 1, 2, and 4 (fig. 2) continue to flow after the snow has melted in their respective drainage basins. Two of these perennial or nearly perennial streams (inflows 1 and 2, fig. 2) are split parts of a larger stream that drains along the main axis of the basin; this drainage accounts for 60 percent of the total inflow area. For modeling purposes, the drainage area of inflows 1 and 2 was designated as a single unit. Outflow from the watershed seems to occur only through the lake outlet. The geomorphometry of the lake suggests that the low, transverse ridge of bedrock that runs along the master joint trace on the valley floor dams the lake and is impermeable. There is no visual evidence of ground-water seepage at land surfaces downslope of the lake. Seepage losses from the watershed are assumed to be negligible.

Background

Alpine watersheds pose a number of challenges to the construction of hydrochemical models. Snow accounts for the vast majority of the hydrologic inputs to these systems, and therefore, snowmelt and not rain, is the important hydrologic event to be modeled. Previous models, such as the Birkenes model (Christophersen and others, 1984) and the Integrated Lake-Watershed Acidification Study (ILWAS) model (Chen and others, 1983), have used degree-day formulations to predict snowmelt. These models are appropriate for forested catchments where solar radiation and wind are not substantial components in the energy budget of the snowpack (Miller, 1950). In sparsely vegetated alpine watersheds, a more complete evaluation of the energy budget is required, but measurement of the necessary radiation values is logistically difficult. A major part of the field effort at Emerald Lake was devoted to obtaining these data, which have been used in an energy balance model of snowmelt at one point (Marks, 1988). The application of the point model to the complex terrain of the Emerald Lake watershed is an area of active research (Dozier, 1988).

Even with the prediction of meltwater generation at the surface of the snowpack, routing of water through the pack is difficult to simulate. In watersheds for which models have been developed (for example, the Adirondacks, southern Norway) and where snowpacks are on the order of 1 m deep, observations can be made as to whether the majority of snowmelt infiltrates the soil (i.e., there is no basal ice layer) or flows overland. Such observations are difficult, if not impossible, to make in alpine terrain; thus water movement through the pack is not well understood and modeling this process remains a matter of conjecture.

Moreover, existing process-oriented hydrochemical models (the ILWAS and Birkenes models) rely on a hydrologic accounting scheme that requires an accurate measure of the timing and quantity of the entry of snowmelt or rain into the watershed. Predicting or measuring snowmelt is much more difficult than rainfall, particularly in deep snowpacks. Any error in the estimates of timing and magnitude of meltwater generation will lead to errors in parameterization of the hydrologic part of the model, such as the capacity of soil layers or the rates of release from the soil layers (for example, their conductivity or other rate constant). Thus, errors in the hydrologic model can influence chemical parameterization and predictions.

A hydrologic accounting model also requires that a hydrologic budget, which divides the water input to the watershed into outflow, evapotranspiration losses and change in storage within the watershed, can be constructed with little error. For water year 1986, an acceptably precise hydrologic budget (less than 5 percent error) was obtained for the entire melt period (Dozier and others, 1989). However, when finer time steps were considered, the errors were too large for modeling purposes. A simple extrapolation of the point-snowmelt model (Marks, 1988) to the entire watershed yielded incorrectly timed meltwater inputs. Furthermore, the water yield from the snowpack, as determined from sequential snow surveys, was inconsistent with the measured lake outflow (Dozier and others, 1989). This lack of agreement probably was due both to errors in the snow survey, which were exacerbated owing to the great depth of the pack, and to errors in the stage-discharge relation, which had to be extrapolated to high stage values that were beyond the range of calibration. The difficulties in achieving an acceptable hydrologic budget led to consideration of a simpler model structure that did not depend on a hydrologic budget, but that could still be used to achieve the objectives of the modeling program.

Acknowledgments

The authors would like to thank K.A. Tonnessen of the California Air Resources Board for her persistence and organizational abilities. J.M. Melack and Jeffrey Dozier provided useful criticisms of the model and insights into the hydrologic and chemical mechanisms operating at Emerald Lake. Mark Williams, Richard Kattelmann, and J.O. Sickman responded to our many data requests and assisted in the interpretation of these data. This work was performed in cooperation with the California Air Resources Board.

MODEL STRUCTURE

ALF consists of a coupled hydrologic and chemical model. The hydrologic model, which provides the basic structure for the model, determines the quantity of discharge from each terrestrial subbasin to the lake and the volume of the lake epilimnion into which the streamwater mixes. The chemical model calculates the equilibrium concentration of the solutes for the streamwater and the lake water based upon primary mineral weathering and dissolution of carbon dioxide. These models are described in more detail in the following sections.

Hydrologic Model

To eliminate the need for a hydrologic accounting model, discharge was assumed to equal snowmelt on a daily basis. This assumption implies that there is no change in lake or ground-water storage during the snowmelt period, which is not so unrealistic as it may seem. The lake stage varies between 1 and 2 millimeters (mm) on a weekly basis (Richard Kattelmann, Univ. of California, Santa Barbara, written commun., 1987), which is equivalent to a volume of approximately 50 cubic meters (m³). Daily outflow from the lake during snowmelt ranges between 1,000 and 35,000 m³. Ground-water storage in unconsolidated surficial deposits has been estimated at 100,000 ± 50,000 m³ (Dracup and others, 1989). During the first part of snowmelt, some meltwater recharges the ground-water system, which is depleted during winter. A schematic of this simple hydrologic model is shown in figure 3. The total daily runoff from the terrestrial part of the watershed is set equal to the measured daily discharge from the lake.



Figure 3.—Schematic diagram of hydrologic model.

Because the snowpack melts at different rates depending on the aspect and slope of the land surface, the watershed was divided into seven subbasins (fig. 2). The proportion, P_i (t), of total discharge, Q(t), for each subbasin *i* on day *t* is determined by the relative amount of solar radiation to that subbasin

$$P_{i}(t) = \frac{r_{i}(t)A_{i}}{\sum_{i=1}^{n} r_{i}(t)A_{i}}$$
(1)

The variable r_i is the potential solar radiation per unit area as determined by an algorithm of Swift (1976), and it is a function of latitude, slope inclination, aspect, and day of year. The variable A_i is the snow-covered area of the *i*th subbasin; and *n* is the total number of subbasins in the watershed. Each subbasin is credited at the start of the snowmelt season with part of the total seasonal discharge from the lake during the entire snowmelt period based on the proportion of the subbasin area; and each subbasin meets the discharge demand (i.e., $P_i(t)Q(t)$) as long as enough water remains in the subbasin. After the subbasin depletes its store of water, the discharge is apportioned among the remaining subbasins. If the snow-covered area is not known, the subbasin area can be used, although this tends to overestimate the contribution from subbasins with relatively less snow-covered area than other subbasins. Snow-covered area was measured at Emerald Lake, but the values by subbasin were not available in time for this modeling exercise; therefore, the subbasin area was used for the results presented in this report. The values required for the application of equation (1) to the seven subbasins are shown in table 1.

Subbasin	Area (10^3 m^2)	Percent of total area	Slope (degrees)	Aspect (degrees)	
West joint	140	12	34	43	
SW gully	133	11	31	27	
Inflow 3	141	12	28	10	
Inflow 1&2	396	33	34	305	
NE gully	245	20	27	275	
Pinestand	135	11	34	253	
Meadow	9.28	0.8	14	327	

Table 1. Characteristics of terrestrial subbasins

 $[10^3 \text{ m}^2$, thousands of square meters]

Because there is not an accounting model with explicit reservoirs, routing of rainfall that occurs during the melt period had to be determined from the discharge. If the discharge increased on a day when rain was recorded, all of the increase in discharge was assumed to be rain, if there was a sufficient volume of rain. If not, the volume unaccounted for was supplied from the snow pack. If more rain fell than the discharge increased (as was usually the case), the "excess" rain was areally proportioned among the subbasins and added to the snowpack. Rainfall volume was minor in 1986 but provided more than 10 percent of the precipitation in 1987.

The thermocline and ice depth in the lake are determined by linearly interpolating the field values that were measured monthly in winter and biweekly during the melt period and summer. The depths are converted to volumes by using the lake bathymetry provided by J.O. Sickman (Univ. of California, Santa Barbara, written commun., 1987). The ice layer is assumed to be completely contained within the epilimnion, thus reducing the volume of this layer. The volumes of all three layers for the years that were modeled are shown in figure 4. Streams from the terrestrial part of the watershed flow into the epilimnion and the lake outflow is from the epilimnion; the hypolimnion is hydraulically separated from the surface water, and does not contribute to streamflow. Each layer is assumed to be completely mixed.

This simple hydrologic model structure eliminated artificial parameters that would have had to be fitted to the data, as is the case with the previously mentioned hydrochemical models. As long as the assumption holds that there is no change in storage of water in the lake or in the aquifers, this hydrologic model will provide a more powerful test of the proposed chemical mechanisms, in terms of a hypothesis test, than a model that contains artificial parameters.



Figure 4.—Volume of ice layer, epilimnion and hypolimnion, 1986 and 1987.

Chemical Model

The chemical reactions that are contained in ALF can be divided into those which generate alkalinity, those which are part of the biological nitrogen cycle, and others which include carbonate buffering and non-reactive solutes. The representation of these reactions included within ALF are described in the next three sections. The description of the chemical model concludes with a discussion of the numerical techniques used to solve to simultaneous equations which constitute the chemical model.

Alkalinity generation

The dominant sources of alkalinity in watersheds with young soils that contain fresh mineral faces and few sesquioxides, are primary mineral weathering and cation exchange (Drever, 1988, p. 207-231). Weintraub (1986) characterized the mineralogy and chemistry of soils, surficial material, and bedrock from Emerald Lake, and conducted laboratory leaching studies of undisturbed soil columns. Weintraub (1986) concluded that weathering was the dominant alkalinity-generating mechanism in the watershed. Lund and others (1987) further characterized the soil chemical characteristics and they concluded that the Emerald Lake soils are acidic (pH from 4.4 to 4.8) and have very low percent base saturation, which may only provide limited buffering of acidic inputs. In addition, the interrelation of solute compositions of streamwaters were evaluated by Lund and others (1987) from which they concluded that primary mineral weathering is a primary source of alkalinity. We, therefore, included mineral weathering in the model but did not include cation exchange.

The mineralogy of the bedrock, which determines the rate and stoichiometry of primary weathering, is dominated by quartz and feldspar and contains minor amounts of biotite and hornblende (Weintraub, 1986). A variety of secondary vermiculite minerals were identified in Emerald Lake soils (Weintraub, 1986). Incongruent weathering of aluminosilicate minerals produces secondary minerals, base cations, and dissolved silica (SiO₂) either while consuming hydrogen ion or releasing alkalinity. In the absence of detailed mineral-weathering information, the stoichiometry of the mineral weathering was simplified by combining the individual aluminosilicate minerals into one reactant called rock. Rather than tracking each base cation individually, the base cations (Na⁺, K⁺, Ca²⁺, and Mg²⁺) were combined as the sum of base cations, herein called SBC. The slope of the least squares regression of SBC concentration (in microequivalents per liter (μ eq L⁻¹)) on dissolved silica (in micromoles per liter(μ mol L⁻¹)) for the 1986 inflow data (fig. 5) is 1.2 (r² is 0.77), and therefore, the following incongruent dissolution reaction is consistent with these data

Rock +
$$1.2H^+ \rightarrow 1.2SBC^+ + SiO_2$$
 (aq)+ secondary mineral. (2)

This reaction, however, is subject to kinetic constraints. The hydrologic model does not permit an explicit calculation of the residence time of water within the terrestrial system, so a standard kinetic model cannot be fit to the data. Instead, we use the lake discharge, averaged over the 10 days prior to the sample collection, as a surrogate for residence time. A regression of the log of the reaction constant, K, where





$$K = \frac{[SBC^+]^{1.2}[SiO_2]}{[H^+]^{1.2}}$$
(3)

on the average discharge (fig. 6) yields the following equation,

$$K = 0.0082 \cdot 10^{-0.000044} Q' \tag{4}$$

where Q' is the 10-day average discharge. Equation 4 explains 80 percent of the variation in the data. Because this is the central equation in the model, two different intercepts, 0.005 and 0.02, that bracket the observations were used to assess the sensitivity of the results to this equation (fig. 6). Model results based on the lower line are referred to as the "lower bound" case because these results have lower concentrations of alkalinity, SBC, and silica than the "upper bound" case. The watershed is more sensitive to acidification under the lower bound case because less of the hydrogen ion is neutralized by the weathering reaction.



concentration and 10-day average discharge.

Nitrogen cycle

A simplified representation of the nitrogen cycle is included in the model. Ammonium concentration in the main inflow is very low, generally much less that 1 μ eq/L. Therefore during biotically active seasons, all ammonium is assumed to be taken up by the biota; each mole of ammonium is exchanged for one mole of hydrogen ion. In addition, a part of the nitrate, v, also is retained along with an equivalent amount of hydrogen ion. When the biota are not active, all the ammonium is oxidized to nitrate, which is passed through the system, and to two moles of hydrogen ion; no nitrate is taken up by the biota. After biotic uptake, if any, nitrate is treated as a conservative species. We assumed that the biota were active throughout the snowmelt period and set v to 0.10 (that is, 10 percent of the nitrate is retained by the biota).

Other chemical species

A standard representation of the carbonate buffering system is included. When the lake is covered by ice, the system is assumed to be closed to the atmosphere; during the lake's ice-free season and during all times in the terrestrial part of the watershed, the system is assumed to be open to the atmosphere. The partial pressure of carbon dioxide (CO_2), to which water equilibrates in the open system, is a parameter in the model to be fit. Chloride and sulfate are treated as conservative species in the model.

Solution technique

A modified Newton's method (Morel and Morgan, 1972) was used to solve the system of nonlinear simultaneous equations that describe weathering and carbonate buffering in the terrestrial subbasins. There are four variables in the system, hydrogen ion, bicarbonate, silica, and SBC, that are determined from charge balance constraint, weathering equilibrium, weathering stoichiometry, and carbonate equilibrium. Temperature corrections are made for the carbonate system and for the hydrolysis of water; activity corrections are not made owing to the low ionic strength of the surface waters. The mathematical solution seems to be identical to an equilibrium solution, yet the equilibrium "constant" for the weathering reaction changes as discharge changes. These equations are solved on a daily time step by using the input concentration of the meltwater. On days when rain fell, the "excess" rain, whose chemistry is an input to the model, is assumed to react with the rock and then mix with the meltwater which has also reacted with the rock. This mixture is then re-equilibrated before flowing into the lake.

Within the lake, no chemical reactions other than carbonate buffering occur. The volume of the epilimnion is increased by inflow from the terrestrial system, thinning of the ice layer, and deepening of the thermocline. The appropriate quantities of solutes are advected with the water, and the system is re-equilibrated assuming that the epilimnion is completely mixed. The time step concludes with water of the same quality as the epilimnion flowing out of the lake.

Within the chemical model, there are only five parameters that must be calibrated: the two regression coefficients for the weathering reaction (equation 4), that part of nitrate retained by the biota (v), the date at which the biota become active, and the partial pressure of CO₂. This latter parameter must be fitted because surface waters frequently contain more dissolved CO₂ than would be predicted from equilibrium with atmospheric levels of CO₂. Typical partial pressures range from two to five times the atmospheric partial pressure of CO₂. Dissolved inorganic carbon was not measured in surface-water samples collected from Emerald Lake. The equilibrium constants and enthalpy values (used for temperature correction) for the carbonate system and for water hydrolysis are considered to be well known and are not subject to calibration.

Model Inputs

ALF was designed to utilize data collected for the Integrated Watershed Study. However, its input requirements are modest enough to allow its application to less intensively studies sites. Table 2 shows the hydrologic and chemical inputs necessary to run the model. The most critical measurements are the daily discharge from the lake and the mass of solutes contained in the snowpack at maximum accumulation. Rain water quality primarily affects stream and lake water quality on the day which the event happens. However, if these is any "excess" rain as defined above, the solutes associated with this rain are incorporated into the snowpack. Thermocline depth and ice thickness determine the hydraulic mixing volume of the lake and are of secondary importance. Water temperatures are used to correct thermodynamic equilibrium constants and also are not critical. The remaining inputs are used to determine initial conditions for the model.

Гable	2.	Model	Inputs
-------	----	-------	--------

Hydrologic	Chemical
Daily discharge from lake	Total mass of solutes at maximum
Thermocline depth ¹	snowpack accumulation
Ice thickness ¹	Rainwater quality
Temperature of inflows ¹	Initial lake water quality
Temperature of epilimnion ¹	Initial chemical composition of ice on lake

¹ May be interpolated from field measurements to derive daily values necessary to run model.

Although the mass of solutes in the snowpack is an input to the model, the meltwater quality during the snowmelt must be calculated. The first meltwater from the snowpack typically contains a high concentration of solutes (Johannessen and Henriksen, 1978). This first "pulse" of solutes from acidic snowpacks is believed to be the cause for the sharp pH depressions during the spring snowmelt (Skartveit and Gjessing, 1979), and has been linked to fish kills in Norway (Leivested and Muniz, 1976). Therefore, the intensity of the acidic pulse is a key factor in determining the impact of acidic pollutants contained within the snowpack on aquatic biota. Both the Birkenes and ILWAS models assume an exponential decline in the release of solutes from the snowpack. Specifically, the concentration factor (the ratio of the concentration of the meltwater to the bulk concentration of the snowpack) is assumed to decline exponentially. Because the pattern of solute release is so critical, particularly when considering different depositional scenarios, we chose to parameterize the solute release differently. An elution rule in the form of x percent of the solutes released in the first y percent of the meltwater (for example, 80/20 rule means 80 percent of the solutes are released in the first 20 percent of the meltwater) and a maximum concentration factor for the first day (the ratio of concentration on that day to the average concentration of the solute in the snowpack) are specified. Two exponential curves, one for the period when the first y percent of the water drains from the pack and a second curve for the remainder of the melt period, define the meltwater concentration. These curves meet the specified elution rule and concentration factor along with a continuity condition that the concentration does not change abruptly between the two curves. The results of this approach are compared with the experimental data of R. C. Bales (Univ. of Arizona, written commun., 1988) on a cumulative basis in figure 7. All solutes are treated identically; no provisions are made for "preferential" elution (that is, different solutes being released at different rates), which has been reported by Brimblecombe and others (1987).



Figure 7.—Laboratory and modeled solute release from snowpack.

Because snowpacks melt at different rates in each subbasin owing to different radiation inputs, meltwater concentration curves are calculated for each subbasin. Consequently, ionic pulses from subbasins where snowmelt occurs later in the season are contributed to surface waters after earlier ionic pulses from subbasins where snowmelt has already occurred. This distribution of meltwater generation leads to a much less intense ionic pulse to the lake than if the entire snowpack melted as one unit. The total ionic loading in the snowpack for all ions was estimated by Dozier and others (1987) from field observations (table 3). An artificial solute called was created to create an exact charge balance. Typically there is a small cation charge excess which arises from the unmeasured contribution of organic anions to the charge balance of the solution. If this solute were not created, the charge balance constraint in the chemical model would adjust the hydrogen ion concentration to meet the charge balance. The concentration of this artificial solute was always very small.

MODEL RESULTS

Errors in the prediction of one charged solute will affect predictions of other solutes owing to the electroneutrality constraint. Therefore, predictions for the nonreactive solutes, those that were modeled as not participating in any chemical reaction, will be discussed first. Then, solutes which are the products of the weathering reactions will be considered. Finally, the result of different deposition scenarios will be presented.

Nonreactive Solutes

The ionic loadings in the snowpack from field estimates are compared in table 3 with the measured export from discharge through the lake outlet during the snowmelt period. There are large differences between 1986 and 1987 both in the amount of precipitation and in its form. Rain inputs of water and solutes were negligible in 1986, but accounted for 18 percent of the water in 1987 and more than half the deposition of all solutes, except chloride. For the nonreactive solutes (sulfate and chloride in this model), the input should equal the output; clearly, this is not the case for sulfate and chloride during 1986 and 1987. The ionic deposition is calculated as the product of the ionic concentration and the volume of water, both of which are subject to measurement error.

Extensive snow surveys were performed to estimate the water volume of the snowpack. Techniques were refined and sampling intensity increased during 1986 and 1987 over the initial effort in 1985, as is evidenced by the good agreement in the water budget. Snowpack solute concentration was measured at eight points throughout the watershed, although not all sites were sampled for all events. Therefore, the discrepancy in the ionic budgets must be due either to sampling and analytical errors or to additional sources and mechanisms that regulate solute concentrations.

Sulfate input is less than one-half the sulfate output for 1986; yet the output exceeds the input for 1987. The difference between the form of the input (snow in 1986 and predominantly rain in 1987) may account for the difference, but there is insufficient data to make any conclusions. These budgets ignore input due to atmospheric dry deposition of sulfate and internal sources of sulfate. Contributions of sulfate by dry atmospheric deposition were evaluated using deposition to pine branches by Bytnerowicz and Olszyk (1988). An upper-bound estimate for the dry deposition of sulfate may exist. Volcanic glass was identified in the coarse silt fraction of soils at Emerald Lake (Wyles, 1986). Strong acid treatment of these soils resulted in sulfate release, which was attributed to the breakdown of vesicles that possibly contain sulfur dioxide. However, no quantitative estimate of this contribution to the sulfur budget has been made. Modifications to the model await further field observations of sulfur sources and regulating mechanisms.

The chloride budget is also variable, as might be expected, given the numerous opportunities for contamination during sample handling and the analytical precision at the low concentrations. The snow input number for 1986 may be somewhat high because there was a known chloride contamination from the plastic bags used for sampling that year. It was not possible to determine quantitatively the amount of contamination (M.W. Williams, Univ. of California, Santa Barbara, written commun., 1989). The chloride budget for 1987 would suggest, however, that the precision of these load estimates may be as low as ± 50 percent if the variation in the chloride estimates arise primarily from sampling error and not from contamination or analytical imprecision.

Despite these differences in the budgets, the measured snowpack loadings for all solutes were used in the model. Although the differences on a seasonal basis appear to be large, the amount of error introduced in daily concentrations is only a few microequivalents per liter.

Year		Water Volume	SBC	NH4 ⁺	NO3-	SO4 ²⁻	Cl-
		$(10^6 \mathrm{m}^3)$	(10 ³ eq)				
1986	Snow input	1 2.4	7.5	2.9	4.1	3.4	7.4
	Rain input ²	<u>0.05</u>	<u>0.69</u>	<u>0.82</u>	<u>0.82</u>	<u>0.66</u>	0.24
	Total input	2.4	8.2	3.7	4.9	4.1	7.7
	Output ³	<u>2.3</u>	<u>63</u>	<u>0.1</u>	<u>9.1</u>	<u>9.9</u>	<u>5.6</u>
	Difference ⁴	0.1	-55	3.6	-4.2	-5.8	2.1
	Ratio ⁵	.96	7.7	.027	1.9	2.4	0.73
1987	Snow input	¹ 0.80	3.2	3.0	3.4	2.3	1.2
	Rain input ²	<u>0.17</u>	<u>7.3</u>	<u>6.8</u>	<u>5.9</u>	<u>5.6</u>	<u>0.85</u>
	Total input	0.97	10.	9.8	9.3	7.9	2.0
	Output ³	.75	<u>34.</u>	<u>0.01</u>	<u>6.3</u>	<u>5.2</u>	<u>3.2</u>
	Difference ⁴	0.22	-24	9.8	3.0	2.7	-1.2
	Ratio ⁵	0.80	3.4	.001	0.68	0.66	1.6

Table 3. Input-output budget for snowmelt season

[10⁶ m³, millions of cubic meters; 10³ eq, thousands of equivalents]

¹As determined from snowpack loadings at maximum accumulation (Dozier and others, 1989).

²Includes rain and mixed rain and snow events that occurred during the melt period (Dozier and others, 1989) which ran from April 16, 1986 to September 30, 1986 and from March 25, 1987 to July 21, 1987.

³As determined from mass export through lake outlet; water-quality observations were linearly interpolated to a daily basis.

⁴Input less output; a positive value indicates net retention in the watershed and a negative value, a net release.

⁵The ratio of output to input; values greater than one indicate that the watershed is a source of the solute and values less than one indicate that the watershed is a sink.

The results of the model simulations are presented in figures 8 and 9 for the combined inflows 1 and 2 and the lake outflow, respectively. Model results for streamwater quality cease when the model calculates that all the snow in the subbasin has melted; this occurred in mid July 1987. The "spikes" in concentration during the simulation period occur on during rain storms. Clearly these events were more important during 1987 than during 1986 as was also indicated by the budgets. The observations for the combined inflows (fig. 8) are an average of the measurements taken on the individual streams if more than one inflow was sampled on a particular day, or if more than one sample were collected during the day.

The upper three panels of each plot are for solutes that do not participate in the weathering reaction (Cl⁻, NO₃⁻ and SO₄⁻²). Three different elution scenarios are considered: an 80/20 release (80 percent of the solutes in the first 20 percent of the meltwater) with a maximum concentration factor (CF) of 8; a 40/20 release with a CF of 3; and a uniform release. For chloride, the 40/20 release rule seems to fit best; for nitrate, either the 40/20 for 1987 or 80/20 release rule for 1986 fits best. For sulfate, a uniform release fits the pattern best in 1986, although there is a systematic underprediction due to the net export of sulfate during this year. In 1987, the 40/20 rule appears to fit the initial stage of snowmelt best, yet during the rain events, the sulfate concentration does not respond as much as would be predicted by the model using either the 40/20 or 80/20 rule. The uniform release also does not describe the observations. In this case the sulfate added by the rain (twice as much as was in the snow, see table 3), continues to increase the stream concentration to levels above those observed. If there is a differential elution of sulfate (that is the 40/20 or 80/20rule), then the stream sampling missed extreme changes in concentration. However, these concentrations would have to be much larger than those ever observed at Emerald Lake. Alternately, sulfate is not a nonreactive solute and is being retained in the watershed in 1987. Among the three solutes, sulfate exhibits the smallest range in concentration. This pattern with sulfate showing the least pronounced elution is opposite the preferential elution patterns observed by Brimblecombe and others (1987) where sulfate eluted first.

Putting aside the large rain-on-snow events in 1987 for the moment, consider 1986 and the April 1987 when an initial pulse from the snowpack would have been expected but did not occur. Two possible explanations can be advanced. One explanation is that solutes are eluted uniformly from the snowpack and that there are additional sources of nitrate and chloride, which account for the observed increases in concentration in the streamwater. Supporting this explanation is the observation that the peak nitrate concentrations observed in the inflow streams are roughly equivalent in 1986 and April, 1987 (approximately 12 μ eq L⁻¹), yet the average concentration in the snowpack (assuming 10 percent and 100 percent biotic retention of nitrate and ammonium, respectively) is twice as large in 1987 (5 μ eq L⁻¹) as in 1986 (2.5 μ eq L⁻¹). The observed chloride pulse also may come from the soil solution where chloride from weathering or from the decomposition of organic matter may be concentrated during the winter (Peters, 1987).

A second explanation is that chloride and nitrate pulses arise from differential elution from the snowpack, but sulfate pulses from the pack are not observed in the stream owing to regulation by adsorption/desorption processes in the soils. Although soils cover only a small part of the watershed area, most flowpaths to the lake probably intersect the soils at some point. The constancy of the sulfate concentrations during the rain events in 1987 would also support sulfate regulation by the soils. However, such a regulating mechanism for sulfate has not been identified at Emerald Lake and would be unusual in such young soils.

The fundamental conclusion, regardless of the specific mechanism that accounts for the observed concentrations, is that the effective "acid pulse" to surface waters from this alpine snowpack is not as severe as would be predicted from laboratory studies or from point measurements as determined by snow lysimeters. For evaluation of the weathering model and the subsequent scenario analysis, the uniform and the 40/20 release rules will be considered.









Weathering Products

The fourth through the seventh panels in figures 8 and 9 show the observed and the model predicted values for solutes affected by the weathering reactions (alkalinity, hydrogen ion, SBC, and silica). Two different solute release rules, uniform release and 40/20 release with a CF of 3, are considered. Model predictions are depicted as two lines that represent the upper and the lower bounds for predictions when the two different intercepts are used in the empirical weathering formulation as described above. As would be expected, the lines for the two different release rules cross each other during the melt. The 40/20 rule is lower at the beginning of the season for alkalinity because of the initial acid pulse, but ends higher than the uniform rule because the solutes already have been eluted, thus leaving less acidic water than the uniform elution case. Similarly, the 40/20 rule exhibits larger changes in concentration during rain events than the uniform release rules rule.

Considering the model predictions for the inflows during the calibration year 1986, the general pattern of dilution and recovery is reproduced for all solutes (fig. 8). Both silica and SBC are overpredicted during the latter part of the melt period by approximately 10 μ eq L⁻¹; this overprediction could reflect biotic influences which are not included in the model such as diatomaceous uptake of silica and biotic uptake of calcium, magnesium and potassium during this period. Similarly, model predictions for the lake outflow (fig. 9) show the same general patterns, although SBC and silica are systematically underpredicted by 5 to 10 μ eq L⁻¹ during the first three months of the simulation. This underprediction may result from errors in the distributed melt formulation or may indicate an additional source for these solutes such as ground water.

During the validation year 1987, however, the observed patterns of solute concentrations are not so well reproduced. The minimum alkalinity observed during snowmelt is correctly predicted, but the pattern of dilution during the initial phases of snowmelt is not captured correctly. In the inflows for the uniform release case (which better predicts alkalinity), silica is slightly overpredicted and SBC is substantially underpredicted. In addition, alkalinity is underpredicted during the early part of the snowmelt. These observations taken together suggest that cation exchange, not weathering, is supplying SBC and alkalinity (through consumption of hydrogen ion) during this first phase of melt. Cation exchange was not obvious during 1986 owing to the large volume of water that flowed through the system and to less intensive sampling of surface waters during the beginning of snowmelt. This additional alkalinity-generating mechanism would further ameliorate the acid pulse at the onset of snowmelt. Current model predictions overestimate the loss of alkalinity, and hence, provide a conservative prediction.

Deposition Scenarios

To assess the sensitivity of the watershed to different loadings of acidic solutes, two different atmospheric deposition scenarios were considered: one in which both sulfuric acid and ammonium nitrate loadings were doubled, and one in which both of these loadings were halved. Factor analysis indicates that the cations associated with sulfate and nitrate are hydrogen and ammonium, respectively (Lowell Ashbaugh, California Air Resources Board, written commun., 1988). Owing to the nitrogen cycle representation in ALF in which ammonium is exchanged for hydrogen ion, the scenarios are equivalent to changing the nitric acid loading to the system for this model. The results of these scenarios for the lake outlet are shown in figure 10. Again, each case is represented with a pair of lines bounding the predictions. The solid line is the half-loading case and the stippled line is the double-loading case. In figure 10, 1986 is indicative of a heavy snow year and 1987 represents a light snow year. The alkalinity of the system is not very sensitive to changes in loadings. The initial acid pulse causes the alkalinity in the outflow to decrease from about 20 μ eq L⁻¹ under current loadings (uniform release case in figure 9) to about 15 μ eq L⁻¹. The acid pulse is more pronounced under the 40/20 release, as expected, with a minimum observed alkalinity of 8 μ eq L⁻¹ for 1986 conditions and 12 μ eq L⁻¹ for 1987 conditions as compared with minimum predicted alkalinities under current deposition of 10 and 15 μ eq L⁻¹ for 1986 and 1987, respectively. The snowpack loadings required to exhaust the alkalinity of the lake epilimnion, expressed in multiples of the current snow loading for 1986 and 1987, are summarized in table 4. No rain-on-snow events are assumed to occur.

and sulfuric acid required to exhaust alkalinity of Emerald Lake						
Solute release rule	Heavy snow year (1986)	Light snow year (1987)				
Uniform release	4 - 7	11 - 18				
40/20 release	4 - 6	7 - 11				
80/20 release	2 - 4	2 - 5				

	Table	4.	Multiples	of	current	deposition	loadings	of	ammonium	nitra	te
--	-------	----	-----------	----	---------	------------	----------	----	----------	-------	----



FIGURE 10.—The effect of halving and doubling current nitrate and sulfate deposition on alkalinity of the lake outflow.

23

SUMMARY

This simple process-oriented model is able to capture the basic solute patterns during snowmelt in an alpine catchment where ground water is a minor contributor to streamflow. It includes an empirical representation of primary mineral weathering as the only alkalinity-generating mechanism. The model seems to be adequate during a heavy snow year, such as the one that served as the calibration year for this model. During a light snow year, there is evidence that cation exchange may be a substantial source of alkalinity during the initial phases of snowmelt. By not including cation exchange, the model overestimates the acidification during this period. The minimum observed alkalinity, which occurs through dilution, is predicted correctly by the model. The representation of the lake as simply a mixing volume with no additional chemical reactions is supported by the observations.

Model results indicate a change of 2 to 5 μ eq L⁻¹ in the minimum alkalinity if the deposition of both ammonium nitrate and sulfuric acid is doubled. This estimate represents an upper bound on the alkalinity change because cation exchange is ignored. The predicted change in alkalinity would be insufficient to exhaust the alkalinity of the lake. Deposition would have to increase between two and 18 times the current loadings to accomplish that; the precise increment depends on hydrologic conditions and on the pattern of solute release from the snowpack. Although these increases in deposition seem to be very large, one should remember that current loadings to the system are low. The most likely scenario for acidification of Emerald Lake under conditions ranging from current to double the present-day deposition is an acidic rainstorm occurring during the latter part of snowmelt when the lake is at its minimum alkalinity owing to dilution from the meltwater. Such an event was observed in summer 1984. The infrequency of rainstorms in the Sierra Nevada, however, makes intensive sampling of storms difficult.

CONCLUSIONS

From this analysis of the data collected at Emerald Lake, additional avenues of research can be suggested both for modeling and field investigation. The simple form of ALF and its modest data requirements should allow for its widespread application in watersheds that are less intensively studied than Emerald Lake. The model can be calibrated with only water-quality and quantity measurements from the outflow of the lake. To run the model for any particular year requires an estimate of solute loadings in the pack. If there are no internal sources or sinks of sulfate and chloride in the watershed, these loadings can be estimated from the mass export from the watershed. Application of ALF to other watersheds in the Sierra Nevada would permit a regional assessment of the risk of episodic acidification.

The pattern of solute release from the snowpack is not consistent across the "nonreactive" solutes. In particular, sulfate, typically the mobile anion in such watersheds, has a surprisingly constant concentration in streamwater given the small amount of soil in the watershed. At a high altitude catchment in Scotland, sulfate concentration varied between 20 and 80 µeq/L during spring and summer, a factor of 4 (Cooper and others, 1987). Point measurements at Emerald Lake (Williams and Melack, 1988) and laboratory studies suggest that an acidic pulse should come from the pack at the beginning of snowmelt. However, stream concentrations of chloride, the least reactive solute, indicate a less intense ionic pulse than point measurements. Additional field observations and experiments need to be conducted to determine whether there are regulating mechanisms in the soil which control the sulfate concentration or whether these packs, due to the steep slope, ice lenses, mixing of waters at different stages of solute elution, or some other factor, do not in effect produce an acid pulse on the catchment scale. If the pack effectively elutes its solutes uniformly at this scale, an additional source for the observed nitrate and chloride peak must be identified. The apparent attenuation of the acid pulse, whatever its cause, renders this watershed much less sensitive to acidification from an acidified snowpack than if such a pulse existed.

The input-output budgets for sulfate and chloride were inconsistent between the two years of intensive study at Emerald Lake. The chloride input for 1986 is suspect due to sample contamination, but the sulfate budget deserves further study because the differences are too large to be attributable solely to sampling or analytical error. The most important distinction between the two years may be the size of the snowpack. The large amount of water in 1986 may have been able to leach more sulfate out of the watershed than in 1987, resulting in a net export in 1986 but a net retention in 1987. If this explanation is true, the chemical dynamics at Emerald Lake are more complicated than first thought both because sulfate is a reactive ion and because the flowpath through the watershed would determine the streamwater chemistry.

Paired mini-catchments having an area of several hundred square meters need to be identified at Emerald Lake to investigate sulfate regulation mechanisms. These catchments would represent the extremes in soil interaction; that is, one should have only bare bedrock and the other should be predominantly covered with soil. Meltwater draining the bare rock catchment would be affected by soil or talus, and thus, the existence or absence of an acidic pulse from the snowpack could be ascertained. In addition to runoff, soil solutions prior to and during the initial phases of melt need to be monitored to determine the effects of the soil on the "nonreactive" solutes.

Under current depositional loads and even with double the current loads, the Emerald Lake watershed is more susceptible to acidification from an acidic rainstorm which falls during the late melt period when the lake is its most dilute than to acidification from a acidic snowpack. The infrequency of such events in the Sierra Nevada makes sampling difficult. One field effort that is feasible and that may shed some light on the watershed's response to acidic pulses is intensive sampling over the daily melt cycle. Sampling during early, mid and late melt season would provide information on the watershed's chemical response to more acidic pulses, to clean pulses, and to clean pulses after the system has been strongly leached. In addition if sampling were carried out at points above and below the inflow meadow, then the influence of vegetation could be assessed.

REFERENCES CITED

- Brimblecombe, P., Clegg, S.L., Davies, T.D., Shooter, D., and Tranter, M., 1987, Observations of the preferential loss of major ions from melting snow and laboratory ice: Water Research, v. 21, p. 1279-1286.
- Bytnerowicz, Andrew and Olszyk, D.M., 1988, Measurement of atmospheric dry deposition at Emerald Lake in Sequoia National Park: Sacramento, Ca., California Air Resources Board, Final Report, Contract A7-32-039, 48 p.
- California Air Resources Board, 1986, The fourth annual report to the governor and the legislature on the Air Resources Board's acid deposition research and monitoring program: Sacramento, CA, 374 p.
- Chen, C.W., Gherini, S.A., Hudson, R.J.M., and Dean, J.D., 1983, The integrated lakewatershed acidification study: Model principles and application procedures: Lafayette, Ca., Tetra Tech, Inc., EPRI Report EA-3221, v. 1, RP 1109-5, 188 p.
- Christophersen, Nils, Dymbe, L.H., Johannessen, Merete, and Seip, H. M., 1984, A model for sulphate in streamwater at Storgama, southern Norway: Ecological Modelling, v. 21, p. 35-61.
- Clow, David, 1987, Geologic controls on the neutralization of acid deposition and on the chemical evolution of surface and ground waters in the Emerald Lake watershed, Sequoia National Park, California: Fresno, Ca., unpublished Masters Thesis, California State University, 216 p.
- Cooper, D.M., Morris, E.M. and Smith, C.J., 1987, Precipitation and streamwater chemistry in a subarctic Scottish catchment: Journal of Hydrology, v. 93, p. 221-240.
- Dozier, Jeff, 1988, Topographic distribution of solar radiation over an alpine, snow-covered drainage basin [abs]: Eos Transactions of American Geophysical Union, v. 69, p. 1198.
- Dozier, Jeff, Melack, J.M., Marks, Daniel, Elder, Kelly, Kattelmann, Richard, and Williams, Mark, 1989, Snow deposition, melt, runoff, and chemistry in a Sierra Nevada watershed: Sacramento, Ca., California Air Resources Board, Final Report, Contract A6-147-32, 268 p.
- Dracup, J.A., Melack, J.M., Marks, Daniel, Kattelmann, Richard, and Sickman, J. O., 1989, The hydrologic mass balance of the Emerald Lake basin, Sequoia National Park, California: Sacramento, Ca., California Air Resources Board, Final Report, Contract A4-123-32, 160 p.
- Drever, J.I., 1988, The Geochemistry of Natural Waters (2nd ed.): Englewood Cliffs, New Jersey, Prentice-Hall, 437 p.
- Johannessen, Merete and Henriksen, Arne, 1978, Chemistry of snow meltwater—changes in concentration during melting: Water Resources Research, v. 14, p. 615-619.
- Leivested, H. and Muniz, I.P., 1976, Fish kill at low pH in a Norwegian river: Nature, v. 259, p. 391-392.
- Lund, L.J., Brown, A.D., Lueking, M.A., Nodvin, S.C., Page, A.L., Sposito, Garrison, 1987, Soil processes at Emerald Lake Watershed: Sacramento, Ca., California Air Resources Board, Final Report, Contract A3-105-32, 114 p.

- Marks, Daniel, 1988, Climate, energy exchange, and snowmelt in Emerald Lake watershed, Sierra Nevada: Santa Barbara, CA, unpublished Ph.D. thesis, Univ. of California, 158 p.
- Melack, J.M., Cooper, S.D., Jenkins, T.M., Jr., Barmuta, Leon, Hamilton, Scott, Kratz, Kimberly, Sickman, James, and Soiseth, Chadwick, 1989, Chemical and biological characteristics of Emerald Lake and the Streams in its watershed, and the responses of the lakes and streams to acidic deposition: Sacramento, Ca., California Air Resources Board, Final Report, Contract A6-184-32, 377 p.
- Miller, S.A., 1950, Some snow-melt runoff characteristics: Eos Transactions of the American Geophysical Union, v. 31, p. 741-745.
- Morel, François and Morgan, J.J., 1972, A numerical method for computing equilibria in aqueous chemical systems: Environmental Science and Technology, v. 6, p. 58-66.
- Nishida, A.I. and Schnoor, J.L., 1989, Steady state model to determine lake resources at risk to acid deposition in the Sierra Nevada, California: Sacramento, Ca., California Air Resources Board, Final Report, Contract A7-32-036, 189 p.
- Peters, N. E., 1987, Chloride cycling in forested lake watersheds in the west-central Adirondack Mountains, New York [abs]: Prague, Czechoslovakia, GEOMON, International Workshop on Geochemistry and Monitoring in Representative Basins, May, 1987: Geological Survey, Prague, p. 264-266.
- Skartveit, A. and Gjessing, K.T., 1979, Chemical budgets and chemical quality of snow and runoff during spring snowmelt, Nordic Hydrology: v. 10, p. 141-154.
- Sorooshian, Soroosh, Bales, R.C., Gupta, V.K., Noppe, P.A., and Wolford, R.A., 1989, Development of watershed models for Emerald Lake watershed in Sequoia National Park and for other lakes of the Sierra Nevada: Sacramento, Ca., California Air Resources Board, Final Report, Contract A732-035, 64 p.
- Stephenson, N.L., 1988, Climatic control of vegetation distribution—The role of the water balance with examples from North American and Sequoia National Parks: Ithaca, New York, unpublished Ph.D. dissertation, Cornell University, 295 p.
- Swift, L.W., 1976, Algorithm for solar radiation on mountain slopes:Water Resources Research, v. 12, p. 108-112.
- Weintraub, Jill, 1986, An assessment of the susceptibility of two alpine watershed to surface water acidification—Sierra Nevada, California: Bloomington, Ind., unpublished Masters Thesis, Indiana University, 186 p.
- Williams, M. W. and Melack, J.M., 1988, Snowpack runoff contributions to the major solute chemistry of lake and stream waters in a high-altitude watershed in the Sierra Nevada [abs], Eos Transactions of the American Geophysical Union, v. 69, p. 1198.
- Wyles, W.S., 1986, The buffering capacity of Sierra Nevadan soils exposed to simulated acid precipitation: Davis, Ca, Univ. of California, unpublished Masters Thesis, 188 p.

APPENDIX A.

Sample input/output from computer model

This attachment contains sample input which are necessary to run the computer model and samples of the output which the computer generates. The files are annotated by using Times-Roman typeface, the same typeface as this is written in. The contents of the files and any prompts which the computer types are written in Helvetica typeface, as these words are.

As the model runs, it executes a series of prompts for the names of the input files. Note that no output file name is requested since they are based on the name of the main input file. A sample dialog follows.

RUN DIST.86.CPL *User types the appropriate system command to execute the program

15 Mar 89 11:16:36 Wednesday ENTER NAME OF MAIN INPUT FILE: ILWAS>IN.WY86>I861/18A3.5 *User enters names of files are requested ENTER NAME OF RAIN INPUT FILE: ILWAS>IN.WY86>RAIN.WY86 ENTER NAME OF ICE ELEV INPUT FILE: ILWAS>IN.WY86>ICELEV.WY86 ENTER NAME OF PRE 10 DAYS Q INPUT FILE: ILWAS>IN.WY86>Q10DAY ENTER NAME OF OBS STREAM INPUT FILE: ILWAS>IN.WY86>IN.1 ENTER NAME OF OBS LAKE OUTPUT FILE: ILWAS>IN.WY86>OUT.CONC ENTER NAME OF OBS STREAM TEMP FILE: ILWAS>IN.WY86>IN.TEMP ENTER NAME OF OBS THERMOCLINE: ILWAS>IN.WY86>THERMO.WY86 JD= 198 *Computer indicates which day it is working on JD= 199 JD= 200 JD= 362 JD= 363 JD= 364 JD= 365 **** STOP
The required input files follow.

Main Input File С С Ĉ С *These spaces are reserved for notes about the input С *Lake initial cond: Julian day, ANC, CL, H, NH4, NO3, SBC, SIO2, SO4 С С 2 *kr flag=> kr =F(Q) formula to use:1=actual,2=upper,l=lower 0.001106 0.1 100 *partial pressure of CO2, % NO3 retained, day for Biotic influnce 196 0.00 3.9 3.5 1.8 4.1 5.1 0.00 2.6 *ice chem initial condition 196 43.0 7.7 0.81 0.4 6.8 61.47 20.0 6.5 *epilimnion chem initial condition 196 37.0 3.6 1.66 1.9 12.7 58.88 20.0 6.6 *hypolimnion chem initial condition 198 365 202 365 1 *begin, end day of simulation, begin, end snowmelt hydrograph 198 2100. *julian day and discharge 199 1800. 200 1600. 363 880. 364 1100. 365 1800. *Total ionic loading of the snow pack in Equivalents: SBC, CL, NO3, NH4, SO4 8517.0 5619.7 4970.0 4107.0 9894.9 *Number of sub-units, total area, respective areas 7 1200000. 140298.82 132776.09 141176.47 396447.62 245115.45 134907.53 9278.03 *Sub-unit #, slope, aspect, lattitude, elution (100 100=uniform), concentration factor 1 33.7 42. 36. 100 100 8. 2 30.6 27. 36. 100 100 8. 3 28.0 10. 36. 100 100 8. 4 33.8 305. 36. 100 100 8. 5 27.2 275. 36. 100 100 8. 6 33.7 253. 36. 100 100 8. 7 13.5 327. 36. 100 100 8. *Representive values of inflow before snow melt. *J-day, discharge,SBC,CL,NO3,NH4,SO4,H 198 2100.8 1.5 1.99 1.07 0.77 0.94 1.73 199 1744.5 1.5 1.99 1.07 0.77 0.94 1.73 200 1551.3 1.5 1.99 1.07 0.77 0.94 1.73 201 1732.2 1.5 1.99 1.07 0.77 0.94 1.73

Rain data input file

297 14.8741 5.3597 22.9 24.94 20.2 14.79 *julian day, SBC CL NO3 NH4 SO4 H 324 24.2725 10.437 25.8 8.869 9.369 4.467 353 10.191 2.2567 2.9 4.435 2.498 2.951

Ice elevation file (elevation of ice/epilimnion interface):

196 4.5 *julian day, elevation 197 4.53125 198 4.5625

364 10.5 365 10.5

.

10-day average flow file (assumed to cover entire water year):

2763.2 2198.7 1915.1 1606.7 2170.1 1962 1791 1616.7 1715.2 2053.2

Chemical observations from main inflow:

198 36 8.77 8.04 7.34 1.4791 0.8 61.4 33.5 *julian day, ANC, Cl, NO₃, SO₄, H, NH₄, SBC, SIO₂ 213 27.7 6.045 10.9675 7.6025 1.1977 0.325 47.5225 37.175 214 26.2 6.175 11.39 7.79 0.89149 0.15 48.69 36.35 217 26.8 4.795 9.79 6.66 1.2466 0 49.31 40.85

- ۵
- •

Chemical Observations for lake outflow:

213 5.98 1.0471E-06 0.2 32.5 5.8 10.15 8 45.65 56.4 *J-DAY,pH,H,NH4,ANC,CL,NO3,SO4,SIO2,SBC 214 5.98 1.0471E-06 0.1 35 5.4 9.9 7.2 45 53.2 218 6.04 9.1201E-07 0.1 35 6.9 10.6 6.6 43.9 56.7

- •
- .
- .

Inflow stream temperature file:

198 0.13875 *Julian day and temperature 199 0.185 200 0.23125

- •
- .

-

Thermocline elevation file:

0.0 0.0 *elevation, julian day. Code interpolates to obtain daily values.

0.0 25.0 7.6 72.0 8.5 99. 8.0 126. 2.5 191. 2.5 230. 0. 275. 0. 302. 5.5 317. 3.5 337. 1.0 351. 0. 357. 0. 365.

<u>Output files</u> There are fourteen output files produced by the program. All have a leading asterisk to allow for importation to a graphics package. Fields are tab delimited. The following is a list of these files with an explanation of what each contains.

D1861/18A3.5	sub-unit #1
D2861/18A3.5	sub-unit #2
D3861/18A3.5	sub-unit #3
D3861/18A3.5	sub-unit #4
D5861/18A3.5	sub-unit #5
D6861/18A3.5	sub-unit #6
D7861/18A3.5	sub unit #7
I861/18A3.5	snowmelt, calc inflow, observed inflow, for constructing bar
	graphs
L861/18A3.5	lake output
RL861/18A3.5	calculated vs observed, lake outflow
RS861/18A3.5	calculated vs observed, inflow
S861/18A3.5	combined stream inflow (d1 through d7)
V861/18A3.5	a text file for printing various daily information for visual
	review
VOL861/18A3.5	jday, total, ice, epilimnion, and hypolinion volumes

D1861/18A3.5

JDAY 202 203 204 205 206 207 208	Q 368.094 716.812 908.620 709.459 700.403 903.974 948.512	ANCST 34.467 33.910 33.140 32.575 32.039 31.345 30.626	CLST 2.439 2.439 2.439 2.439 2.439 2.439 2.439 2.439	DOCST 0.000 0.000 0.000 0.000 0.000 0.000 0.000	HST 0.598 0.607 0.621 0.631 0.641 0.655 0.669	NH4ST 0.000 0.000 0.000 0.000 0.000 0.000 0.000	NO3ST 1.941 1.941 1.941 1.941 1.941 1.941 1.941 1.941	SBCST 43.141 42.584 41.814 41.249 40.713 40.018 39.300	SIO2ST 39.444 38.887 38.117 37.552 37.016 36.322 35.604	SO4ST 4.295 4.295 4.295 4.295 4.295 4.295 4.295	+/-BAL1 0.000 0.000 0.000 0.000 0.000 0.000 0.000	PH 6.223 6.217 6.207 6.200 6.193 6.184 6.174
208	948.512	30.626	2.439	0.000	0.669	0.000	1.941	39.300	35.604	4.295	0.000	6.174

1 hearth 1

<u>1861/18A3.5</u>:

IONS	J DAY	SBC	ANC	CL	DOC	H	NO3	SO4	NH4
CATIONSINPUT(S)	198	1.50	0.00	0.00	0.00	1.74	0.00	0.00	0.77
ANIONSINPUT(S)	198	0.00	0.00	1.99	-0.01	0.00	1.08	0.94	0.00
CATIONSCALC	198	40.78	0.00	0.00	0.00	0.56	0.00	0.00	0.00
ANIONSCALC	198	0.00	36.89	1.99	-0.01	0.00	0.97	0.94	0.00
CATIONSOBS	198 198	61.40 0.00	0.00 36.00	0.00 8.77	0.00 -3.53	1.48 0.00	0.00 8.04	0.00 7.34	0.80
CATIONSINPUT(S)	199	1.50	0.00	0.00	0.00	1.74	0.00	0.00	0.77
ANIONSINPUT(S)	199	0.00	0.00	1.99	-0.01	0.00	1.08	0.94	0.00
CATIONSCALC	199	40.82	0.00	0.00	0.00	0.56	0.00	0.00	0.00
ANIONSCALC	199	0.00	36.93	1.99	-0.01	0.00	0.97	0.94	0.00

<u>L861/18A3.5</u>:

*											
JDAY	′Q	ANC LK	CL LK	DOCLK	HLK	NH4 LK	NO3 LK	SBC LK	SIO2 LK	SO4 LK	PH LK
198	2100.8	41.9	7.3	-1.2	0.8	0.4	6.4	60.0	21.4	6.1	6.1
199	1744.5	40.8	6.9	-1.1	0.8	0.4	6.0	57.8	22.0	5.7	6.1
200	1551.3	39.7	6.6	-1.1	0.9	0.4	5.7	55.9	22.5	5.4	6.1
201	1732.2	38.8	6.3	-1.0	0.9	0.4	5.4	54.1	23.0	5.1	6.0
202	3500.0	37.7	5.8	-0.9	0.9	0.4	5.0	52.1	24.3	5.0	6.0
203	6800.0	36.5	5.2	-0.7	0.9	0.3	4.5	49.6	26.6	4.8	6.0
204	8600.0	35.3	4.6	-0.5	0.9	0.3	3.9	47.3	28.8	4.7	6.1
205	6700.0	34.3	4.2	-0.4	0.9	0.3	3.6	45.6	29.9	4.6	6.1
206	6600.0	33.5	3.9	-0.4	0.9	0.2	3.3	44.2	30.6	4.5	6.1
207	8500.0	32.7	3.6	-0.3	0.9	0.2	3.0	42.8	31.4	4.4	6.1
208	8900.0	31.9	3.3	-0.2	0.9	0.2	2.8	41.5	31.9	4.4	6.1

APPENDIX B.

Model Listing

The following is a listing of the program code. It has been written in standard FORTRAN-77. The only system-specific statements are the \$INSERT statements, which are called \$INCLUDE in some operating systems.

```
С
       MODIFIED FOR RAIN ON SNOWMELT.
С
С
       ADDED CALLS TO SUBROUTINES AFTER 150 CONTINUE
С
С
С
С
С
С
С
C.... MAIN.F77
$INSERT COMMON/MAIN/
      COMMON/INIT/ RELO
      REAL*8 CN(9), SPCT(9), CROS(9)
      REAL*8 RS, RA, RL
      REAL*8 F1, F2, F3, F4, F5, F6, F7, F8, F9
      REAL*8 RELO
      REAL*8 QDIFF
      REAL*8 ANCAM, CLAM, DOCAM, HAM, NH4AM
      REAL*8 NO3AM, SBCAM, SIO2AM, SO4AM
      INTEGER DATE
      CHARACTER FNAME*30, HEAD*80
      EQUIVALENCE (ANC, CN(1))
      EQUIVALENCE (ANCT, SPCT(1))
      TAB=CHAR(137)
С
C... INITIAL VALUES
С
       TO = REF. TEMP
С
         KHO = HENRYS K FOR CO2(q) AT TO
С
         R = UNIVERSAL GAS CONST
С
         RELO = INIT VALUE FOR % REL IONS IN FUNCTION RELFUNC
С
      IRAIN1=0
      RELO =0.D0
      R = 1.987
      KHO=10.**(-1.47)
      TO = 25.+273.16
      INIT=0
      IO=1
      XAVE=0.0
      DO 21 IA=1,10
        QSUM(IA) = 0.0
        CMELT(IA) = 0.0
        CMELTO(IA)=0.0
        CREL(IA) = 0.0
        CRELO(IA) = 0.0
        IOFF(IA) = 1
        IOFF2(IA)=1
        IOFF3(IA) = 1
        IOFF4(IA) = 1
21
     CONTINUE
C...Integer Lake/Stream Observed Day (ILOD, ISOD) COUNTERS USED WRITING
C TO A FILE WHICH CONTAINS OBSERVED DATA VS. CALCUATED DATA (R^2)
```

С

```
ILOD=1
      ISOD=1
C...IR IS A DAILY COUNTER FOR RAIN DAY CONCENTRATIONS
      IR=1
С
C... CALL FILER TO OPEN IN/OUTPUT FILES, READ INITIAL CONDITIONS
      CALL FILER
C... CALL LAKE AND GIVE IT THESE CONDITIONS!
      CALL LAKE(1)
C... CALL DISTM AND DISTRIBUT THE DISCHARGE AND LONS, CALC RELEASE RULE
      CALL DISTM
С
C... BEGIN DAILY TIME STEPS
      IPC=1
      NPTS=JDN-JD1+1
      DO 50 I=JD1, JDN, 1
       NRR=0
       XDAT(IPC)=FLOAT(I)
        JDAY=I
        WRITE(*,*)' JD=', JDAY
        WRITE(*,*)QSU(4,JDAY)
С
C...READ DAILY INFLOW TEMP IN CELCIUS
        READ(20, *) JY, ST
C
C...CONVERT INFLOW TEMP TO KELVINS
       ST=ST+273.16
С
C...MEMORY, FOR Q vs. Kr
      IF (MEM.EQ.1) THEN
       QT=0.0
       DO 52 J=0,9
52
          QT=QT+SH(I-J)
       QT=QT/10.
      ELSE
        QT=SH(I)
      ENDIF
С
C... ARE WE BEFORE, DURING, OR AFTER THE MELT.
      IF (JDAY.LT.JBM) THEN
      GOTO 200
      ELSEIF (JDAY.GT.JEM) THEN
      GOTO 300
      ENDIF
С
С
C*******
C... HERE WE DO DISTRIBUTED MELT, JDAY .GE. JBM .AND. JDAY .LT. JEM
C... THE Q FOR EACH SUB UNIT IS IN ARRAY AQSU(IA,ND) WHERE IA IS THE
C... SUB UNIT AND ND IS THE DAY. ANOTHER ARRAY, ND(IA) HAS THE DAY
C... THAT EACH SUB UNIT DRIES UP
С
C... FIRST WE CHECK FOR RAIN ON SNOW.
С
       IF(RR(I).GT.0.0)THEN
         SBC=SBCR(IR-1)
         CL=CLR(IR-1)
         NO3=NO3R(IR-1)
         NH4 = NH4R(IR-1)
         SO4 = SO4R(IR-1)
         H = HR(IR - 1)
         DOC=DOCR(IR-1)
         IF (DOC.LT.0.0) THEN
             DOC = 0.0
```

۰.

```
WRITE (*, *) ' WARNING: NEGATIVE DOC ON RAIN JDAY =', JDAY
         ENDIF
         WRITE(*,*)'CALL NRWEATH 1'
С
         CALL NRWEATH
         DO 152 IJK=1,9
           CROS(IJK) =CN(IJK)
152
         CONTINUE
       ENDIF
С
C... CALCULATE PRECENTAGE OF IONS RELEASED FROM EACH SUB UNIT
C... CONVERT EQ/M**3 TO uEQ/L, WHERE QSU IS IN METERS CUBED
      DO 150 IA=1,NSU
         IF (JDAY.GT.ND(IA,1)) GOTO 150
         DO 151 IC=1,9
            IF (IC.EQ.1.OR.IC.EQ.8) THEN
              CN(IC) = 0.D0
              GOTO 151
            ENDIF
            T=TIME (JDAY, IA, IC)
            IF (JDAY.GT.TSTAR (JDAY, IA, IC) ) THEN
C.. UNITS uEQ/L
               IF (IC.EQ.2.AND.IA.EQ.4) WRITE (*,*) 'AFTER T*'
С
              CN(IC) = AB(JDAY, IA, IC, 3) * EXP(-AB(JDAY, IA, IC, 4) * T) * 1000.
            ELSE
               IF (IC.EQ.2.AND.IA.EQ.4) WRITE (*,*) 'BEFORE T*'
С
              CN(IC) = AB(JDAY, IA, IC, 1) * EXP(-AB(JDAY, IA, IC, 2) * T) * 1000.
            ENDIF
             IF (IC.EQ.9.AND.IA.EQ.4) WRITE (*, *) AB (JDAY, IA, IC, 1),
С
С
      &
                                                  AB(JDAY, IA, IC, 2),
С
      &
                                                  AB(JDAY, IA, IC, 3),
С
      &
                                                  AB(JDAY, IA, IC, 4),
С
      &
                                                  TSTAR (JDAY, IA, IC)
151
         CONTINUE
            IF (IA.EQ.4) THEN
С
         PRINT OUT MELT WATER, BEFORE ROCK REACTION
       WRITE (*, 33) JDAY, TAB, ANC, TAB, CL, TAB, DOC, TAB, H,
С
      &TAB,NH4,TAB,NO3,TAB,SBC,TAB,SIO2,TAB,SO4,TAB,CB,TAB,PH
С
33
      FORMAT(I3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
     & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
     & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
     & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
     & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3)
            ENDIF
C... CALC HYDROGEN ST THE ELUDED IONS HAVE CHARGE BALANCE
           H=CL+NO3+SO4-SBC-NH4
С
C... SEND VALUES IN uEQ/L
С
           DOC=FD(H,SBC,NH4,SO4,CL,NO3)
          IF (DOC.LT.0.0) THEN
            WRITE(*,*)' '
            WRITE (*, *) ' WARNING: NEGATIVE DOC ON JDAY =', JDAY
            WRITE (*, *) ' DOC = ', DOC
            DOC=0.0
          ENDIF
          IF (ANC.LT..01) ANC=0.D0
          IF (SIO2.LT..01) SIO2=0.D0
С
C... THESE VALUES ARE FOR AFTER MELT (AM) WHEN INFLOW CONC TO LAKE IS CONST.
С
     THESE VALUES ARE FROM THE LAST CONTRIBUTING SUBUNIT, IF TWO SUBUNITS
     DRY UP ON THE SAME DAY, THE THE ONE INPUTED LAST IS USED.
С
С
      ANCAM=ANC
      CLAM=CL
      DOCAM=DOC
      НАМ≕Н
```

```
35
```

```
NH4AM=NH4
      NO3AM=NO3
      SBCAM=SBC
      SI02AM-SI02
      SO4AM=SO4
С
С
         WRITE(*,*)' '
С
         WRITE(*,*)'IA --',IA
         WRITE(*,*) 'ANC=', ANC
С
         WRITE(*,*)'CL=',CL
С
С
         WRITE(*,*)'DOC=',DOC
С
         WRITE(*,*)'H=',H
С
         WRITE(*,*)'NH4=',NH4
С
         WRITE(*,*)'NO3=',NO3
С
         WRITE(*,*)'SBC=',SBC
С
         WRITE(*,*)'SIO2=',SIO2
С
         WRITE(*,*)'SO4=',SO4
С
          IF(IA.EQ.4)THEN
С
             WRITE(*,*)' '.
С
             WRITE(*,*)'CL=',CL
С
          END IF
        CALL NRWEATH
С
          IF(IA.EQ.4)THEN
С
            WRITE(*,*)'CL=',CL
С
             WRITE(*,*)' '
С
          ENDIF
С
         WRITE(*,*)'AFTER CALL TO NRWEATH'
С
         WRITE(*,*)'ANC=',ANC
С
         WRITE(*,*)'CL=',CL
С
         WRITE(*,*)'DOC=',DOC
С
         WRITE(*,*)'H=',H
         WRITE(*,*)'NH4=',NH4
С
         WRITE(*,*)'NO3=',NO3
С
С
         WRITE(*,*)'SBC=',SBC
         WRITE(*,*)'SIO2=',SIO2
С
С
         WRITE(*,*)'SO4=',SO4
С
         WRITE(*,*)' '
С
C... HERE WE TRACE THE OUTFLOW OF EACH SUB UNIT, CHECK FOR RAIN FIRST
        IF (RR (JDAY).GT.0.0) THEN
          XCENT=AREA(IA)/AREAT
          DO 169 IC=1,9
             CN(IC) = ((CN(IC) *QSU(IA, JDAY)) +
                    (CROS(IC)*XCENT*RR(JDAY)))/
     &
                    (QSU(IA, JDAY) +RR(JDAY) *XCENT)
     &
169
          CONTINUE
          CALL NROPEN
        ENDIF
        PH=-LOG10 (H/100000)
        WRITE (28+1A, 43) JDAY, TAB, QSU(IA, JDAY), TAB, ANC, TAB, CL, TAB, DOC,
     &TAB, H, TAB, NH4, TAB, NO3, TAB, SBC, TAB, STO2, TAB, SO4, TAB, CB, TAB, PH
С
        DO 170 IC=1,9
          SPCA(IA, LC) = CN(IC)
170
        CONTINUE
150
      CONTINUE
С
C... MIX ALL SUB UNIT OUTPUT BEFORE MIXING INTO LAKE
      DO 180 IC-1,9
        CN(IC)=0.0
        QCSUM=0.0
        DO 185 TA=1,NSU
```

```
IF (JDAY.GT.ND(IA, 1)) THEN
             IF (RR (JDAY) .GT.0.0) THEN
              XCENT=AREA(IA)/AREAT
              CN(IC) = (CROS(IC) * RR(JDAY) * XCENT+QCSUM*CN(IC))
     &
                      / (QCSUM+RR (JDAY) *XCENT)
              QCSUM=QCSUM+RR(JDAY) *XCENT
             ENDIF
         ELSE
687
           CN(IC) = (SPCA(IA, IC) *QSU(IA, JDAY) +QCSUM*CN(IC))
                   /(QCSUM+QSU(IA, JDAY))
     &
           QCSUM=QCSUM+QSU(IA, JDAY)
          ENDIF
185
        CONTINUE
180
      CONTINUE
С
C... TEST FOR LOW ANC AND PRINT OUT PH
C... UNITS ARE MICRO EQUIVALENTS
С
                            123456789
      PH=-LOG10(H*1000000.0)
       IF (ABS (ANC-H) .LT.0.5) THEN
С
С
          WRITE(*,*)'
                        1
          WRITE(*,*)' ANC = H'
С
          WRITE(*,*)' JDAY =',JDAY
С
          WRITE(*,*)' ANC =', ANC
С
          WRITE(*,*)' H =',H
С
С
          WRITE(*,*)' PH =',PH
       ELSEIF (ANC.LT.H) THEN
С
         WRITE(*,*)' '
С
          WRITE (*, *) ' ANC < H'
С
С
          WRITE(*,*) ' JDAY =', JDAY
          WRITE(*,*)' ANC =', ANC
С
          WRITE(*,*)' H =',H
С
С
          WRITE(*,*)' PH =',PH
С
       ENDIF
С
C... WRITE COMBINED STREAMS OUTPUT TO FILE 13
С
      PH = -LOG10(H/1000000)
      WRITE (13, 43) JDAY, TAB, QI (JDAY), TAB, ANC, TAB, CL, TAB, DOC, TAB, H,
     &TAB, NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, CB, TAB, PH
С
C... IF STREAM OBSERVATIONS EXIST FOR THIS DAY, WRITE OUT CALC AND OBS DATA
С
    FOR COMPARISON (R^2, TAB DELIMITED, OUTPUT FILE)
С
       IF (JDAY.EQ.OSJD(ISOD)) THEN
          OSH (ISOD) = OSH (ISOD) / 1000000
          OSPH=-LOG10 (OSH(ISOD))
          OSH(ISOD) = OSH(ISOD) *1000000
          WRITE(7,43) JDAY, TAB, QI(JDAY), TAB, ANC, TAB, CL, TAB, H
     &
          , TAB, NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, PH
     &
          , TAB, OSANC (ISOD), TAB, OSCL (ISOD), TAB, OSH (ISOD), TAB,
     8
          OSNH4 (ISOD), TAB, OSNO3 (ISOD), TAB, OSSBC (ISOD), TAB, OSIO2 (ISOD)
          , TAB, OSSO4 (ISOD), TAB, OSPH
      8
          ISOD=ISOD+1
       ENDIF
4
      FORMAT(2X, 'ANC', 4X, 'CL', 3X, 'DOC', 4X, 'H', 4X,
     &'NH4', 3X, 'NO3', 3X, 'SBC', 2X, 'SIO2', 3X, 'SO4')
43
      FORMAT(I3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
      & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
      & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
      & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
      & F10.3,A1,F10.3,A1,F10.3,A1,F10.3,A1,F10.3)
3
      FORMAT(13(F5.1,1X),/)
```

```
С
```

```
C... BEGIN LAKE MIXING
      GOTO 500
С
C... HERE WE TAKE CONCENTRATION FROM INPUT PREVIOUS TO MELT
С
200 CONTINUE
C...SNOW MELT DATA: jday,Q(M**3),SBC(uEQ/L),CL,NO3,NH4,SO4,H
        READ(9,*)DAY,QI(I),SBC,CL,NO3,NH4,SO4,H
         WRITE(*,*)DAY,QI(I),SBC,CL,NO3,NH4,SO4,H
С
        DOC=FD(H,SBC,NH4,SO4,CL,NO3)
С
         WRITE(*,*)'CALL RAIN 1'
       CALL RAIN
С
         WRITE(*,*)'CALL NRWEATH 3'
       CALL NRWEATH
      PH=-LOG10(H/100000)
      WRITE(13,43) JDAY, TAB, QI(JDAY), TAB, ANC, TAB, CL, TAB, DOC, TAB, H,
     &TAB, NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, CB, TAB, PH
С
C...IF STREAM OBSERVATIONS EXIST FOR THIS DAY, WRITE OUT CALC AND OBS DATA
    FOR COMPARISON (R^2, TAB DELIMITED, OUTPUT FILE)
С
С
      IF (JDAY.EO.OSJD(ISOD)) THEN
          OSH(ISOD) = OSH(ISOD) / 1000000
          OSPH=-LOG10(OSH(ISOD))
          OSH(ISOD) = OSH(ISOD) * 1000000
         WRITE (7,43) JDAY, TAB, QI (JDAY), TAB, ANC, TAB, CL, TAB, H
          , TAB, NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, PH
     &
          , TAB, OSANC(ISOD), TAB, OSCL(ISOD), TAB, OSH(ISOD), TAB,
     &
         OSNH4(1SOD), TAB, OSNO3(ISOD), TAB, OSSBC(ISOD), TAB, OSIO2(ISOD)
     &
         , TAB, OSSO4 (ISOD), TAB, OSPH
     æ
         ISOD≕ISOD+1
      ENDIF
      GOTO 500
     CONTINUE
300
С
C... THESE VALUES ARE THE LAST VALUES OF SNOW MELT
C... AND ARE USED TO DRIVE THE MODEL WHEN ALL SUB UNITS
C... HAVE DRIED UP.
      ANC=ANCAM
      CL=CLAM
      DOC=DOCAM
      H=HAM
      NH4=NH4AM
      NO3=NO3AM
      SBC=SBCAM
      SIO2=SIO2AM
      SO4=SO4AM
400
      CONTINUE
С
С
C... EQUILIBRATE USING NEWTON-RAPHSON METHOD
         WRITE(*,*)'CALL RAIN 2'
С
        CALL RAIN
С
         WRITE(*,*)'CALL NRWEATH 4'
        CALL NRWEATH
С
      PH=-LOG10(H/1000000)
      WRITE (13, 43) JDAY, TAB, Q1 (JDAY), TAB, ANC, TAB, CL, TAB, DOC, TAB, H,
     &TAB, NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, CB, TAB, PH
С
C...IF STREAM OBSERVATIONS EXIST FOR THIS DAY, WRITE OUT CALC AND OBS DATA
```

```
C FOR COMPARISON (R^2, TAB DELIMITED, OUTPUT FILE)
```

```
С
      IF (JDAY.EQ.OSJD(ISOD)) THEN
         OSH(ISOD) = OSH(ISOD) / 1000000
         OSPH=-LOG10(OSH(ISOD))
          OSH(ISOD) = OSH(ISOD) * 1000000
          WRITE(7,43) JDAY, TAB, QI(JDAY), TAB, ANC, TAB, CL, TAB, H
         , TAB, NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, PH
     &
     &
          , TAB, OSANC (ISOD), TAB, OSCL (ISOD), TAB, OSH (ISOD), TAB,
     &
         OSNH4 (ISOD), TAB, OSNO3 (ISOD), TAB, OSSBC (ISOD), TAB, OSIO2 (ISOD)
         , TAB, OSSO4 (ISOD), TAB, OSPH
     &
          ISOD=ISOD+1
      ENDIF
С
C... HERE WE JUMP BACK IN AFTER THE DISTRIBUTED MELT, MELT ENDED
C... AND BEFORE THE BEGINNING OF MELT
500
      CONTINUE
С
C... OUTPUT IONIC STRENGTHS OF INPUT (SNOW MELT OR RAIN),
C... CALCULATED LAKE INFLOW, AND OBS STREAM INFLOW.
С
С
         WRITE(*,*)' CALL IONIC 1'
        CALL IONIC
С
C... MIX WITH LAKE VOLUME
С
         WRITE(*,*)' CALL LAKE 1'
        CALL LAKE(0)
С
C... PRINT OUTPUT
        DO 40 J=1,9
40
             CN(J)=QI(JDAY)*CN(J)*0.001
       WRITE(15,*)' OUTFLOW LOADING,
С
                                        EQ'
С
       WRITE(15,4)
С
       WRITE (15,7) (CN(IJI), IJI=1,9)
7
      FORMAT(13(F8.1,1X),/)
88
      FORMAT(13(F8.2,A2))
        IPC=IPC+1
      IRAIN1=0
50
      CONTINUE
С
С
С
       CLOSE(6)
      CLOSE(18)
      CLOSE(17)
      CLOSE(8)
      CLOSE(9)
      CLOSE(10)
      CLOSE(11)
      CLOSE(12)
      CLOSE(13)
      CLOSE(14)
С
       CLOSE(15)
      CLOSE(5)
      CLOSE(7)
      CLOSE(19)
      CLOSE(20)
C*** USED TO CHECK LAKE MIXING
      CLOSE (29)
      CLOSE(30)
      CLOSE(31)
      CLOSE (32)
      CLOSE (33)
      CLOSE(34)
```

```
CLOSE(35)
      CLOSE (36)
      STOP
      END
      DOUBLE PRECISION FUNCTION RELFUNC (XM, IA, MS)
      REAL*8 XM
С
C... THE EQUATIONS
C... DEPENDING ON THE VALUE OF IA, A SUB UNITS HAS AN RELEASE EQUATION
C... THE XM IS CUMILATIVE MELT(%), AND REL IS CUMULATIVE RELEASE (%)
С
      IF (MS.EQ.11) THEN
        RELFUNC=XM
      ELSEIF (MS.EQ.82) THEN
        IF (XM.LE..2) THEN
          RELFUNC=4.DO*XM
        ELSE
          RELFUNC=0.25D0*XM+0.75D0
        ENDIF
         WRITE(*,*)'RELFUNC 82 =', RELFUNC
С
      ENDIF
      RETURN
      F:ND
      SUBROUTINE DISTM
$INSERT COMMON/MAIN/
      REAL*8 CN(9), SPCT(9)
      REAL*8 TMASS, PMASS, PWAT
      EQUIVALENCE (ANCT, SPCT(1))
С
      HERE WE DIVIDE MASS OF SNOW BY AREA, CALCULATE DAILY RADIATION
С
      DAILY DISCHARGE FROM EACH SUB UNIT, COEFFICENTS FOR THE RELEASE
С
      RULE FOR IONS, A1, A2, B1, B2, NDAY
С
С
C... DIVIDE TOTAL MASS OF SNOW AMONG SUB UNITS BASED ON
C... RELATIVE AREAS
      DO 66 IA=1,NSU
        QSUTOT (IA) = (AREA (IA) /AREAT) *QTOT
         WRITE(*,*)' DISTM QSUTOT(',IA,')=',QSUTOT(IA)
С
        DO 66 IC=1,9
           IF(IC.EQ.1.OR.IC.EQ.8)GOTO 66
              SPCAT(IA, IC) = (AREA(IA) / AREAT) * SPCT(IC)
         CONTINUE
66
С
         WRITE(*,*)' SPCAT(',IA,IC,')=', SPCAT(IA,IC)
С
C... DAILY LOOP FOR SOLAR RADITATION AND DISCHARGE
С
      DO 100 ID=JBM, JEM
       QS=SH(ID)
         WRITE(*,*)' JDAY', ID
CC
        WRITE(*,*)' MAIN Q(',ID,')=',QS
СС
С
C... CALCULATE PRCENTAGE INSOLATION ON SUB UNITS PER UNIT AREA
5
      SRAD = 0.0
      DO 110 IA=1,NSU
        IF(IOFF(IA).EQ.0)GOTO 110
         RSB=RSLOPE(IA)
          RA=RASPECT(IA)
          RL=RLAT(IA)
          CALL SOLRAD (ID, RSB, RA, RL, RAD)
          SRAD = SRAD + RAD
```

```
40
```

```
RSU(IA) = RAD
110
      CONTINUE
С
C... CALCULATE AREA INTO MELT
      DEM=0.0
      DO 115 IA=1,NSU
        IF(IOFF(IA).EQ.0)GOTO 115
         D=AREA(IA) *RSU(IA)/SRAD
         DEM=DEM+D
115
      CONTINUE
С
C... CALCULATE THE PRECENT DISCHARGE FOR EACH SUBUNIT
      DO 120 IA=1,NSU
        IF (IOFF (IA) .EQ.0) GOTO 120
        RCENT = (AREA(IA) *RSU(IA) /SRAD) /DEM
        QSU(IA, ID) =RCENT*QS
120
        CONTINUE
С
C... CHECK IF SUB UNIT CUMULATIVE DISCHARGE HAS EXCEEDED TOTAL WATER
CC
        WRITE(*,*)' '
      IRT=1
      DO 130 IA=1, NSU
CC
          WRITE(*,*)' QSUM(',IA,')=',QSUM(IA)
        IF(IOFF(IA).EQ.0)GOTO 130
        TQSUM(IA) = QSUM(IA) + QSU(IA, ID)
СС
          WRITE(*,*)' MAIN TQSUM(',IA,')=',TQSUM(IA)
СС
          WRITE(*,*)' QSUTOT',QSUTOT(IA)
        IF (TQSUM(IA).GT.QSUTOT(IA)) THEN
          IOFF(IA) = 0
          ND(IA, 1) = ID
СС
        WRITE(*,*)' ND(',IA,',1)=',ND(IA,1)
CC
        WRITE(*,*)' QSUTOT(',IA,')=',QSUTOT(IA)
CC
        WRITE(*,*)' QSUM(',IA,')=',QSUM(IA)
        WRITE(*,*)'
СС
          QSU(IA, ID) = QSUTOT(IA) -QSUM(IA)
          QS =QS - QSU(IA, ID)
          IRT = 0
СС
            WRITE(*,*)' DRIED UP NEW QSU(',IA,')=',QSU(IA,ID)
СС
        WRITE(*,*)' '
        ENDIF
        IF (IOFF (IA) .NE.O) THEN
          ND(IA,1) = ID
        ENDIF
130
     CONTINUE
C
C... RECALCUATE PRECENT INSOLATION PER SUB UNIT IF A UNIT 'DRYS UP'
      IF(IRT.EQ.0)GOTO 5
С
C... UPDATE THE SUM OF WATER LEAVING A SUB UNIT
      DO 140 IA=1, NSU
        IF(IOFF2(IA).EQ.0)GOTO 140
        QSUM(IA) =QSUM(IA) +QSU(IA, ID)
          WRITE(*,*)' MAIN QSUM(',IA,')=',QSUM(IA)
CC
        IF (IOFF(IA).EQ.0) IOFF2(IA) = 0
140
      CONTINUE
100
      CONTINUE
С
       DO 131 IA=1,NSU
C131
           WRITE(*,*)' ND(',IA,',1)=',ND(IA,1)
С
        DO 301 I=JBM, JEM
C301
         WRITE(*,*)'QSU(1)=',QSU(1,I)
С
C... CALC THE PARAMETERS FOR CONCENTRATION
С
      DO 300 IA=1,NSU
```

```
41
```

C	DO 250 IC=1,9 WRITE(\star \star) TA= TA + IC= TC
Ċ	KI=1
	JBS=JBM
45	0 CONTINUE
с.	NO AND AND SIOZ CONTRIBUTED, DOC IS CALCULATED LATER
	IF (IC.EQ.I.OR.IC.EQ.8.OR.IC.EQ.3) GOID 230
	PWAT=DBLF(MSFL(IA 2))/100
	TMASS=SPCAT(IA, IC)
	NQ=ND(IA,1)
	CFS=CF(IA)
	CALL ELUTR(QSU, JBS, NQ, IA, TMASS, PWAT, PMASS, CFS, A1, B1, A2, B2, NDAY)
	ND(IA,2)=NDAY
C	WRITE (*,43) JBS, TAB, IA, TAB, IC, IAB, AI, IAB, BI, IAB
47	FORMAT (13, A1, 11, A1, 11, 5 (A1, E12, 6E2), A1, I3)
1.	T=1.
	IF(MSEL(IA,1).EQ.MSEL(IA,2))THEN
	QSUM(IA) = 0.0
	DO 399 I=JBS, JEM
	QSUM(IA) = QSUM(IA) + QSU(IA, I)
35	9 CONTINUE ENDIE
	DO 400 I=JBS.JEM.1
	AB(I, IA, IC, 1) = A1
	AB(I,IA,IC,2)=B1
	AB(I, IA, IC, 3) = A2
	AB(I, IA, IC, 4) = B2
	IF(MSEL(IA, I) EQ.MSEL(IA, 2))THEN AD(T_TA_TC_1)-TMASS(OSHM(TA))
	AB(1, 1A, 1C, 2) = 0.0
	AB(I, IA, IC, 3) = TMASS/QSUM(IA)
	AB(I, IA, IC, 4)=0.0
	ENDIF
	IF (I.GT.ND($I\Lambda$, 2)) THEN
	CN(IC) = AB(1, IA, IC, 3) * EXP(-AB(1, IA, IC, 4) * T)
	ELSE $CN(TC) = AB(T TA TC 1) \times EXP(-AB(T TA TC 2) \times T)$
	ENDIF
	TIME(I, IA, IC) = T
	TSTAR(I, IA, IC) = ND(IA, 2)
	IF(I.EQ.ND(IA,2))T=0
	SPCAT(IA, IC) = SPCAT(IA, IC) - CN(IC) * QSU(IA, I)
C	WRITE (* 57) L. TAB. SPCAT (IA. IC), TAB. XGANG
5	FORMAT (13, A1, E12, 6E2, A1, E12, 6E2)
-	IF(I.EQ.JDR(KI)) GOTO 410
4	DO CONTINUE
	GOTO 250
4	10 IF (IC.EQ.2) THEN
C	BANG≈CLR(KI)
C	WRITE(^,) DANG FLSFIF(IC FO 4) THEN
	BANG=HR(KI)
	ELSEIF (IC.EQ.5) THEN
	BANG=NH4R(KI)
	ELSEIF (IC.EQ.6) THEN
	BANG=NO3R(KI)
	ELSEIF (IC.EQ. /) THEN
	RANGESRICK(VI)
	BANG=SO4R(KI)

. .

```
ENDIF
C... ADD RAIN STORAGE * RAIN CONC TO SNOW PACK ON THIS RAIN DAY
         BANG=RS(I) * (AREA(IA) /AREAT) *BANG/1000.
С
          WRITE(*,*)BANG
      SPCAT(IA, IC) = SPCAT(IA, IC) + BANG
         WRITE(*,*)SPCAT(IA,IC)
С
         JBS=JDR(KI)+1
С
          WRITE(*,*)'RAIN DAY =-', JDR(KI)
         KI = KI + 1
         IF (JBS.GT.ND(IA, 1)) THEN
           GOTO 250
         ELSE
           GOTO 450
         ENDIE
250
       CONTINUE
300
     CONTINUE
С
     RETURN
     END
С
     SUBROUTINE ELUTR (Q, JBM, NQ, IA, TMASS, PWAT, PMASS, CF, A1, B1, A2, B2, NDAY)
     IMPLICIT DOUBLE PRECISION (A-H, O-Z)
     DIMENSION Q(10,1000)
     REAL*8 MPR, MPR2
C This program takes an elution rule (e.g., 80% of the ions in 20% of *
C the melt), and the concentration on a given day and returns the two \;\;\star\;
C parameters for an exponentially declining concentration. A simple
C Newton's method of solution is used.
С
       TWAT=0
      DO 10 I=JBM, NQ
       TWAT=TWAT+Q(IA,I)
10
       CONTINUE
      WRITE(*,*)'TWAT=',TWAT,' JBM=',JBM,' NQ=',NQ
С
      SUMQ=0
      DO 20 I=JBM,NQ
        SUMQ=SUMQ+Q(IA,I)
        IF (SUMQ.LT.PWAT*TWAT) THEN
         GOTO 20
        ELSE
         GOTO 30
        ENDIF
20
        CONTINUE
      WRITE (*, *) 'ERROR IN CALC. OF DAYS REQD FOR FLOW.'
      STOP
C* ROUND UP THE NUMBER OF DAYS REQUIRED FOR WATER TO FLOW OUT
30
       NDAY=I
С
         WRITE(*,*)'NDAY= ',NDAY
C* ready to begin Newton's method solution; MAKE INITIAL GUESS
      B1 = .03
      CAVE=TMASS/TWAT
      MPR=TMASS*PMASS
      CO=CF*CAVE
50
      XSUM=0.0
      FBR=0.0
      J=JBM
      CK=NDAY-JBM+1
      DO 100 I=1,CK
         CN = CO*DEXP(-B1*DBLE(I))
         XSUM=XSUM + CN*Q(IA, J)
         FBR =FBR + DBLE(I)*Q(IA, J)*CO*DEXP(-B1*DBLE(I))
         J=J1
```

```
100 CONTINUE
       F=MPR-XSUM
       C=F-FBR*B1
       IF (DABS (F).GT.1.D-06) THEN
          B1=-C/FBR
          GOTO 50
       ENDIF
       A1=CF*CAVE
C******
      CO = CN
      MPR = TMASS -MPR
      B2 = .025
      JK≕J
      J=JK
150
      XSUM-0.0
      FBR=0.0
      DO 110 I=1, NQ-NDAY
         CN == Q(IA, J) *CO*DEXP(-B2*DBLE(I))
         XSUM-XSUM+ CN
         FBR =FBR + DBLE(1)*Q(IA, J)*CO*DEXP(-B2*DBLE(I))
         J = J + 1
      CONTINUE
110
       F=MPR-XSUM
       C=F-FBR*B2
       IF (DABS (F).GT.1.D-06) THEN
          B2=-C/FBR
          GOTO 150
       ENDIF
       A2=C0
      RETURN
      END
      SUBROUTINE SOLRAD (JDAY, 1, A, LO, R4)
      COMMON/RAD/ JD,D
      REAL*8 D
      REAL*8 1, L, L1, L0, A, L2, X, Y, R1, E, T
      REAL*8 T6, T7, T1, T0, T2, T3, D1, R4
      INTEGER JDAY
С
C... THIS IS FROM SWIFT 1976, TO CALCULATE POTENTIAL SOLAR RADIATION.
С
с...
                    1, LO, A, ARE CONVERTATED FROM DEGREES TO RADIANS
      I=DR(I)
      LO=DR(LO)
      A=DR(A)
С
      X= DCOS(I)*DSIN(LO)+DSIN(I)*DCOS(LO)*DCOS(A)
      L1=DASIN(X)
      X= DSIN(I)*DSIN(A)
      Y= DCOS(I)*DCOS(LO)-DSIN(I)*DSIN(LO)*DCOS(A)
      L2 = DATAN(X/Y)
С
      L1=RD(L1)
      L2=RD(L2)
                                   L1 AND L2 ARE NOW IN DEGREES
с...
С
       WRITE(*,*)'L1 =',L1
       WRITE(*,*)'L2 =',L2
С
C... CONVERT JDAY BEGINNING OCT 1, TO JULTAN DAY BEGINNING JAN 1
      IF (JDAY.LE.92) THEN
        JD=JDAY+273
      ELSEIF (JDAY.GT.92) THEN
        JD=JDAY-92
      ENDIF
С
```

```
с...
      D=FUNC1(0.4D0,23.3D0,10.D0)
      WRITE(*,*)' D (DEG)=',D
С
C... CONVERT LO FROM RADIANS TO DEGREES
      LO=RD(LO)
С
      E=FUNC1(1.0D0,0.0167D0,-3.0D0)
      R1=60.0D0*1.95/(E*E)
      WRITE(*,*)'R1=',R1
С
      T=FUNC2(L1)
      T7 = T - L2
      T6=-T-L2
      T=FUNC2(LO)
      T1 = T
      TO = -T
      IF (T7.LT.T1) THEN
        T3=T7
      ELSE
        T3=T1
      ENDIF
      IF (T6.GT.TO) THEN
        T2 = T6
      ELSE
        T2 = TO
      ENDIF
       WRITE(*,*)'L2,L1,T3,T2',L2,L1,T3,T2
С
      R4=FUNC3(L2,L1,T3,T2,R1)
С
       WRITE(*,*)'R4 =-',R4
С
      RETURN
      END
с.
      DOUBLE PRECISION FUNCTION FUNC3 (V, W, X, Y, R1)
      COMMON/RAD/ JD,D
      REAL*8 D
      REAL*8 V,W,X,Y,PI,R1
      REAL*8 VV, WW, XX, YY, A, B, DD
      PI=3.141592654D0
      DD=DR(D)
      WW=DR(W)
      XX=DR(X)
      YY = DR(Y)
      VV=DR(V)
      A=DSIN(DD)*DSIN(WW)*(X-Y)/15.0D0
      B=DCOS(DD)*COS(WW)*(DSIN(XX+VV)-DSIN(YY+VV))*12.D0/PI
      WRITE(*,*)'R1=',R1
С
      FUNC3=R1*(A+B)
      RETURN
      END
      DOUBLE PRECISION FUNCTION FUNC2 (Y)
      COMMON/RAD/ JD,D
      REAL*8 D
      REAL*8 X, DD, YY, R1, Y
      DD=DR(D)
      YY≃DR(Y)
      X = (-DTAN(YY) * DTAN(DD))
      IF (X.GT.1.0D0) THEN
        WRITE(*,*)' SOLRAD X=',X
        X=1.D0
      ENDIF
      IF(X.LT.-1.0DO)THEN
        WRITE(*,*)' SOLRAD X=',X
        X=-1.D0
```

```
ENDIF
      X=DACOS(X)
      FUNC2=X*180.0D0/3.141592654D0
      END
      DOUBLE PRECISION FUNCTION FUNC1 (W, X, Y)
C... ESTIMATE DECLINATION, RADIUS VECTOR OF SUN, FUNC OF JULIAN DAY
      COMMON/RAD/ JD,D
      REAL*8 D
      REAL*8 W, X, Y, V, R1
      V=(DBLE(JD)+Y)*0.986D0
      V=DR(V)
      A = COS(V)
      FUNC1=W-X*A
      RETURN
      END
      DOUBLE PRECISION FUNCTION DR(I)
C... CONVERT DEGREES TO RADIANS
      REAL*8 L
      DR=1*3.141592654D0/180D0
      RETURN
      END
      DOUBLE PRECISION FUNCTION RD(1)
C... CONVERT DEGREES TO RADIANS
      REAL*8 I
      RD=1*180D0/3.141592654D0
      RETURN
      END
      SUBROUTINE FILER
$INSERT COMMON/MAIN/
$INSERT COMMON/THERM/
      REAL*8 CN(9), SPCT(9)
      REAL*8 XH, XD
      INTEGER DATE
      CHARACTER FNAME*35, HEAD*80, NAME*30, DN(10)*2
      EQUIVALENCE (ANC, CN (1))
      EQUIVALENCE (ANCT, SPCT(1))
      DATA DN/'D1','D2','D3','D4','D5','D6','D7','D8','D9','D0'/
      TAB=CHAR(137)
С
C... OPEN INPUT FILES
С
C...PARAMETERS AND DAILY SNOWMELT DATA
      WRITE(*,*) 'ENTER NAME OF MAIN INPUT FILE:'
      READ(*, '(A30)') FNAME
      NAME=FNAME
      OPEN (9, FILE=FNAME, STATUS='OLD')
C...RAIN CHEMISTRY DATA
      WRITE(*,*) 'ENTER NAME OF RAIN INPUT FILE:'
      READ(*, '(A30)') FNAME
      OPEN(10, FILE=FNAME, STATUS='OLD')
C...ICE LAYER LEVELS DATA
      WRITE(*,*) 'ENTER NAME OF ICE ELEV INPUT FILE:'
      READ(*,'(A30)') FNAME
      OPEN(18, FILE=FNAME, STATUS='OLD')
C...PREVIOUS 10 DAYS OF DISCHARGE DATA FOR MEMORY OF SYSTEM
      WRITE(*,*) 'ENTER NAME OF PRE 10 DAYS Q INPUT FILE:'
       READ(*, '(A30)') FNAME
      OPEN(17,FILE=FNAME,STATUS='OLD')
C...OBSERVED STREAM DATA TO COMPARE IONIC STRENGTHS AND R^2
      WRITE (*, *) 'ENTER NAME OF OBS STREAM INPUT FILE:'
       READ(*, '(A30)') FNAME
```

```
OPEN(11, FILE=FNAME, STATUS='OLD')
C...OBSERVED LAKE DATA TO COMPARE R^2
      WRITE(*,*) 'ENTER NAME OF OBS LAKE OUTPUT FILE:'
      READ(*, '(A30)') FNAME
      OPEN (5, FILE=FNAME, STATUS='OLD')
C...DAILY INFLOW TEMPERATURES, BEGIN W/ J-DAY OF BEGIN OF SIMULATION
      WRITE(*,*) 'ENTER NAME OF OBS STREAM TEMP FILE:'
      READ(*, '(A30)') FNAME
      OPEN (20, FILE=FNAME, STATUS='OLD')
C...THERMOCLINE OBSERVATIONS
      WRITE(*,*) 'ENTER NAME OF OBS THERMOCLINE:'
      READ(*, '(A30)') FNAME
      OPEN (29, FILE=FNAME, STATUS='OLD')
C
C... READ THERMOCLINE ELEVATIONS (MAXIMUM OF 14 VALUES), AND THE J-DAY
C
       TC=1
43
       READ(29,*,END=10)XH(IC),XD(IC)
       TC=TC+1
       GOTO 43
10
         CONTINUE
С
C... OUTPUT FILE NAMES, CREATED BASED ON NAME OF MAIN INPUT FILE
С
C...MAIN OUTPUT FILE (VISUAL)
       FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>V'//NAME(17:30)
       OPEN(15, FILE=FNAME, STATUS='UNKNOWN')
C...STREAM CALCULATIONS FILE (TAB DELIMITED FOR MACINTOSH)
       FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>S'//NAME(17:30)
       OPEN(13, FILE=FNAME, STATUS='UNKNOWN')
C...STREAM CALC. VS. OBSERVATIONS FILE (TAB DELIMITED FOR MACINTOSH)
       FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>RS'//NAME(17:30)
       OPEN (7, FILE=FNAME, STATUS='UNKNOWN')
C...LAKE CALC. VS. OBSERVATIONS FILE (TAB DELIMITED FOR MACINTOSH)
       FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>RL'//NAME(17:30)
       OPEN(6, FILE=FNAME, STATUS='UNKNOWN')
C...LAKE CALCULATIONS FILE (TAB DELIMITED FOR MACINTOSH)
       FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>L'//NAME(17:30)
       OPEN (12, FILE=FNAME, STATUS='UNKNOWN')
C...IONIC STRENGTH OF STREAM, MACINTOSH FORMAT, CRICKET BAR GRAPH STYLE
       FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>I'//NAME(17:30)
       OPEN(14, FILE=FNAME, STATUS='UNKNOWN')
C...LAKE LAYER VOLUMES
       FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>VOL'//NAME(17:30)
      OPEN(19, FILE=FNAME, STATUS='UNKNOWN')
С
C... INITIALIZE OUTPUT FILES
С
      WRITE (12.1)
1
      FORMAT(1H*)
      WRITE(12, '(13(A,A1))')'JDAY', TAB, 'Q', TAB, 'ANC LK', TAB, 'CL LK', TAB,
     &'DOC LK', TAB, 'H LK', TAB, 'NH4 LK', TAB, 'NO3 LK', TAB, 'SBC LK',
     &TAB,'SIO2 LK', TAB, 'SO4 LK', TAB, 'PH LK'
      WRITE (14, *) 'IONS', TAB, 'J DAY', TAB, 'SBC', TAB, 'ANC', TAB,
     &'CL', TAB, 'DOC', TAB, 'H', TAB, 'NO3', TAB, 'SO4', TAB, 'NH4'
      WRITE(13,1)
      WRITE(13, '(13(A, A1))')'JDAY', TAB, 'Q', TAB, 'ANC ST',
     &TAB, 'CL ST', TAB, 'DOC ST', TAB, 'H ST'
     &, TAB, 'NH4 ST', TAB, 'NO3 ST', TAB, 'SBC ST', TAB
     &, 'SIO2 ST', TAB, 'SO4 ST', TAB, '+/- BALANCE ST',
     &TAB, 'PH'
С
       WRITE(7,*)'JDAY ST', TAB, 'Q ST', TAB, 'ANC ST',
C
      &TAB, 'CL ST', TAB, 'H ST'
С
      &, TAB, 'NH4 ST', TAB, 'NO3 ST', TAB, 'SBC ST', TAB
```

```
47
```

```
&,'SIO2 ST', TAB, 'SO4 ST', TAB, 'PH ST'
С
      &, TAB, 'ANC OB', TAB, 'CL OB', TAB, 'H OB', TAB,
С
      &'NH4 OB', TAB, 'NO3 OB', TAB, 'SBC OB', TAB, 'SIO2 OB'
С
С
      &, TAB, 'SO4 OB', TAB, 'PH OB'
       WRITE(6,*)'JDAY LK', TAB, 'Q LK', TAB, 'ANC LK',
С
      &TAB, 'CL LK', TAB, 'H LK'
С
С
      &, TAB, 'NH4 LK', TAB, 'NO3 LK', TAB, 'SBC LK', TAB
      &, 'SIO2 LK', TAB, 'SO4 LK', TAB, 'PH LK'
С
      &, TAB, 'ANC OB', TAB, 'CL OB', TAB, 'H OB', TAB,
С
      &'NH4 OB', TAB, 'NO3 OB', TAB, 'SBC OB', TAB, 'SIO2 OB'
С
      &, TAB, 'SO4 OB', TAB, 'PH OB'
С
С
C... READ INITIAL DATA VALUES
C... READ OBSERVED VALUES FOR COMPARISONS
С
C...OBS STREAM DATA TO COMPARE FONIC STRENGTHS, uEQ/L
С
       T = 1
6
      READ(11, *, END=11)OSJD(I), OSANC(I), OSCL(I), OSNO3(I),
     &OSSO4(I),OSH(I),OSNH4(Í),OSSBC(I),OSIO2(I)
С
      OSDOC(I) = OSANC(I) + OSCL(I) + OSSO4(I) + OSNO3(I) - OSH(I) - OSNH4(I)
                -OSSBC(T)
     æ
С
       I = I + 1
      GOTO 6
      CONTINUE
11
С
C...OBS LAKE DATA TO COMPARE R^2, uEQ/L
С
       I = 1
7
       READ(5, *, END=12)OLJD(I), OLPH(I), OLH(I), OLNH4(I), OLANC(I),
      &OLCL(I),OLNO3(I),OLSO4(I),OLSIO2(I),OLSBC(I)
      OLDOC(I) = OLANC(I) + OLCL(I) + OLSO4(I) + OLNO3(I) - OLH(I) - OLNH4(I)
                 -OLSBC(I)
      &
      I = I + 1
       GOTO 7
      CONTINUE
12
С
C... READ RAIN DATA jday SBC(uEQ/L), CL, NO3, NH4, SO4, H, CALCULATE DOC
С
        TC=1
        READ(10, '(A80)') HEAD
8
        READ(10, *, END=9) JDR(IC), SBCR(IC), CLR(IC), NO3R(IC), NH4R(IC)
                          ,SO4R(IC),HR(IC),QFALL(IC)
      &
       DOCR(IC) = FD(HR(IC), SBCR(IC), NH4R(IC), SO4R(IC), CLR(IC), NO3R(IC))
        TC = IC + 1
        GOTO 8
9
        CONTINUE
С
C... BEGIN TO READ INPUT
С
       READ(9, '(A80)') HEAD
С
        WRITE(15,'(A80)') HEAD
C. KR, KN NEG. EXPONENTIALS, PO2, PCO2
       DO 13 I=1,7
           READ (9, '(A30) ') HEAD
13
       READ(9,*)IMOD
C... AA, BB, PO2 ARE NO LONGER USED
       READ(9, *) AA, BB, PO2, PCO2, BAP, JDBIO
C... READ INITIAL COND. OF LAKE, UEQ/L, FIRST ICE, EPILIMNION, THEN HYPOLIMNION
C... DATE, ANC, CL, H, NH4, NO3, SBC, SIO2, SO4
       READ(9,*)DATI, ANCI, CLI, HI, NH4I, NO31, SBCI, SIO2I, SO41
       READ(9,*)DATE, ANCE, CLE, HE, NH4E, NO3E, SBCE, SIO2E, SO4E
```

```
READ (9, *) DATE, ANCH, CLH, HH, NH4H, NO3H, SBCH, SIO2H, SO4H
C...NO TEMP CORRECTIONS HERE YET
      DOCE=HE+SBCE+NH4E-SO4E-CLE-NO3E-ANCE-(10.**(-2.))/HE
      DOCI=HI+SBCI+NH4I-SO4I-CLI-NO3I-ANCI-(10.**(-2.))/HI
      DOCH=HH+SBCH+NH4H-SO4H-CLH-NO3H-ANCH-(10.**(-2.))/HH
С
C... FIRST DAY, LAST DAY OF SIMULATION, FIRST DAY AND LAST DAY OF MELT
C... AND IF USE MEMORY (=1, USE MEM)
с...
      READ(9, *) JD1, JDN, JBM, JEM, MEM
С
C... READ ALL DISCHARGES IN METERS CUBED
C... SNOW MELT DATA: jday,Q(M**3)
      DO 22 I=JD1, JDN, 1
22
             READ(9,*)DAY,QI(1)
С
C... CREATE SNOW HYDROGRAPH SH()
             RAIN STORAGE
С
                             RS()
С
             RAIN RUNOFF
                             RR()
С
             SH - RS
                           SHRS()
С
       DO 199 I=1,365
         RS(I) = 0.
199
         RR(I) = 0.
       KT=1
       DO 200 I= JBM, JEM , 1
         IF (I.EQ.JDR (KI)) THEN
            RR(I) = QI(I) - QI(I-1)
            IF(RR(I).LT.0.0) RR(I)=0.0
            IF(RR(I).GT.QFALL(KI)) RR(I)=QFALL(KI)
            SH(I) = QI(I) - RR(I)
            RS(I) =QFALL(KI) -RR(I)
            SHRS(I) = SH(I) - RS(I)
            KI = KI + 1
         ELSE
            SH(I) = QI(I)
         ENDIF
С
          WRITE (*, 137) I, TAB, QI (I), TAB, SH (I), TAB, RR (I), TAB, RS (I)
137
         FORMAT(I3, A1, 4(F12.2, A1))
200
       CONTINUE
С
C... SUM DISCHARGE FOR THE MELT TIME
      QTOT=0.D0
      DO 23 I=JBM, JEM, 1
23
          QTOT=QTOT+SH(I)
С
C... BEGIN ADDITION OF DIST. MELT MODIFICATIONS
С
C... READ TOT LOAD OF IONS IN SNOWPACK IN EQUIVALENTS
      READ(9,*)SBCT,CLT,NO3T,NH4T,SO4T,HT
C... CALC THE HYDORGEN SUCH THAT THE SNOW PACK MASS HAS CHARGE BALANCE
С
       HT=CLT+NO3T+SO4T-SBCT-NH4T
       WRITE(*,*)' FILER QTOT=',QTOT
С
      DOCT=HT+SBCT+NH4T-SO4T-CLT-NO3T
      WRITE(*,*)'DOC OF SNOWPACK = ',DOCT
с.
C... READ # OF SUB UNITS, TOTAL AREA, AND THEIR RESPECTIVE AREAS
      READ(9,*)NSU, AREAT, (AREA(IA), IA=1, NSU)
С
       WRITE(*,*)'NSU=',NSU
С
C... CREATE NAME OF SUB UNITS FOR THE OUTPUT FILES
      TWP = 29
      DO 111 I=1,NSU
        FNAME='ILWAS>OUT.WY'//NAME(17:18)//'>'//DN(I)//NAME(17:30)
```

```
OPEN (1WP, FILE=FNAME, STATUS='UNKNOWN')
       WRITE (IWP, 1)
       WRITE(IWP, '(13(A,A1))')'JDAY', TAB, 'Q', TAB, 'ANC ST',
     & TAB, 'CL ST', TAB, 'DOC ST', TAB, 'H ST'
     & ,TAB,'NH4 ST', TAB, 'NO3 ST', TAB, 'SBC ST', TAB
     & ,'SIO2 ST', TAB, 'SO4 ST', TAB, '+/- BALANCE ST',
     & TAB, 'PH'
      IWP=IWP+1
111
    CONTINUE
С
C... CHECK AREA INPUTS
     DO 5 IA=1,NSU
5
       AT=AT+AREA (IA)
      IF ((AT/AREAT).GT.1.1.OR. (AT/AREAT).LT..9) THEN
        WRITE(*,*)' THE AREAS OF THE SUB UNITS DO NOT SUM UP'
          ZZZ=AT/AREAT
          WRITE(*,*)' RATIO ASUM/AREAT=',ZZZ
        DO 71 I-1,NSU
71
          WRITE(*,*)AREA(I)
        WRITE(*,*)' AREA TOTAL INPUT =', AREAT
        STOP
      ENDIF
С
C... READ IN SUB UNITS' SLOPE, ASPECT AND LATITUDE
      DO 67 IA=1,NSU
      READ(9,*) DUMMY, RSLOPE(1A), RASPECT(IA), RLAT(IA)
               ,MSEL(IA,1),MSEL(IA,2),CF(TA)
     &
67
     CONTINUE
C... END ADDITION FOR DIST MELT
C
C... OPEN PREVIOUS 10 DAYS OF DISCHARGE DATA FOR MEMORY OF SYSTEM
С
      DO 91 IM=JD1-10,JD1-1
         READ(17, *)QI(IM)
91
         SH(IM)=QI(IM)
С
C... READ ICE LEVEL DATA
      DO 92 ICE=JD1, JDN
92
            READ(18, *) IID, BICE(1CE)
C92
         WRITE(*, *) ICE, BICE(ICE)
С
C... ECHO INPUTS SOFAR
С
       WRITE(15,*)' '
С
       WRITE(15,*)' KR=10**-',AA
С
       WRITE(15,*)' KN=10**-',BB
С
       WRITE (15, *) ' PCO2=', PCO2
С
       WRITE(15,*)' PO2--',PO2
С
       WRITE(15,*)''
С
С
       WRITE(15,*)' BEGIN DAY= ',JD1
С
       WRITE(15,*)' END DAY= ', JDN
      WRITE(15,*)' '
С
      XXXX=10.5-BICE1
С
      WRITE(15,*)' DEPTH OF ICE= ',XXXX
      PH=-LOG10(HE/1000000.)
С
      WRITE(15,*)' INITIAL LAKE CONC. uEQ/L,
                                                   PH=- PH
       WRITE(15,4)
С
4
      FORMAT(2X, 'ANC', 4X, 'CL', 3X, 'DOC', 4X, 'H', 4X,
     &'NH4', 3X, 'NO3', 3X, 'SBC', 2X, 'SIO2', 3X, 'SO4')
С
       WRITE (15, 3) ANC, CL, DOC, H, NH4, NO3, SBC, SIO2, SO4, PH
      FORMAT(14(F5.1,1X),/)
3
С
      RETURN
      END
```

```
DOUBLE PRECISION FUNCTION FD (H, SBC, NH4, SO4, CL, NO3)
      REAL*8 H,SBC,NH4,SO4,CL,NO3
C...CONVERT uEQ/L TO EQ/L (WHICH IS SAME AS MOLES, IN THIS CASE)
        H=H/1000000.
        SBC=SBC/1000000.
        NH4=NH4/1000000.
        SO4=SO4/1000000.
        CL=CL/1000000.
        NO3=NO3/1000000.
C...CALCULATE DOC AND CONVERT BACK TO uEQ/L
        FD=(H+SBC+NH4-SO4-CL-NO3-(10.**(-14.))/H)*1000000.
        H=H*1000000.
        SBC=SBC*1000000.
        NH4=NH4*1000000.
        SO4=SO4*1000000.
        CL=CL*1000000.
        NO3=NO3*1000000.
      RETURN
      END
 _
      SUBROUTINE RAIN
$INSERT COMMON/MAIN/
      REAL*8 CN(9), SPCT(9)
      EQUIVALENCE (ANC, CN(1))
      EQUIVALENCE (ANCT, SPCT(1))
      TAB=CHAR(137)
C... TEST FOR RAIN
C... IF ITS A RAIN DAY, REPLACE SNOWMELT CHEM WITH RAIN CHEM
        IFLAG=0
С
         WRITE(*,*)' JDAY, JDAY OF RAIN', JDAY, JDR(IR)
        IF (JDAY.EQ.JDR(IR)) THEN
          IRAIN1=1
          SBC=SBCR(IR)
          CL=CLR(IR)
          NO3=NO3R(IR)
          NH4=NH4R(IR)
          SO4=SO4R(IR)
          H=HR(IR)
          DOC=DOCR(IR)
          IF (IYEST.NE.JDAY) THEN
            IR=IR+1
          ENDIF
          IYEST=JDAY
          IFLAG=1
          WRITE(15,*)'
                         RAIN DAY !
        END IF
        WRITE(15,*)'
                       JULIAN DAY', JDAY
        WRITE(15,*)'
                        DISCHARGE M**3', QI(JDAY)
        WRITE(15,*)' '
2
        FORMAT(6X,6(F5.1,1X),6X,F5.1,/)
      FORMAT(2X, 'ANC', 4X, 'CL', 3X, 'DOC', 4X, 'H', 4X,
4
     &'NH4', 3X, 'NO3', 3X, 'SBC', 2X, 'SIO2', 3X, 'SO4')
C... THESE ARE THE INPUT DATA USED TO COMPARE IONIC STRENGTH
        IPDOC=DOC
         IPSBC=SBC
        IPCL=CL
        IPNO3≕NO3
        IPNH4=NH4
        IPSO4=SO4
        IPH=H
С
        RETURN
```

```
END
```

_

```
SUBROUTINE NRWEATH
С
      *** NEWTON RALPHSON: WEATHERING ***
С
С
С
       REMOVED NITROGEN EQUATION; NOW ALL NH4=>H
С
С
       INPUT: PCO2, PH, NH4, SBC, CL, SO4, NO3, ANC
       REEQUILIBRATE: H, NO3, NH4, HCO3, CO3
С
       CALCULATE: SIO2, SBC
С
С
С
      INPUT IS IN EQ/L. OUTPUT IS IN EQ/L.
       EXCEPTION: SIO2 IS ALWAYS IN MOLES/L.
С
       INTERNAL EQUILIBRIUM CALCULATIONS ARE IN MOLES/L.
С
С
С
          X,Y,Z = UNKNOWNS
С
               Х = Н+
С
С
               Y - NO3-
С
               Z = SiO2
С
С
       Kx = EQUILIBRIUM CONST. H - HENRY'S CO2
С
                                R - ROCK WEATHERING
                                W - WATER
С
                              1,2 - pK'S FOR CARBONATE SYSTEM
С
С
С
      PCO2 = PRESSURE OF ATM. CO2 GAS
С
С
С
       C = NH4 - CL - NO3 - SO4 - DOC
С
С
       Y1, Y2, Y3 = GOVERNING EQUATIONS = 0.
С
       Y1X = DERIVATIVE OF Y1 W/R/T X
С
С
С
С
С
       Y3Z = DERIVATIVE OF Y3 W/R/T Z
С
       DV = V(OLD) - M(NEW), etc.
С
С
С
       GIVEN: INITIAL ESTIMATES FOR UNKNOWNS
С
       SOLVE BY NEWTON-RAPHSON TO FIND NEW V, W, X, Y, Z
С
С
       CALCULATE NEW YI'S AND TEST FOR EQUALITY TO ZERO
С
С
       IF TEST FAILED, THEN NEW V,W,X,Y,Z VALUES ARE
С
       INITIAL ESTIMATES FOR NEXT ITERATION
С
С
С
$INSERT COMMON/MAIN/
      REAL*8 CN(9)
      REAL*8 HCO3, NO3ST, NO3B1
      EQUIVALENCE (ANC, CN(1))
      TAB=CHAR(137)
      BUG=0
C... OPEN OUTPUT FILE
      OPEN (16, FILE='IT.LIST', STATUS='UNKNOWN')
С
C... ITERATION COUNTER
      IC = 1
C... TOLERANCE OF Y CONVERGENCE TEST
```

```
TOL=0.00000000000001
C... RATE CONSTANTS
C... EVALUATE TEMPERATURE
Ċ
        TO = REF. TEMP
С
        KHO = HENRYS K FOR CO2(g) AT TO
С
        ST = STREAM TEMP (KELVIN)
с...
С
       KW=10.**(-14.)
       K1=10.**(-6.35)
С
       K2=10.**(-10.33)
С
С
      K1=10.0**((-3404./ST) - 0.0328*ST + 14.844)
      K2=10.0**((-2902./ST) - 0.0238*ST + 6.498)
      KH=EXP((-4850/R)*(1/TO-1/ST) + DLOG(KHO))
      KW=10.0**(-4470.99/ST - 0.01706*ST + 6.0875)
С
C... THIS IS OLD STUFF, AND ISN'T USED. RUN THE MODEL WITH MEMORY (MEM=1)
      IF (MEM.EQ.0) THEN
      IF (IMOD.EQ.1) THEN
        C SERIES USES KR VALUE IN INPUT FILE (EQUILIBRIUM MODEL)
C...
        KR=10.**(-AA)
      ELSEIF (IMOD.EQ.2) THEN
с...
        THIS IS WHERE WE LET K=F(Q) (KINETIC MODEL, NO MEM)
        R^2 = 0.734
        KR=0.30104*QT**(-.63452)
      ELSE
        THIS IS WHERE WE LET K=F(Q) (KINETIC MODEL)
с...
        R^2=0.721
                     EXPONENTIAL FIT
С
        KR=3.052*10.**(-3.0)*10.**(-4.0773*10.**(-5.)*QT)
      ENDIF
С
      ELSEIF (MEM.EQ.1) THEN
с...
        HERE WE USE 10 DAY MEMORY IF MEM=1
        IMOD = 1, ACTUAL EQUATION
С
С
        IMOD = 2, LOWER BOUND
        IMOD = 3, UPPER BOUND
С
        BOUNDS ON THE SLOPE OF .002 AND .006
С
C...NEW STOICHIOMETRY OF 1.2 H : 1.2 SBC, SBC/SIO2 = 1.2
C...NEW KR, 89/03/02 ACTUAL
C...NEW KR,89/03/02, UPPER BOUND
C...NEW KR,89/03/02, LOWER BOUND
        IF (IMOD.EQ.1) THEN
          KR=(0.0081653D0)*10.**(-0.000043561D0 * QT)
        ELSEIF (IMOD.EQ.2) THEN
          KR=(0.015D0)*10.**(-0.000043561D0 * QT)
        ELSEIF (IMOD.EQ.3) THEN
          KR=(0.005D0)*10.**(-0.000043561D0 * QT)
        ELSEIF (IMOD.EQ.1) THEN
          WRITE(*,*) 'CHECK INPUT: LOG OR EXP FIT, Q AVERAGE'
          STOP
        ENDIF
      ENDIF
C... NEED DUMMY VALUE FOR SIO2
      SI02=1.0
C... CONVERT uEQ/L TO EQ/L
С
      DO 5 JK=2,9
5
        CN(JK) = (CN(JK) / 1000000.)
       WRITE(*,*)'ANC=',ANC
С
       WRITE(*,*)'CL=',CL
С
       WRITE (*, *) 'DOC=', DOC
С
С
       WRITE(*,*)'H=',H
       WRITE(*,*)'NH4=',NH4
С
       WRITE (*, *) 'NO3=', NO3
С
```

```
WRITE(*,*)'SBC=',SBC
С
       WRITE(*,*)'SIO2=',SIO2
С
       WRITE(*,*)'SO4=',SO4
С
С
С
C... ALL NH4 GOES TO H
C... THEN A % OF NO3 IS RETAINED BY BIOLOGICAL ACTIVITY.
C... THIS VARIABLE IS Biological Activity Percent (BAP)
с...
      CAT=NH4+SBC+H
      ANI=CL+NO3+SO4
      CB=(CAT-ANI) / (CAT+ANI)
C... NITROGEN CYCLE
           NO3ST = NO3 LEFT IN STREAM
с...
           NO3BI = NO3 CONSUMED BY BIOTA
с...
      IF (JDAY.LT.JDBIO) THEN
        NO3 = NO3 + NH4
        H = H + 2. * NH4
        NH4 = 0.0
      ELSEIF (JDAY.GE.JDBIO) THEN
        NO3ST = (1.-BAP) * NO3
        NO3BI = (BAP) * NO3
        NO3 = NO3ST
        H = H - NO3BI
        H = H + NH4
        NH4 = 0.
      ENDIF
      C=NH4-CL-SO4-NO3-DOC
       WRITE(*,*)' '
С
С
       WRITE(*,*)' NRWEATH C=',C
C... INITIAL CONDITION, UNKNOWNS
      X=10.**(-5.1)
      Z=SIO2
C... THE Y EQUATIONS
      A = K1*KH*PCO2
10
      Y2= Z*(Z+SBC)**1.2 - KR*X**1.2
      Y3 = Z + SBC + X + C - \Lambda / X - A * K2 / X * *2 - KW / X
       IF (BUG.EQ.1) THEN
С
          WRITE(16,*)'INITIAL Y2=',Y2
          WRITE(16,*)'
                              ¥3=',¥3
С
         WRITE(16,*)' '
С
      ENDIF
C... THE PARTIALS OF Y1, ..., Y3 W/R/T X, ..., Z
       Y2X= -1.2*KR*X**0.2
       Y2Z= (Z+SBC) **1.2 + 1.2*Z*(Z+SBC) **0.2
       Y3X= 1.+A/X**2.+2.*A*K2/X**3.+KW/X**2.
       Y3Z= 1.
       IF (BUG.EQ.1) THEN
          WRITE(16,*)'Y2X=',Y2X
С
С
          WRITE(16,*)'Y2Z=',Y2Z
С
          WRITE(16, *) 'Y3X=', Y3X
          WRITE(16,*)'Y3Z=',Y3Z
С
         WRITE(16,*)''
С
       ENDIF
C... SOLVE FOR DX, DY, DZ
      DETA=DET (Y2X, Y2Z, Y3X, Y3Z)
       IF (DETA.LT.1E-10) WRITE (*,*) 'DETA=', DETA
С
       DX=DET( Y2, Y2Z, Y3, Y3Z)
       DZ=DET(Y2X,Y2,Y3X,Y3)
       IF (BUG.EQ.1) THEN
С
          WRITE(16,*)'DET=',DETA
С
          WRITE(16,*)'DETX=',DX
          WRITE(16, *) 'DETZ=', DZ
С
       ENDIF
```

```
DX=DX/DETA
      DZ=DZ/DETA
      IF (BUG.EQ.1) THEN
С
         WRITE(16, *) 'DX=', DX
С
         WRITE (16, *) 'DZ=', DZ
С
         WRITE(16,*)' '
      ENDIE
C... SOLVE FOR NEW VALUES OF W, X, Y, Z
      IF (BUG.EQ.1) THEN
С
         WRITE(16,*)' THE OLD X',X
         WRITE(16,*)' SOLVING FOR NEW X'
С
      ENDIF
      X=BETA(X,DX)
      IF (BUG.EQ.1) THEN
С
         WRITE(16, *) 'THE NEW X', X
         WRITE(16,*)' '
С
         WRITE(16,*)' THE OLD Z',Z
С
С
         WRITE (16, *) ' SOLVING FOR NEW Z'
      ENDIF
C... TEST TO ASSURE NIETHER SIO2 OR SBC CONC. DROPS BELOW ZERO
      Z = BETA(Z, DZ)
      IF (BUG.EQ.1) THEN
С
         WRITE (16, *) 'THE NEW Z', Z
      ENDIF
C... TEST FOR CONVERGENCE
      Y2= Z*(Z+SBC)**1.2 - KR*X**1.2
      Y3= Z+ SBC+X+C-A/X-A*K2/X**2.-KW/X
      IF (ABS(Y2).LT.TOL.AND.ABS(Y3).LT.TOL) THEN
         PH = -LOG10(X)
         CO3 = K1 * K2 * KH * PCO2 / X * * 2.
         HCO3 =K1*KH*PCO2/X
         Н
             =X
         ANC = HCO3+2*CO3+KW/H-H
C**
            ANC= HCO3
         SBC =SBC+Z
         SIO2 =Z
C... CONVERT FROM EQUIVALENTS TO MICRO EQUIVALENTS
        DO 6 JK=1,9
6
          CN(JK)=CN(JK)*1000000.
С
C... PROGRAM OUTPUT
      FORMAT(2X, 'ANC', 4X, 'CL', 3X, 'DOC', 4X, 'H', 4X,
4
     & 'NH4', 3X, 'NO3', 3X, 'SBC', 2X, 'SIO2', 3X, 'SO4')
с...
      FORMAT(13(F5.1,1X),/)
3
        RETURN
      ELSE
36
        IC=IC+1
      ENDIF
C... RUNAWAY CATCHER
      IF (IC.GT.190) THEN
        WRITE(*,*)' MAX ITERATIONS EXCEEDED'
        STOP
      ENDIF
      GOTO 10
      END
      DOUBLE PRECISION FUNCTION DET(A, B, C, D)
      INTEGER BUG
      REAL*8 A, B, C, D
C... CALCULATES THE DETERMINATE OF A 2 X 2 MATRIX
      DET=A*D-C*B
      RETURN
      END
```

```
DOUBLE PRECISION FUNCTION BETA(XC, DXC)
SINSERT COMMON/MAIN/
     REAL*8 XC,DXC,U
11
     U=XC
      IF (BUG.EQ.1) THEN
        WRITE (16, *) 'DU=', DXC
C
     ENDIF
      U=U-DXC
     IF (BUG.EQ.1) THEN
       WRITE(16,*)'U=',U
С
      ENDIF
      IF (U.LT.O.) THEN
       DXC=DXC/10.
       GOTO 11
      ELSE
       BETA=U
      ENDIF
      RETURN
      END
 _
      SUBROUTINE NROPEN
С
С
     *** NEWTON RALPHSON: OPEN CARBONATE
С
С
           UNKNOWNS
С
С
               X = H +
               Z = ANC
С
С
       Kx = EQUILIBRIUM CONST. H - HENRY'S CO2
С
С
                                R - ROCK WEATHERING
                                W - WATER
С
                              1,2 - pK'S FOR CARBONATE SYSTEM
С
С
С
      PCO2 = PRESSURE OF ATM. CO2 GAS
С
С
       C = NH4 - CL - NO3 - SO4 - DOC
С
С
С
       Y1, Y2 = GOVERNING EQUATIONS = 0.
С
       Y1X == DERIVATIVE OF Y1 W/R/T X
С
С
       •
С
       .
С
       Y3Z = DERIVATIVE OF Y3 W/R/T Z
С
С
С
      DV = V(OLD) - M(NEW), etc.
С
       GIVEN: INITIAL ESTIMATES FOR UNKNOWNS
С
С
С
       SOLVE BY NEWTON-RAPHSON METHOD
С
       CALCULATE NEW YI'S AND TEST FOR EQUALITY TO ZERO
С
C
      IF TEST FAILED, THEN NEW V,W,X,Y,Z VALUES ARE
С
       INITIAL ESTIMATES FOR NEXT ITERATION
С
С
С
$INSERT COMMON/MAIN/
      REAL*8 CN(9)
      REAL*8 HCO3
```

```
EQUIVALENCE (ANC, CN (1))
      TAB=CHAR(137)
      BUG=0
C... OPEN OUTPUT FILE
C... ITERATION COUNTER
      IC=1
C... TOLERANCE OF Y CONVERGENCE TEST
      TOL=0.0000000000001
C... RATE CONSTANTS
C... RATE CONSTANTS
C... EVALUATE TEMPERATURE
        TO = REF. TEMP
С
         KHO = HENRYS K FOR CO2(g) AT TO
С
С
         ST = STREAM TEMP
с...
С
      Kl=10.0**((-3404./ST) - 0.0328*ST + 14.844)
       K2=10.0**((-2902./ST) - 0.0238*ST + 6.498)
      KH = EXP((-4850/R) * (1/TO - 1/ST) + DLOG(KHO))
      KW=10.0**(-4470.99/ST - 0.01706*ST + 6.0875)
С
C... CONVERT uEQ/L TO EQ/L
С
      DO 5 JK=1,9
5
       CN(JK) = (CN(JK) / 1000000.)
С
с...
      C=SBC-NH4-CL-SO4-NO3-DOC
C... INITIAL CONDITIONS
      X = H
      Z=ANC
C... THE Y EQUATIONS
      A = K1 * KH * PCO2
10
      Y2 = Z - A/X
      Y3= X+C-Z-A*K2/X**2.-KW/X
      IF (BUG.EQ.1) THEN
С
         WRITE(16, *)'INITIAL Y2=', Y2
         WRITE(16,*)'
С
                             Y3=',Y3
С
         WRITE(16,*)' '
      ENDIF
C... THE PARTIALS OF Y1,...,Y3 W/R/T X,...,Z
      Y2X = A/X * * 2.
      Y2Z= 1.
      Y3X= 1.+2.*A*K2/X**3.+KW/X**2.
      Y3Z= -1.
      IF (BUG.EQ.1) THEN
С
         WRITE(16,*) 'Y2X=',Y2X
С
         WRITE (16, *) 'Y2Z=', Y2Z
         WRITE(16,*)'Y3X=',Y3X
С
         WRITE(16,*)'Y3Z=',Y3Z
С
С
         WRITE(16,*)''
      ENDIF
C... SOLVE FOR DX, DY, DZ
      DETA=DET(Y2X,Y2Z,Y3X,Y3Z)
С
       IF (DETA.LT.1E-10) WRITE (*, *) 'DETA=', DETA
      DX=DET( Y2, Y2Z, Y3, Y3Z)
      DZ=DET(Y2X,Y2,Y3X,Y3)
      IF (BUG.EQ.1) THEN
С
         WRITE(16, *) 'DET=', DETA
С
         WRITE(16, *) 'DETX=', DX
С
         WRITE (16, *) 'DETZ=', DZ
      ENDIF
      DX=DX/DETA
      DZ=DZ/DETA
```

```
IF (BUG.EQ.1) THEN
         WRITE(16,*)'DX=',DX
С
         WRITE (16, *) 'DZ=', DZ
С
С
         WRITE(16,*)' '
      ENDIF
C... SOLVE FOR NEW VALUES OF W, X, Y, Z
      IF (BUG.EQ.1) THEN
         WRITE(16,*)' THE OLD X',X
WRITE(16,*)' SOLVING FOR NEW X'
С
С
      ENDIF
      X = BETA(X, DX)
      IF (BUG.EQ.1) THEN
С
         WRITE(16,*)'THE NEW X',X
         WRITE(16,*)' '
С
          WRITE(16,*)' THE OLD Z',Z
С
          WRITE(16,*)' SOLVING FOR NEW Z'
С
      ENDIF
C... TEST TO ASSURE NIETHER SIO2 OR SBC CONC. DROPS BELOW ZERO
      Z = BETA(Z, DZ)
      IF (BUG.EQ.1) THEN
          WRITE(16,*)'THE NEW Z',Z
С
      ENDIF
C... TEST FOR CONVERGENCE
      Y2 = Z - A / X
       Y3- X+C-Z-A*K2/X**2.-KW/X
       IF (ABS(Y2).LT.TOL.AND.ABS(Y3).LT.TOL) THEN
          PH = -LOG10(X)
          CO3 = K1 * K2 * KH * PCO2 / X * * 2.
          HCO3 =Z
          Н
             ==X
          ANC = HCO3 + 2 CO3 + KW/H - H
          ANC= HCO3
С
C... CONVERT FROM EQUIVALENTS TO MICRO EQUIVALENTS
         DO 6 JK=1,9
           CN(JK)=CN(JK)*1000000.
 6
         RETURN
       ELSE
С
          WRITE(16,*)' '
          WRITE(16,*)' ITERATION NUMBER ', IC
С
                         Y2= ',Y2
          WRITE(16,*)'
С
                           Y3= ',Y3
С
          WRITE(16,*)'
          WRITE(16,*)' H
                              X= ',X
С
          WRITE(16,*)' '
С
36
        IC = IC + 1
C... TEMPORARY STOP
         STOP
С
       ENDIF
C... RUNAWAY CATCHER
       IF (IC.GT.190) THEN
         WRITE(*,*)' MAX ITERATIONS EXCEEDED OPEN CARB SYS'
         STOP
       ENDIF
       GOTO 10
       END
  _
       SUBROUTINE NRCLOSE
 С
         *** NEWTON RALPHSON: CLOSED CARBONATE SYSTEM ***
 С
 С
 С
             UNKNOWNS
 С
                 X = H +
 С
                 Z = ANC
 С
```

```
С
С
       Kx = EQUILIBRIUM CONST. H - HENRY'S CO2
С
                               R - ROCK WEATHERING
С
                               W - WATER
                             1,2 - pK'S FOR CARBONATE SYSTEM
С
С
С
С
      C = NH4 - CL - NO3 - SO4 - DOC
С
С
      Y1, Y2 = GOVERNING EQUATIONS = 0.
С
С
      Y1X = DERIVATIVE OF Y1 W/R/T X
С
С
       .
С
       .
С
С
      Y3Z = DERIVATIVE OF Y3 W/R/T Z
С
С
      DV = V(OLD) - M(NEW), etc.
С
С
      GIVEN: INITIAL ESTIMATES FOR UNKNOWNS
С
С
      SOLVE BY NEWTON-RAPHSON METHOD
С
      CALCULATE NEW YI'S AND TEST FOR EQUALITY TO ZERO
С
С
С
       IF TEST FAILED, THEN NEW V,W,X,Y,Z VALUES ARE
       INITIAL ESTIMATES FOR NEXT ITERATION
С
С
С
$INSERT COMMON/MAIN/
      REAL*8 CN(9)
      REAL*8 HCO3
      EQUIVALENCE (ANC, CN(1))
      TAB=CHAR(137)
      BUG=1
C... OPEN OUTPUT FILE
С
      OPEN (16, FILE='IT.LIST3', STATUS='UNKNOWN')
C... ITERATION COUNTER
      TC=1
C... TOLERANCE OF Y CONVERGENCE TEST
      TOL=0.00000000001
C... RATE CONSTANTS
C... EVALUATE TEMPERATURE
       TO = REF. TEMP
С
С
         KHO = HENRYS K FOR CO2(g) AT TO
С
         ST = STREAM TEMP
с...
С
      K1=10.0**((-3404./ST) - 0.0328*ST + 14.844)
      K2=10.0**((-2902./ST) - 0.0238*ST + 6.498)
      KH=EXP((-4850/R)*(1/TO-1/ST) + DLOG(KHO))
      KW=10.0**(-4470.99/ST - 0.01706*ST + 6.0875)
С
C... CONVERT uEQ/L TO EQ/L
С
      DO 5 JK=1,9
5
       CN(JK) = (CN(JK) / 1000000.)
С
C... CLOSED SYS, CONSERVE TOTAL CARBON (CT)
      CT=ANC+ANC*((H/K1+K2/H))
      WRITE(*,*)'CT=',CT
С
      C=SBC-NH4-CL-SO4-NO3-DOC
C... INITIAL CONDITIONS
```

```
X = H
      Z=CN(1)
      IF (BUG.EQ.1) THEN
         WRITE(16, *)'INITIAL X=',X
С
         WRITE(16,*)'
                               Z=',Z
С
         WRITE(16,*)' '
С
      ENDIF
C... THE Y EQUATIONS
      Y2 = -CT + Z * X / K1 + Z + K2 * Z / X
10
      Y3 = X+C-KW/X-Z-K2*Z/X
      IF (BUG.EQ.1) THEN
          WRITE(16,*)'INITIAL Y2=',Y2
С
С
          WRITE(16,*)'
                              Y3=',Y3
         WRITE(16,*)' '
С
      ENDIF
C... THE PARTIALS OF Y1,...,Y3 W/R/T X,...,Z
      Y2X= Z/K1+K2*Z/X**2.
      Y2Z= X/K1+1.+K2/X
      Y3X= 1.+KW/X**2.+K2*Z/X**2.
      Y3Z- -1.-K2/X
С
       IF (BUG.EQ.1) THEN
           WRITE(16,*)'Y2X=',Y2X
СС
           WRITE(16,*)'Y2Z=',Y2Z
CC
           WRITE(16,*)'Y3X=',Y3X
CC
           WRITE(16,*)'Y3Z=',Y3Z
СС
СС
           WRITE(16,*)' '
С
       ENDIF
C... SOLVE FOR DX, DY, DZ
      DETA=DET (Y2X, Y2Z, Y3X, Y3Z)
       IF (ABS (DETA) .LT.1E-10) WRITE (*, *) 'DETA=', DETA
      DX=DET( Y2, Y2Z, Y3, Y3Z)
      DZ=DET(Y2X,Y2,Y3X,Y3)
       IF (BUG.EQ.1) THEN
С
          WRITE(16, *) 'DET=', DETA
          WRITE(16,*)'DETX=',DX
С
С
          WRITE(16, *) 'DETZ=', DZ
       ENDIF
      DX=DX/DETA
      DZ=DZ/DETA
       IF (BUG.EQ.1) THEN
          WRITE (16, *) 'DX=', DX
С
С
          WRITE(16,*)'DZ=',DZ
          WRITE(16,*)' '
С
      ENDIF
C... SOLVE FOR NEW VALUES OF W, X, Y, Z
       IF (BUG.EQ.1) THEN
           WRITE(16,*)' THE OLD X',X
СС
           WRITE (16, *) ' SOLVING FOR NEW X'
CC
       ENDIF
       X = BETA(X, DX)
       IF (BUG.EQ.1) THEN
CC
           WRITE(16,*)'THE NEW X',X
           WRITE(16,*)' '
СС
СС
           WRITE(16,*)' THE OLD Z',Z
           WRITE(16,*)' SOLVING FOR NEW Z'
CC
       ENDIF
C... TEST TO ASSURE NIETHER STO2 OR SBC CONC. DROPS BELOW ZERO
       Z = BETA(Z, DZ)
       IF (BUG.EQ.1) THEN
           WRITE(16,*)'THE NEW Z',Z
СС
       ENDIF
C... TEST FOR CONVERGENCE
       Y_{2} = -C_{T} + Z \times X / K_{1} + Z + K_{2} \times Z / X
       Y3 = X + C - KW / X - Z - K2 + Z / X
```

```
IF (BUG.EQ.1) THEN
CC
         WRITE(16,*)' NEW
                             Y2=',Y2
          WRITE(16,*)'
                             Y3≔',Y3
СС
          WRITE(16,*)' '
СС
      ENDIF
      IF (ABS(Y2).LT.TOL.AND.ABS(Y3).LT.TOL) THEN
        WRITE(*,*)' FINAL Y2=',Y2
С
         WRITE(*,*)'
С
                            Y3=',Y3
         WRITE(*,*)'X=',X
С
С
         WRITE(*,*)'Z=',Z
         HCO3 =Z
         CO3 = K2 * HCO3
         H =X
         ANC = HCO3+2*CO3+KW/H-H
С
         ANC= HCO3
         PH=-LOG10(X)
C... CONVERT FROM EQUIVALENTS TO MICRO EQUIVALENTS
        DO 6 JK=1,9
         CN(JK)=CN(JK)*1000000.
6
        RETURN
      ELSE
        WRITE(16,*)''
С
         WRITE(16,*)' ITERATION NUMBER ',IC
С
         WRITE(16,*)' Y2= ',Y2
С
         WRITE(16,*)'
                        Y3= ',Y3
С
        WRITE(16,*)' '
С
36
        IC=IC+1
C... TEMPORARY STOP
С
        STOP
      ENDIF
C... RUNAWAY CATCHER
      IF (IC.GT.1490) THEN
        WRITE (*, *) ' MAX ITERATIONS EXCEEDED CLOSED CARB SYS'
        STOP
      ENDIF
      GOTO 10
      END
 ----
      SUBROUTINE REQ(VICE)
С
C... REEQUILIBRATE LAKE EPI LAYER
C... NROPEN = OPEN SYSTEM
C... NRCLOSE = CLOSED SYSTEM
С
      IF (VICE.LE.O.O) THEN
        CALL NROPEN
      ELSE
        CALL NRCLOSE
      ENDIF
      RETURN
      END
      SUBROUTINE LAKE(IKI)
С
      TAKES PREVIOUS CONC OF EP. AND HY. LAYERS AND MIXES THEM
С
       AS THE THERMOCLINE CHANGES.
С
       VT= VOL TOTAL
       VE= VOL EPILIMNION, NEW (TODAY)
С
С
       VE1= VOL EPILIMNION, OLD (YESTERDAY)
       VH= VOL HYPO NEW
С
С
       VH1= VOL HYPO OLD
       VHD= DIFFERENCE IN HYPO VOLUME DUE TO CHANGE IN THERMOCLINE
С
```

```
61
```

```
DIV = DIFFERENC IN ICE VOLUME DUE TO MELT/FREEZING
С
С
       VICE - ICE VOLUME
       BICE() = ELEVATION OF ICE BOUNDARY ABOVE LAKE BOTTOM
С
       CI1() = CHEM CONC. IN ICE
С
С
       CI2() = NEW DAY CONC. IN ICE
С
       TAKES CONC: VALUES IN COMMON AND MIXES THEM WITH CONC.
C
       OF LAKE FROM PREVIOUS TIME STEP, IN ARRAY CE1, CH1, WHICH IS INTIALIZED
С
       IN THE FIRST CALL TO THIS SUB FROM MAIN, WHEN ARGUMENT I = 1.
С
С
       ARRAY CN IS EQUIVALENCED WITH THE IN COMING CONCENTRATIONS
С
       AND THE OUT GOING NEW CONCENTRATIONS.
С
С
       ARRAY CE1, CH1 IS THE MEMORY OF THE LAKE. HERE WE STORE FOR ONE TIME
С
       STEP THE LAST VALUES OF CONCENTRATION OF THE LAKE.WHEN IKI=1
С
С
       THE CAT/ANION VALUES ARE READ INTO CE1, CH1
С
       SUBROUTINE THERM WILL DETERIMINE THE HEIGHT OF THE THERMOCLINE
С
       BASED ON THE DATE, AND RETURN VOLUME OF THE EPILIMNION.
С
$INSERT COMMON/MAIN/
      REAL*8 CN(9)
      EQUIVALENCE (ANC, CN(1))
      TAB=CHAR(137)
      VT=179600.0
С
C... IF BLOCK EXECUTED ONCE IN BEGINNING OF SIMULATION, TO INITIALIZE
C... LAKE CONCENTRATIONS, AND OPEN ANY FILES
С
      IF (IKI.EQ.1) THEN
       OPEN(29,FILE='ILWAS>OUTPUT>MIX.ANC',STATUS='UNKNOWN')
С
C... FOLLOWINGOPEN STATEMENTS USED TO CHECK MIXING OF LAKE.
       WRITE (29, *) 'JDAY', TAB, 'EPI VOL', TAB, 'E CON ', TAB, 'EPI VOLAI'
C
      &, TAB, 'EC AI', TAB, 'EV AH', TAB, 'EC AH', TAB, 'STR VOL', TAB,
С
С
      &TAB, 'STR CONC', TAB, 'SV AM', TAB, 'SC AM'
       OPEN(30, FILE='ILWAS>OUTPUT>MIX.SIO2', STATUS='UNKNOWN')
С
       OPEN(31,FILE='ILWAS>OUTPUT>MIX.SBC',STATUS='UNKNOWN')
С
С
       OPEN(32, FILE='ILWAS>OUTPUT>MIX.H', STATUS='UNKNOWN')
       OPEN(33, FILE='ILWAS>OUTPUT>MIX.CL', STATUS='UNKNOWN')
С
       OPEN(34,FILE='ILWAS>OUTPUT>MIX.NO3',STATUS='UNKNOWN')
С
       OPEN(35,FILE='ILWAS>OUTPUT>MIX.NH4',STATUS='UNKNOWN')
С
       OPEN(36, FILE='ILWAS>OUTPUT>MIX.SO4', STATUS='UNKNOWN')
С
C...ICE
        CI1(1)=ANCI
        CI1(2)=CLI
        CI1(3)=DOCI
        CI1(4)=HI
        CI1(5)=NH4I
        CI1(6) = NO3T
        CI1(7)=SBCI
        CI1(8)=SIO2I
        CI1(9)=SO4I
C...EPILIMNION
         CE1(1) = ANCE
         CE1(2) = CLE
         CE1(3) = DOCE
         CE1(4) = HE
         CE1(5)=NH4E
         CE1(6)=NO3E
         CE1(7) = SBCE
         CE1(8)=SIO2E
         CE1(9)=SO4E
C...HYPOLIMNION
         CH1(1) = ANCH
         CH1(2) = CLH
```

```
CH1(3)=DOCH
         CH1(4)=HH
         CH1(5)=NH4H
         CH1(6)=NO3H
         CH1(7) = SBCH
         CH1(8)=SIO2H
         CH1(9)=SO4H
         RETURN
      ENDIF
С
с.
с..
C... CALL THERMOCLINE TO FIND VOLUME OF EPILIMNION AND ICE LAYER
C
       CALL THERM (JDAY, BICE (JDAY), VE, VICE)
       VH=VT-VE-VICE
       IF (VH.LT.0.0) VH=0.
С
C... CHECK THE MIXING MASS BALANCE HERE
С
       IA=1
        WRITE (29,20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
С
       &CH1(IA), TAB, VH, TAB, CN(IA), TAB, QI(JDAY)
С
        IA=2
        WRITE (30,20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
       &CH1(IA), TAB, VH, TAB, CN(IA), TAB, QI(JDAY)
С
С
        IA=3
С
        WRITE (31, 20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
       &CH1(IA), TAB, VH, TAB, CN(IA), TAB, QI(JDAY)
С
        TA = 4
С
        WRITE (32, 20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
       &CH1(IA), TAB, VH, TAB, CN(IA), TAB, QI(JDAY)
С
        IA=5
        WRITE (33, 20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
       &CH1(IA), TAB, VH, TAB, CN(JA), TAB, QI(JDAY)
С
С
        IA=6
С
        WRITE (34,20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
       &CH1(IA), TAB, VH, TAB, CN(IA), TAB, QI(JDAY)
С
        TA=7
С
        WRITE (35,20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
       &CH1(IA), TAB, VH, TAB, CN(IA), TAB, QI(JDAY)
С
        IA=8
        WRITE (36,20) JDAY, TAB, CI1 (IA), TAB, VICE, TAB, CE1 (IA), TAB, VE, TAB,
С
С
       &CH1(IA), TAB, VH, TAB, CN(IA), TAB, QI(JDAY)
       FORMAT(I3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3
20
      &, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F25.3, A1, F25.3
      &, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3)
С
C... SET INITIAL VALUES FOR VOLUMES, SO NO ICE MELT OR THERMOCLINE MOVEMENT
C... OCCUR ON FIRST DAY
C... VICE1 IS YESTERDAY'S ICE VOLUME
C... VH1 IS YESTERDAYS HYPOLIMNION VOLUME
С
       IF (JDAY.EQ.JD1) THEN
         VICE1=VICE
         VE1=VE
         VH1=VH
       ENDIF
С
C... THE FOLLOWING CHANGES IN VOLUMES CONTROL THE LAKE LAYER MIXING
С
       VHD=VH1-VH
       DIV=VICE-VICE1
CTW
```

```
63
```

```
EVBM=VE1
      ECBM=CE1(1)
       WRITE(*,*)'VE1 BM=',VE1
С
CTW
с.
C... BEGIN ICE LAYER/EPILIMNION INTERFACE MIXING
C
      IF (DIV.LT.-0.1) THEN
C. ADD TO EP, SUBTRACT FROM ICE
        DO 8 I=1,9
          CE2(I) = (CE1(I) *VE1+CI1(I) *ABS(DIV)) / (VE1+ABS(DIV))
          CE1(I) = CE2(I)
           IF ((VICE1-ABS(DIV)).LE.O.) THEN
           CJ1(I) = 0.0
C... ABOVE WE SET CI1 TO ZERO, NO MORE ICE
          ELSE
           CI2(I) = (CI1(I) * VICE1-C11(I) * ABS(DIV)) / (VICE1-ABS(DIV))
           CI1(I) = CI2(I)
          ENDIF
        CONTINUE
8
        VE1=VE1+ABS(DIV)
      ELSEIF (DIV.GT.0.1) THEN
C. ADD TO ICE, SUBTRACT FROM EP
        DO 9 I=1,9
          CI2(I) = (CI1(I) * VICE1 + CE1(I) * ABS(DIV)) / (VICE1 + DIV)
          CI1(I)=C12(I)
          IF((VE1-DIV).LT.0.0)THEN
           CE1(I)=0.0
C... ABOVE WE LET CE1 = ZERO, LAKE MUST BE FREEZING SOLID
          ELSE
           CE2(I) = (CE1(I) *VE1-CE1(I) *ABS(DIV)) / (VE1-DIV)
           CE1(I) = CE2(I)
          ENDIF
9
        CONTINUE
        VE1=VE1-ABS(DIV)
C. NEXT LINE TO CORRECT POSSIBLE ROUND OFF
        IF (VE1.LT.0.0) VE1=0.0
      ENDIF
C. REPLACE OLD VALUES WITH NEW VALUES, ALSO NO CHANGE IN ICE LAYER
        VICE1=VICE
CTW
      EVAI=VE1
      ECAI=CE1(1)
С
       WRITE(*,*)'VE1 AI=',VE1
CTW
с.
C... END ICE
с.
C... BEGIN EPI/HYPO INTERFACE MIXING
с.
С
       WRITE(*,*)'VHD= ',VHD
      IF (VHD.GT.0.1) THEN
C. ADD TO EP, SUBTRACT FROM HY
        DO 10 I=1,9
          CE2(I) = (CE1(I) *VE1+CH1(I) *VHD) / (VE1+VHD)
          CE1(I) = CE2(I)
          IF ((VH1-VHD).LE.O.) THEN
           CH1(I) = 0.0
C... ABOVE WE LET CH1 = ZERO, TURNOVER OR TOTAL FREEZING
          ELSE
           CH2(I)=(CH1(I)*VH1-CH1(I)*VHD)/(VH1-VHD)
           CH1(I)=CH2(I)
           ENDIF
10
        CONTINUE
```
```
VE1=VE1+VHD
      ELSEIF (VHD.LT.-0.1) THEN
C. ADD TO HY, SUBTRACT FROM EP
         DO 11 I=1,9
           CH2(I) = (CH1(I) *VH1+CE1(I) *ABS(VHD)) / (VH1+ABS(VHD))
           CH1(I) = CH2(I)
           IF((VE1-ABS(VHD)).LT.0.0)THEN
            CE1(I)=0.0
C... ABOVE CASE SHOULD NEVER HAPPEN, THERMOCLINE AND ICE/EPI INTERFACE MEET
C... SCREENED OUT IN SUBROUTINE THERM
           ELSE
            CE2(I) = (CE1(I) *VE1-CE1(I) *ABS(VHD)) / (VE1-ABS(VHD))
           CE1(I) = CE2(I)
           ENDIF
11
         CONTINUE
         VE1=VE1+VHD
      ENDIF
C. REPLACE OLD VALUES WITH NEW VALUES, ALSO NO CHANGE IN THERMOCLINE CASE
         VE1=VE
         VH1=VH
C... CHECK VOLUMES
      WRITE (19, 14) JDAY, TAB, VT, TAB, VICE, TAB, VE1, TAB, VH1
C... END CHECK VOLUMES
CTW
      EVAH=VE1
       ECAH=CE1(1)
       WRITE(*,*)'VE1 AH=',VE1
С
CTW
с.
C... END EPI.HYPO INTERFACE MIXING
с.
C... DO THE MIXING OF THE INFLOW
с.
CTW
       SVBM=QI (JDAY)
       SCBM=CN(1)
CTW
       VT=VE+QI (JDAY)
       DO 100 I=1,9
C... MIXED IN uEQ/L
         CE1(I) = (QI(JDAY) * CN(I) + VE1 * CE1(I)) / (VT)
         CN(I) = CE1(I)
100 CONTINUE
С
         WRITE (29,20) JDAY, TAB, CN(1), TAB, CN(2), TAB, CN(3), TAB, CN(4)
С
С
       &, TAB, CN (5), TAB, CN (6), TAB, CN (7), TAB, CN (8), TAB, CN (9)
C... RE-EQUILIBRATE LAKE EPI LAYER
      CALL REQ(VICE)
        WRITE (29,20) JDAY, TAB, CN(1), TAB, CN(2), TAB, CN(3), TAB, CN(4)
С
С
       &, TAB, CN(5), TAB, CN(6), TAB, CN(7), TAB, CN(8), TAB, CN(9)
С
CTW
       SVAM=VT
       SCAM=CN(1)
С
С
        WRITE (29,20) JDAY, TAB, EVBM, TAB, ECBM
С
       &
                          , TAB, EVAI, TAB, ECAI
С
                          , TAB, EVAH, TAB, ECAH
       &
                          , TAB, SVBM, TAB, SCBM
С
       $
С
                          , TAB, SVAM, TAB, SCAM
       &
CTW
С
         FORMAT (I3, A1, F10.0, A1, F10.0, A1, F10.0, A1, F15.0, A1, F15.0)
14
         FORMAT(I3, 3X, 5(4X, F10.1))
15
```

```
С
C...IF LAKE OBSERVATIONS EXIST FOR THIS DAY, WRITE OUT CALC AND OBS DATA
    FOR COMPARISON (R^2, TAB DELIMITED, OUTPUT FILE)
С
С
       IF (JDAY.EQ.OLJD(ILOD)) THEN
          OLH(ILOD) = OLH(ILOD) / 1000000.
          PH=-LOG10(H/1000000.)
          OLH(ILOD) = OLH(ILOD) * 1000000.
           WRITE(6,43)JDAY, TAB, QI(JDAY), TAB, ANC, TAB, CL, TAB, H
С
          , TAB, NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, PH
C
       £
           , TAB, OLANC(ILOD), TAB, OLCL(ILOD), TAB, OLH(ILOD), TAB,
С
       8
С
           OLNH4 (ILOD), TAB, OLNO3 (ILOD), TAB, OLSBC (ILOD), TAB, OLSIO2 (ILOD)
       &
С
           , TAB, OLSO4 (ILOD), TAB, OLPH (ILOD)
       &
          ILOD=ILOD+1
      ENDIF
43
      FORMAT(I3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
      & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
      & F10.3, A1, F10.3, A1, F10, 3, A1, F10.3, A1, F10.3, A1,
      & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1,
      & F10.3, A1, F10.3, A1, F10.3, A1, F10.3, A1, F10.3)
С
        WRITE(15,*)' EPILIMNION VOLUME =', VE
С
        WRITE (15, *) ' EPILIMNION CONC. AFTER MIXING, uEQ/L'
С
С
        WRITE(15.4)
С
        WRITE(15,42)ANC,CL,DOC,H,
C
       &NH4,NO3,SBC,SIO2,SO4
       XPH=-LOG10(H/100000)
      WRITE(12,41)JDAY, TAB, QI(JDAY), TAB, ANC, TAB, CL, TAB, DOC, TAB, H, TAB,
      &NH4, TAB, NO3, TAB, SBC, TAB, SIO2, TAB, SO4, TAB, XPH
41
      FORMAT(I3, A1, F10.1, A1, F10.1, A1, F10.1, A1, F10.1, A1, F10.1,
      &A1,F10.1,A1,F10.1,A1,F10.1,A1,F10.1,A1,F10.1,A1,F10.1,
      &A1, F10.1, A1, F10.1, A1, F10.1, A1, F10.1)
42
      FORMAT (F10.1, F10.1, F10.1, F10.1, F10.1,
      &F10.1,F10.1,F10.1,F10.1,F10.1,F10.1)
      FORMAT(2X, 'ANC', 4X, 'CL', 3X, 'DOC', 5X, 'H', 5X
4
      &, 'NH4', 3X, 'NO3', 3X, 'SBC', 2X, 'SIO2', 3X, 'SO4')
3
       FORMAT(9(F5.1,1X),/)
       RETURN
       END
       SUBROUTINE THERM (JDAY, AL, VL, VICE)
C... AL = HEIGHT OF BOTTOM OF ICE FROM LAKE BOTTOM
$INSERT COMMON/THERM/
С
С
       XH IS OBS THERMOCLINE HEIGHT ON JULIAN DAY XD
С
       REAL*8 VL, AL, VICE, XH, XD
       REAL TC
С
        GIVEN DAY, RETURN VOLUME OF EPILIMNION(VL) AND ICE(VICE)
С
С
       DO 10, IC=1,14
         IF(JDAY.GT.XD(IC)) I=IC
10
       CONTINUE
        TC=TCE(I, JDAY)
C... TO IS ELEVATION OF THERMOCLINE
C... AL IS ELEVATION OF BOTTOM OF ICE LAYER
C... AL CANNOT BE LOWER THAN 'IC
        IF (AL.LT.TC) THEN
          WRITE (*, *) ' ERROR IN ICE AND THERMOCLINE ELEVATIONS !!'
          STOP
        ENDIF
```

С

```
IF(TC.LT..5)I=1
       IF(TC.GE..5) I=2
       IF(TC.GT.1.5) I=3
       IF(TC.GT.2.5)I=4
       IF(TC.GT.3.5) I=5
       IF(TC.GT.4.5) I=6
       IF(TC.GT.5.5)I=7
       IF(TC.GT.6.5) I=8
       IF(TC.GT.7.5) I=9
       IF(TC.GT.8.5) I=10
       IF(TC.GT.9.5) I=11
         VL=VOL(I,TC)
С
          WRITE(*,*)' TC ELEV = ',TC
          WRITE(*,*)' EPI VOL = ',VL
С
       TC=AL
       IF(TC.LT..5)I=1
       IF (TC.GE..5) I=2
       IF (TC.GT.1.5) I=3
       IF(TC.GT.2.5) I=4
       IF(TC.GT.3.5)I=5
       IF(TC.GT.4.5)I=6
       IF(TC.GT.5.5) I=7
       IF(TC.GT.6.5)I=8
       IF(TC.GT.7.5)I=9
       IF(TC.GT.8.5) I=10
       IF(TC.GT.9.5) I=11
         VICE=VOL(I,TC)
          WRITE(*,*)' ICE ELV = ',TC
С
          WRITE(*,*)' ICE VOL = ',VICE
С
         VL=VL-VICE
          WRITE(*,*)' NEW EPI VOL = ',VL
С
      RETURN
      END
С
с...
С
      FUNCTION TCE(I, JDAY)
$INSERT COMMON/THERM/
      REAL*8 XH, XD
C... INTERPOLATE THERMOCLINE ELEVATION BASED ON INPUT VALUES OF
C... THERMOCLINE HEIGHT AND CORRESPONDING DAYS
      XM = (XH(I+1) - XH(I)) / (XD(I+1) - XD(I))
      TCE=XH(I+1)-XM*(XD(I+1)-FLOAT(JDAY))
      RETURN
      END
С
с...
С
      FUNCTION VOL(I,TC)
      DIMENSION T(12), V(12)
      DATA T/0.,.5,1.5,2.5,3.5,4.5,5.5,6.5,7.5,8.5,9.5,10.5/
      DATA V/0.,2200.,8000.,19400.,31400.,46800.,65200.,84000.,104000.,
              130000.,151600.,179600./
     S.
C... INTERPOLATE VOLUME OF THE HYPOLIMNION FROM THE TC ELEVATION
      M = (V(I+1) - V(I)) / (T(I+1) - T(I))
      VOL=V(I+1) - M * (T(I+1) - TC)
C... EPILIMNION VOL = TOTAL VOL - HYPOLIMNION VOL
      VOL=179600-VOL
С
       WRITE(*,*)' VOL',VOL
      RETURN
      END
         SUBROUTINE IONIC
       SUBROUTINE IONIC
$INSERT COMMON/MAIN/
```

TAB=CHAR(137) C... WRITES IN A SPECIFIED FORMAT FOR MAKEING BAR GRAPHS IN CRICKET GRAPH C... A MACINTOSH APPLICATION. USEFUL FOR COMPARING IONIC STRENGTHS С Z=0. с.. IF (IFLAG.EQ.1) THEN WRITE (14, 1) TAB, JDAY, TAB, TPSBC, TAB, Z, TAB &, Z, TAB, Z, TAB, IPH, TAB, Z, TAB, Z, TAB, IPNH4 FORMAT(16HCATIONS INPUT(R),A1,I3,8(A1,F14.2)) 1 WRITE(14,3) TAB, JDAY, TAB, Z, TAB, Z, TAB &, IPCL, TAB, IPDOC, TAB, Z, TAB, IPNO3, TAB, IPSO4, TAB, Z 3 FORMAT(16H ANIONS INPUT(R), A1, I3, 8(A1, F14.2)) ELSE WRITE(14,2) TAB, JDAY, TAB, IPSBC, TAB, Z, TAB &,Z,TAB,Z,TAB,IPH,TAB,Z,TAB,Z,TAB,IPNH4 2 FORMAT(16HCATIONS INPUT(S) ,A1,I3,8(A1,F14.2)) WRITE (14, 4) TAB, JDAY, TAB, Z, TAB, Z, TAB &, IPCL, TAB, IPDOC, TAB, Z, TAB, IPNO3, TAB, IPSO4, TAB, Z FORMAT(16H ANIONS INPUT(S) ,A1,I3,8(A1,F14.2)) 4 ENDIF с... WRITE (14, 5) TAB, JDAY, TAB, SBC, TAB, Z, TAB &, Z, TAB, Z, TAB, H, TAB, Z, TAB, Z, TAB, NH4 5 FORMAT(12HCATIONS CALC , A1, I3, 8(A1, F14.2)) WRITE (14, 6) TAB, JDAY, TAB, Z, TAB, ANC, TAB &, CL, TAB, DOC, TAB, Z, TAB, NO3, TAB, SO4, TAB, Z 6 FORMAT(11HANIONS CALC, A1, I3, 8(A1, F14.2)) с... WRITE (*,*) 'JDAY', JDAY, ' OSJD(',10,') ',OSJD(10) С IF (JDAY.EQ.OSJD(IO)) THEN WRITE (14,7) TAB, OSJD (10), TAB, OSSBC (IO), TAB, Z, TAB &, Z, TAB, Z, TAB, OSH (IO), TAB, Z, TAB, Z, TAB, OSNH4 (IO) -7 FORMAT (11HCATIONS OBS, A1, I3, 8(A1, F14.2)) WRITE(14,8)TAB,OSJD(IO),TAB,Z,TAB,OSANC(IO),TAB &, OSCL(IO), TAB, OSDOC(IO), TAB, Z, TAB, OSNO3(IO), TAB &,OSSO4(IO),TAB,Z IO==IO+1 ENDIF WRITE (14, *) TAB FORMAT(11H ANIONS OBS, A1, I3, 8(A1, F14.2)) 8 RETURN END



101 ARB/R-90/446 2. 3. R ARE PROCLIMENTATION FROME 1. REPORT NO. ARB/R-90/446 2. 3. R Statisting the Response of Emerald Lake, an Alpine Watershed in Sequial Mational Park, California, to Acidification During Snow- melt by Using a Simple Hyrochemical Model 6. Statisting the Response and Address The Companization Name and Address Statistics Name and Address California Air Resources Exard 10. Statistics Name and Address California Air Resources Exard 11. Statistics Name and Address California Air Resources Exard 11. Statistics Name and Address California Air Resources Exard 12. Statistics Name and Address California Air Resources Exard 13. Statistics Name and Address California Air Resources Exard 13. Statistics Name and Address California Air Resources Exard 14. Statistics Name and A	ہ مسیر یہ ہے م	
101 ARB/R=90/446 2 A. R. PAGE ARB/R=90/446 PE Streaming the Response of Emerald Lake, an Alpine Watershed in Alug Arg. Sequoia National Park, California, to Acidification During Snow- 6 ent by Using a Simple Hyrochemical Model 8. P Contraction Name and Address 10. The Boological Survey 11. Science Commission Name and Address 10. Science Commission Name and Address 10. Science Commission Name and Address 10. Science Commission Name and Address 11. S		
RT: DOCUMENTATION 1. REPORT NO. PAGE PE AUGE ALRB/R-90/446 F. Sequeia National Park, California, to Acidification During Snow- neit by Using a Simple Hyrochemical Model F. Stickard P. Hooper, Christopher T. West, and Norman E. Peters 10. String Commission Name and Address 11. Research Division 11. 1800 15th Street 14. Sagramento, CA 95814 14. preference a simple, process-oriented model, called the Alpine Lake Forecast 13. collected during the Integrated Watershed Study at Emerald Lake, Sequoia Nationa 14. preference a loss of ANC and depression in stream water during snownel twee ground water is a minor contributor to stream flow. It includes an empiring rimary mineral wathering as the only alkalinity-generating mechanism. Hydrolo heavy snow year were used to calibrate the model. Matershed processes and in watershed in order to improve the performance of the model. We plan to apply the in the Siterra studied as part of the California Air Resources Board	ecipient's Accession I	No.
Statistics Alg. Security the Response of Emerald Lake, an Alpine Watershed in Security National Park, California, to Acidification During Snow- Alg. Security Simple Hyrochemical Model Statistics Simple Hyrochemical Model 8. P Statistics New York Statistics 10. Statistics New York Statistics 10. Statistics 11. 11. Statistis 12. 12.	390266511	
Sequeia National Park, California, to Acidit for During Show- e. Ent by Using a Simple Hyrochemical Model Sichard P. Hooper, Christopher T. West, and Norman E. Peters The Constructed Process and Address The Constructed a simple Hyrochemical Model We constructed a simple, process-oriented model, called the Alpine Lake Forecast Collected during the Integrated Watershed Study at Emeral Lake, Sequeia National Signard Address Signard Address Collected during the Integrated Watershed Study at Emeral Lake, Sequeia National Signard Address Iter a by Using a Simple Hyrochemical Model We constructed a simple, process-oriented model, called the Alpine Lake Forecast Collected during the Integrated Watershed Study at Emeral Lake, Sequeia Nationa designed to capture the basic solute patterns in stream water during snownelt where ground water is a minor contributor to stream flow. It includes an empiri primary mineral weathering as the only alkalinity-generating mechanism. Hydrolo heavy snow year were used to calibrate the model. Watershed processes during a be different.from the calibration year. The model forecast concludes that stree likely to experience a loss of AVK and depression in pH during spring rain storm snownelt dilution phase. Additional data are needed on snownelt.processes and i watershed in order to improve the performance of the model. We plan to apply th in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Xecument Analysis a Descriptors lakes, models, acidification, hydrology b. Kentifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequeia National f Unclassified	ust 1990	
Sichard P. Hooper, Christopher T. West, and Norman E. Peters 8. F Sichard P. Hooper, Christopher T. West, and Norman E. Peters 10. Sichard P. Hooper, Christopher T. West, and Norman E. Peters 10. Sichard P. Hooper, Christopher T. West, and Norman E. Peters 10. Sichard P. Hooper, Christopher T. West, and Norman E. Peters 10. Sichard P. Hooper, Christopher T. West, and Norman E. Peters 10. Sichard P. Hooper, Christopher T. West, and Norman E. Peters 10. Sichard P. Hooper, Christopher T. West, and Norman E. Peters 10. We constructed Integrated Naters and Address California Air Resources Board Research Division 13. B00 J5th Street 14. Sacramento, CA S5814 14. Sacramento, CA S5814 14. Sacramento, CA, S5814 14. Sacramento, CA, S5814 16. Sierra Nevada, CA. 15. ALF is able to capture the basic solute patterns in stream water during snownelt where ground water is a minor contributor to stream flow. It includes an empiring minary mineral weathering as the only alkalinity-generating mechanism. Hydrolog the performance of the model. Waring spring rain storm snownelt dilution phase. Additional data are needed on snownelt. processes and in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Secure Analysis		
Noted F. Rodes, Wasterney 10. The Gongitzetion Name and Address 10. Water Resources Investigations 11. Water Resources Investigations 12. Water Resources Investigations 12. W	erforming Organizatio	on Rept. No.
J.S. Geological Survey water Resources Investigations 11. (C)	Project/Task/Werk Un	nit No.
(c) (Contract(C) or Grant(G) No.
(c) (A732-034	E Constantino de Const
Program Organization Name and Address 13. California Air Resources Board 14. Reseach Division 14. Sacramento, CA 95814 14. upplementary Notes 14. wateract (Limits 200 words) 14. wateract during the Integrated Watershed Study at Emerald Lake, Sequoia National designed to answer questions concerning the impact of acid deposition on high-ell Sierra Nevada, CA. ALF is able to capture the basic solute patterns in stream water during snownelt where ground water is a minor contributor to stream flow. It includes an empirin primary mineral weathering as the only alkalinity-generating mechanism. Hydrolc heavy snow year were used to calibrate the model. Watershed processes during a be different from the calibration year. The model forecast concludes that stream is snownelt dilution phase. Additional data are needed on snownelt. We plan to apply the mechanism. Hydrolc heavy snow gear were used to calibrate the Ealifornia Air Resources Board's Atmospheri Program. Xecument Analysis a. Descriptore lakes, models, acidification, hydrology s. Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F walability Statement lase Unimited, Available from National Technical 19. Security Class (This R		
Alignmenta Air Resources Board Reseach Division 1800_15th, Street Sacramento, CA 95814 Implementary Notes Automatic Advances	Type of Report & Per	riod Covered
Research Division 14. Sacramento, CA 95814 14. ipplementary Notes 14. we constructed a simple, process-oriented model, called the Alpine Lake Forecast collected during the Integrated Watershed Study at Emerald Lake, Sequeia Nationa designed to answer questions concerning the impact of acid deposition on high-el Sierra Nevada, CA. ALF is able to capture the basic solute patterns in stream water during snownelt where ground water is a minor contributor to stream flow. It includes an empiring rimary mineral weathering as the only alkalinity-generating mechanism. Hydrolc heavy snow year were used to calibrate the model. Watershed processes during a be different from the calibration year. The model Torecast concludes that stream snownelt dilution phase. Additional data are needed on snownelt processes and in watershed in order to improve the performance of the model. We plan to apply the in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Socument Analysis . Descriptors Takes, models, acidification, hydrology >. Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F Unclassified	Final Parant	
Incomplete Section Incomplete Section Secramento, CA 95814 Incomplete Section Septementary Notes Incomplete Section Setrect (Limits 200 words) Incomplete Section Section (Limits 200 words) Section (Limits 200 words) Section (Limits 200 words)	Final Report	
We constructed a simple, process-oriented model, called the Alpine Lake Forecast collected during the Integrated Watershed Study at Emerald Lake, Sequoia National designed to answer questions concerning the impact of acid deposition on high-el Sierra Nevada, CA. ALF is able to capture the basic solute patterns in stream water during snownelt where ground water is a minor contributor to stream flow. It includes an empiri primary mineral weathering as the only alkalinity-generating mechanism. Hydrolc heavy snow year were used to calibrate the model. Watershed processes during a be different from the calibration year. The model forecast concludes that stree likely to experience a loss of ANC and depression in pH during spring rain storm snownelt dilution phase. Additional data are needed on snownelt. We plan to apply th in the Sierra studied as part of the California Air Resources. Board's Atmospheri Program. Xecument Analysis a. Descriptors lakes, models, acidification, hydrology . Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F unclassified 19. Security Class (This R Unclassified	t itr	900 - 5 00 0 1 3111
We constructed a simple, process-oriented model, called the Alpine Lake Forecast collected during the Integrated Watershed Study at Emerald Lake, Sequoia Nationa designed to answer questions concerning the impact of acid deposition on high-el Sierra Nevada, CA. ALF is able to capture the basic solute patterns in stream water during snowmelt where ground water is a minor contributor to stream flow. It includes an empiri- primary mineral weathering as the only alkalinity-generating mechanism. Hydrolo heavy snow year were used to calibrate the model. Watershed processes during a be different.from the calibration year. The model forecast concludes that streat likely to experience a loss of ANC and depression in pH during spring rain storm snowmelt dilution phase. Additional data are needed on snowmelt.we plan to apply the in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Xecument Analysis a. Descriptors lakes, models, acidification, hydrology . Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F usates the limited. Available from National Technical 19. Security Class (this R Unclassified		
We constructed a simple, process-oriented model, called the Alpine Lake Forecast collected during the Integrated Watershed Study at Emerald Lake, Sequoia Nationa designed to answer questions concerning the impact of acid deposition on high-el Sierra Nevada, CA. ALF is able to capture the basic solute patterns in stream water during snowmelt where ground water is a minor contributor to stream flow. It includes an empiri primary mineral weathering as the only alkalinity-generating mechanism. Hydrolo heavy snow year were used to calibrate the model. Watershed processes during a be different from the calibration year. The model forecast concludes that stree likely to experience a loss of ANC and depression in pH during spring rain storm snowmelt dilution phase. Additional data are needed on snowmelts processes and in watershed in order to improve the performance of the model. We plan to apply th in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Jocument Analysis a Descriptors lakes, models, acidification, hydrology > Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F Unclassified	م د الدي الم الح الم الح	
We constructed a simple, process-oriented model, called the Alpine Lake Forecast collected during the Integrated Watershed Study at Emerald Lake, Sequoia Nationa designed to answer questions concerning the impact of acid deposition on high-el Sierra Nevada, CA. ALF is able to capture the basic solute patterns in stream water during snowmelt where ground water is a minor contributor to stream flow. It includes an empiri- primary mineral weathering as the only alkalinity-generating mechanism. Hydrolc heavy snow year were used to calibrate the model. Watershed processes during a be different from the calibration year. The model forecast concludes that streed likely to experience a loss of ANC and depression in pH during spring rain storm snowmelt dilution phase. Additional data are needed on snowmelt processes and in watershed in order to improve the performance of the model. We plan to apply th in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Socument Analysis a Descriptors lakes, models, acidification, hydrology . Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F Unclassified 19. Security Class (This R Unclassified		
We constructed a simple, process-oriented model, called the Alpine Lake Forecast collected during the Integrated Watershed Study at Emerald Lake, Sequoia Nationa designed to answer questions concerning the impact of acid deposition on high-el Sierra Nevada, CA. ALF is able to capture the basic solute patterns in stream water during snowmelt where ground water is a minor contributor to stream flow. It includes an empiri primary mineral weathering as the only alkalinity-generating mechanism. Hydrolc heavy snow year were used to calibrate the model. Watershed processes during a be different from the calibration year. The model forecast concludes that stree likely to experience a loss of ANC and depression in pH during spring rain storm snowmelt dilution phase. Additional data are needed on snowmelt processes and i watershed in order to improve the performance of the model. We plan to apply th in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Xecument Analysis • Descriptors lakes, models, acidification, hydrology • Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F Unclassified * COSATI Field/Group	ينيس جين ۽ پ	a ana ana ana ana ana ana ana ana ana a
be different from the calibration year. The model forecast concludes that stree likely to experience a loss of ANC and depression in pH during spring rain storm snowmelt dilution phase. Additional data are needed on snownelt processes and i watershed in order to improve the performance of the model. We plan to apply th in the Sierra studied as part of the California Air Resources Board's Atmospheri Program. Document Analysis a Descriptors lakes, models, acidification, hydrology >. Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F : COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical 19. Security Class (This R Unclassified	ical representat agic and chemica light snow year	ion of 11 data from a appeared to
Document Analysis a. Descriptors lakes, models, acidification, hydrology b. Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National H c. COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical	ion-exchange in ijs model at oth ic Acidity Prote	ring the soils of this yer, watersheds
lakes, models, acidification, hydrology b. Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F c. COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical		
lakes, models, acidification, hydrology b. Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National K c. COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical		
 Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National F COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical 19. Security Class (This R	••••	• • • • • •
 Identifiers/Open-Ended Terms Kapiloff Acid Deposition Program; Integrated Watershed Study; Sequoia National I COSATI Field/Group Availability Statement Lease Unlimited. Available from National Technical 19. Security Class (This R		
COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical	Park - Emerald L	Lake
c. COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical Unclassified	·· ····	
c. COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical Unclassified	: میست. اینان	
c. COSATI Field/Group Availability Statement lease Unlimited. Available from National Technical Unclassified		1997 - 1997 -
Availability Statement lease Unlimited. Available from National Technical Unclassified	(eport) 21. N	to. of Pages
	2014 2014	68
formation Service. 5285 Port Royal Rd. Springfield, VA 20. Security Class (This P	'age)	hice
	OPTIO	NAL FORM 272 (4-

.

.

×