

APPENDIX D

ATMOSPHERIC LIFETIME AND FATE OF EPOXYETHANE

A Report Submitted in Partial Fulfillment of  
California Air Resources Board Contract No. A732-107

"Lifetimes and Fates of Toxic Air Contaminants in  
California's Atmosphere"

Roger Atkinson

August 1989

Statewide Air Pollution Research Center  
University of California  
Riverside, CA 92521



## TABLE OF CONTENTS

	<u>Page</u>
A. Introduction.....	D-3
B. Chemical Loss Processes.....	D-5
1. Ozone Reaction.....	D-5
2. OH Radical Reaction.....	D-5
3. NO <sub>3</sub> Radical Reaction.....	D-7
4. Photolysis.....	D-8
C. Physical Loss Processes.....	D-8
D. Atmospheric Formation Processes.....	D-9
E. Atmospheric Lifetimes.....	D-10
F. Ambient Concentration.....	D-11
G. References.....	D-11



#### A. Introduction

The atmospheric fates of chemical compounds emitted into the troposphere are governed by a number of chemical and physical removal processes. As the result of laboratory studies carried out over the past 20 years, the potentially important gas-phase chemical reaction pathways which must be considered includes:

- Photolysis during daylight hours
- Reaction with the hydroxyl (OH) radical during daylight hours
- Reaction with the hydroperoxyl ( $\text{HO}_2$ ) radical, mainly during afternoon/evening hours
- Reaction with the nitrate ( $\text{NO}_3$ ) radical during nighttime hours
- Reaction with nitrogen dioxide ( $\text{NO}_2$ )
- Reaction with ozone ( $\text{O}_3$ )
- Reaction with gaseous nitric acid ( $\text{HNO}_3$ )

For the great majority of organic compounds, the most important of these remain photolysis and reaction with hydroxyl (OH) and nitrate ( $\text{NO}_3$ ) radicals and with ozone ( $\text{O}_3$ ).

However, for a limited number of organic compounds, one or more of the other reactive chemical species in the troposphere may react with organic compounds at significant rates. For example,  $\text{HO}_2$  radicals react with formaldehyde and, more slowly, with acetaldehyde and glyoxal;  $\text{NO}_2$  reacts with dialkenes; and gaseous  $\text{HNO}_3$  reacts with basic compounds such as the amines.

Additionally, for chemical compounds present in the adsorbed phase, photolysis and reaction with  $\text{O}_3$ ,  $\text{N}_2\text{O}_5$ ,  $\text{NO}_2$ ,  $\text{HNO}_3$ ,  $\text{HONO}$ ,  $\text{H}_2\text{SO}_4$  and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) may contribute to their degradation. For certain of these reactions involving adsorbed organics, synergistic interactions may be of importance.

The physical removal processes can be defined in a simplistic manner as accretion (or coagulation) of particles, and dry and wet deposition of gases and particles. Removal of gases and particles at ground surfaces (including snow and other moist surfaces) is referred to as dry deposition, while removal of these species by precipitation is referred to as wet deposition. Our current understanding of these processes, which are

dynamic in nature, must be considered to be semi-quantitative at the present time, and in general there is a lack of experimental data concerning these processes for most organic compounds (Atkinson, 1988; Bidleman, 1988).

The atmospheric lifetime  $\tau$  of a chemical is defined as the time to decay to a concentration of  $1/e$  ( $=0.368$ ) of that initially present. Also,

$$\frac{1}{\tau} = \frac{1}{\tau_{\text{chemical}}} + \frac{1}{\tau_{\text{physical}}} \quad (1)$$

where  $\tau_{\text{chemical}}$  and  $\tau_{\text{physical}}$  are the lifetimes of the chemical with respect to only chemical reaction and only physical loss processes, respectively. These chemical and physical lifetimes are composites of those for the individual loss processes, for example

$$\frac{1}{\tau_{\text{chemical}}} = \frac{1}{\tau_{\text{photolysis}}} + \frac{1}{\tau_{\text{OH}}} + \frac{1}{\tau_{\text{NO}_3}} + \frac{1}{\tau_{\text{O}_3}} \dots \quad (2)$$

where  $\tau_{\text{photolysis}}$ ,  $\tau_{\text{OH}}$ ,  $\tau_{\text{NO}_3}$  and  $\tau_{\text{O}_3}$  are the lifetimes with respect to photolysis, reaction with the OH radical, reaction with the  $\text{NO}_3$  radical, and reaction with  $\text{O}_3$ , respectively. In turn, these reactive loss processes are determined by the rate constants,  $k_x$ , for reaction and the ambient atmospheric concentrations,  $[X]$ , of the reactive intermediates. For example

$$\frac{1}{\tau_{\text{OH}}} = k_{\text{OH}}[\text{OH}]$$

For photolysis, the photolysis rate depends on the absorption cross section ( $\sigma$ ), the photolysis quantum yield ( $\phi$ ), and the radiation intensity ( $J$ ), all of which are wavelength dependent:

$$\frac{1}{\tau_{\text{photolysis}}} = k_{\text{photolysis}} = \int_{\sim 290 \text{ nm}}^{\sim 800 \text{ nm}} \sigma_{\lambda} \phi_{\lambda} J_{\lambda} d\lambda$$

In this report, we describe the available literature data concerning the chemical and physical removal processes for epoxyethane ( $\text{CH}_2-\text{CH}_2$ ) and estimate its resulting atmospheric lifetime. We will also summarize the atmospheric chemistry of this compound. Since epoxyethane is gaseous under atmospheric conditions, we consider only the gas-phase processes.

## B. Chemical Loss Processes

### 1. Ozone Reaction

No experimental data have been reported concerning the gas-phase reaction of  $\text{O}_3$  with epoxyethane (Atkinson and Carter, 1984). However, based upon the magnitude of the rate constant for the gas-phase reaction of the OH radical with epoxyethane and the literature data for the reactions of  $\text{O}_3$  with saturated organic compounds such as the alkanes and carbonyls not containing  $\text{>C=C<}$  bonds (Atkinson and Carter, 1984), it is expected that the room temperature rate constant for the reaction of  $\text{O}_3$  with epoxyethane will be

$$k(\text{O}_3 + \text{epoxyethane}) < 10^{-20} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

at 298 K.

### 2. OH Radical Reaction

The available kinetic data are given in Table 1. At room temperature, the absolute rate constants of Zetzsch (1980), Lorenz and Zellner (1984) and Wallington et al. (1988) exhibit a spread of a factor of 1.8, but are consistent with the upper limit to the rate constant derived by Klöpffer et al. (1986). The only temperature-dependent study is that of Lorenz and Zellner (1984), who observed a rapid increase in the rate constant above 435 K, leading to marked non-Arrhenius behavior. Atkinson (1989) used a unit-weighted average of the room temperature rate constants of Zetzsch (1980), Lorenz and Zellner (1984) and Wallington et al. (1988) to recommend

$$k(\text{OH} + \text{epoxyethane}) = 7.6 \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } 298 \text{ K,}$$

with an estimated overall uncertainty of  $\pm 50\%$ .

Table 1. Rate Constants k and Temperature-Dependent Parameters,  $k = Ae^{-B/T}$ , for the Gas-Phase Reaction of the OH Radical with Epoxyethane (from Atkinson, 1989)


$10^{12} \times A$ ( $\text{cm}^3 \text{mole}^{-1} \text{s}^{-1}$ )	B (K)	$10^{12} \times k_1$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{s}^{-1}$ )	at T (K)	Technique <sup>a</sup>	Reference
$11 \pm 4$	$1460 \pm 150$ (297-435 K)	$0.053 \pm 0.01$	295	FP-RF	Zetzsch (1980)
			297	LP-RF	Lorenz and Zellner (1984)
			377		
		$0.080 \pm 0.016$ $0.18 \pm 0.04$ $0.40 \pm 0.08$	435		
			501		
			515		
		$1.6 \pm 0.1$ $2.7 \pm 0.5$			
		$< 0.10$	300	RR [relative to k(propane), $\bar{1.17} \times 10^{-12}$ ] <sup>b</sup>	Klopffer et al. (1986)
		$0.095 \pm 0.005$	296	FP-RF	Wallington et al. (1988)

<sup>a</sup>LP  $\equiv$  laser photolysis; FP  $\equiv$  flash photolysis; RF  $\equiv$  resonance fluorescence; RR  $\equiv$  relative rate.  
<sup>b</sup>Recommendation of Atkinson (1989).




Because of the high magnitude of the temperature dependence measured by Lorenz and Zellner (1984), leading to an Arrhenius pre-exponential factor of  $1.2 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  for rate data obtained over the range 297-435 K and a markedly higher value for data obtained over the temperature range 435-515 K, no temperature dependence was recommended by Atkinson (1989).


This OH radical reaction almost certainly proceeds by H atom abstraction from the C-H bonds



The initially formed radical is expected to rapidly undergo ring cleavage



Using laser induced fluorescence detection to monitor the vinoxy ( $\text{CH}_2\text{CHO}$ ) radical, Lorenz and Zellner (1984) measured  $\text{CH}_2\text{CHO}$  yields at 298 K of  $0.08 \pm 0.03$  and  $0.23 \pm 0.08$  at 10 and 60 Torr total pressure of helium diluent, respectively. The reaction products of the OH radical reaction with epoxyethane under atmospheric conditions are not presently known. [The vinoxy radical reacts rapidly with  $\text{O}_2$ , with a room temperature rate constant of  $2.6 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , with the reaction products possibly being (Gutman and Nelson, 1983; Lorenz et al., 1985)



### 3. $\text{NO}_3$ Radical Reaction

No data are currently available concerning the gas-phase reaction of the  $\text{NO}_3$  radical with epoxyethane. Based on the kinetic data for the corresponding OH radical reaction (see above) and for the gas-phase reactions of the  $\text{NO}_3$  radical with alkanes, ethers and alcohols (Wallington et

al., 1986a,b, 1987; Atkinson et al., 1988), it is expected that the room temperature rate constant for the gas-phase reaction of the NO<sub>3</sub> radical with epoxyethane will be

$$k_3(\text{NO}_3 + \text{epoxyethane}) < 5 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

at 298 K.

#### 4. Photolysis

Epoxyethane will not undergo photolysis in the actinic region, since epoxyethane does not absorb radiation above 290 nm. The long wavelength limit to the absorption is reported to be at 212 nm (Calvert and Pitts, 1966).

#### C. Physical Loss Processes

There are no published literature data available concerning dry deposition of epoxyethane. Under California Air Resources Board funding, the Statewide Air Pollution Research Center at the University of California, Riverside, carried out a series of experiments to determine the loss rates of epoxyethane as a function of the water vapor concentration in a 5800 liter Teflon-coated chamber (Winer et al., 1987). Epoxyethane and water vapor concentrations were monitored by in situ long pathlength Fourier transform infrared (FT-IR) absorption spectroscopy. The initial epoxyethane concentrations were  $\sim 1.4 \times 10^{14}$  molecule cm<sup>-3</sup> ( $\sim 6$  parts-per-million mixing ratio), and all experiments were carried out at  $298 \pm 2$  K.

The experimental conditions and observed loss rates of epoxyethane are given in Table 2. Since the experiment carried out at the lowest relative humidity exhibited the highest epoxyethane loss rate, these data presented in Table 2 show that under these experimental conditions there was no observable hydrolysis of epoxyethane (the small losses of epoxyethane were attributed to a slow removal at the chamber walls). These data yield an upper limit to the loss rate for epoxyethane at  $\sim 50\%$  relative humidity and room temperature of  $\leq 4 \times 10^{-7} \text{ s}^{-1}$ , corresponding to a minimum lifetime of epoxyethane due to hydrolysis of 29 days, with the likelihood of a much longer lifetime due to this removal process.

Table 2. Experimental Conditions and Results for the Behavior of Epoxyethane in the Presence of Water Vapor

Diluent Gas	Relative Humidity (%)	Observed Epoxyethane Loss	Loss Rate ( $s^{-1}$ )
N <sub>2</sub>	<1	1.5% over 4 hrs	$\leq 1 \times 10^{-6}$
N <sub>2</sub>	49	<0.5% over 2 hrs	$< 7 \times 10^{-7}$
Air	49	4% over 29 hrs	$\leq 4 \times 10^{-7}$

Washout ratios, W, where

$$W = \text{Concentration}_{\text{rain}} / \text{Concentration}_{\text{air}}$$

have been determined for epoxyethane at 278 K and 288 K (Dana et al., 1985), with values of  $W = 3.8$  at 278 K and  $6.2$  at 288 K being measured. When these values of  $W$  for epoxyethane are compared to the washout ratios of  $W = 10^4 - 10^6$  for species which are very efficiently scavenged by rainwater, such as phenol and particles (Eisenreich et al., 1981; Atkinson, 1988), it is clear that epoxyethane is inefficiently removed from the troposphere by wet deposition. Indeed, the values of  $W$  for epoxyethane are very similar to those for methylchloroform ( $\text{CH}_3\text{CCl}_3$ ) [ $W \sim 5$  at 298 K (Dana et al., 1985)], for which wet deposition plays no role in its tropospheric removal [the methylchloroform lifetime in the troposphere is 6-7 years, and is due to reaction with the OH radical (Prinn et al., 1987)]. Thus, it is expected that wet deposition will not lead to a significant tropospheric loss rate of epoxyethane.

#### D. Atmospheric Formation Processes

No chemical formation processes for epoxyethane in the troposphere are expected with the current knowledge of atmospheric chemistry of organic compounds (see, for example, Atkinson and Lloyd, 1984). Two possible reactions leading to formation of epoxyethane in the troposphere are the gas-phase reactions of ethene with the  $\text{O}(^3\text{P})$  atom and  $\text{O}_3$ , and at

atmospheric pressure and room temperature and below neither of these reactions leads to the formation of epoxyethane (Atkinson and Lloyd, 1984).

#### E. Atmospheric Lifetimes

The rate constants, or upper limits thereof, cited above for the gas-phase reactions of OH and NO<sub>3</sub> radicals and O<sub>3</sub> with epoxyethane can be combined with measured or estimated ambient atmospheric concentrations of OH and NO<sub>3</sub> radicals and O<sub>3</sub> to calculate the atmospheric lifetimes of epoxyethane with respect to these potential gas-phase reactions. For this purpose, we use ambient atmospheric concentrations of  $1.5 \times 10^6$  molecule cm<sup>-3</sup> for OH radicals during a 12-hr daytime period (Prinn et al., 1987) (an average tropospheric concentration for the northern and southern hemispheres derived from a knowledge of the total worldwide emissions of CH<sub>3</sub>CCl<sub>3</sub> and the present atmospheric burden of CH<sub>3</sub>CCl<sub>3</sub>),  $2.4 \times 10^8$  molecule cm<sup>-3</sup> for NO<sub>3</sub> radicals during a 12-hr nighttime period (Platt et al., 1984; Atkinson et al., 1986) and  $7 \times 10^{11}$  molecule cm<sup>-3</sup> for O<sub>3</sub> during a complete 24-hr day (Logan, 1985). The calculated lifetimes due to these gas-phase reactions are given in Table 3.

Table 3. Calculated Atmospheric Lifetimes of Epoxyethane with Respect to Gas-Phase Room Temperature Reaction with OH and NO<sub>3</sub> Radicals and O<sub>3</sub>

Reaction with	Lifetime $\tau$
OH <sup>a</sup>	200 days
NO <sub>3</sub> <sup>b</sup>	>5 years
O <sub>3</sub> <sup>c</sup>	4.5 years

<sup>a</sup>For a 12-hr average daytime OH radical concentration of  $1.5 \times 10^6$  molecule cm<sup>-3</sup> (Prinn et al., 1987).

<sup>b</sup>For a 12-hr average nighttime NO<sub>3</sub> radical concentration of  $2.4 \times 10^8$  molecule cm<sup>-3</sup> (Platt et al., 1984; Atkinson et al., 1986).

<sup>c</sup>For a 24-hr average O<sub>3</sub> concentration of  $7 \times 10^{11}$  molecule cm<sup>-3</sup> [30 ppb] (Logan, 1985).

Clearly, the dominant gas-phase chemical removal process is the reaction with the OH radical, leading to a calculated lifetime of ~200 days. This calculation uses a room temperature rate constant for the OH radical reaction; use of a rate constant for the OH radical reaction at the temperatures applicable to the mid-troposphere (~270 K) would lead to a longer calculated lifetime of ~300 days. With a lifetime of epoxyethane of this magnitude, wet deposition is expected to be of no significance as a tropospheric removal process, and the overall lifetime of epoxyethane is calculated to be in the range 200-300 days. With a lifetime of this magnitude, epoxyethane will be distributed over a global scale.

F. Ambient Concentration

To our knowledge, no ambient air data have been reported in the literature for epoxyethane.

G. References

- Atkinson, R. and W. P. L. Carter (1984) Kinetics and mechanisms of the gas-phase reactions of ozone with organic compounds under atmospheric conditions. *Chem. Rev.*, 84, 437-470.
- Atkinson, R. and A. C. Lloyd (1984) Evaluation of kinetic and mechanistic data for modeling of photochemical smog. *J. Phys. Chem. Ref. Data*, 13, 315-444.
- Atkinson, R., A. M. Winer and J. N. Pitts, Jr. (1986) Estimation of night-time  $N_2O_5$  concentrations from ambient  $NO_2$  and  $NO_3$  radical concentrations and the role of  $N_2O_5$  in night-time chemistry. *Atmos. Environ.*, 20, 331-339.
- Atkinson, R. (1988) Atmospheric transformations of automotive emissions. In "Air Pollution, the Automobile and Public Health," Eds. A. Y. Watson, R. R. Bates and D. Kennedy, National Academy Press, Washington, D.C., pp. 99-132.
- Atkinson, R., S. M. Aschmann and J. N. Pitts, Jr. (1988) Rate constants for the gas-phase reactions of the  $NO_2$  radical with a series of organic compounds at  $296 \pm 2$  K. *J. Phys. Chem.*, 92, 3454-3457.
- Atkinson, R. (1989) Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. *J. Phys. Chem. Ref. Data*, Monograph 1, 1-246.
- Bidleman, T. F. (1988) Atmospheric processes. *Environ. Sci. Technol.*, 22, 361-367.

- Calvert, J. G. and J. N. Pitts, Jr. (1966) Photochemistry. Wiley, New York, NY.
- Dana, M. T., R. N. Lee and J. M. Hales (1985) Hazardous Air Pollutants: Wet removal rates and mechanisms. EPA-600/3-84-113, Research Triangle Park, NC, January.
- Eisenreich, S. J., B. B. Looney and J. D. Thornton (1981) Airborne organic contaminants in the Great Lakes ecosystem. Environ. Sci. Technol., 15, 30-38.
- Gutman, D. and H. H. Nelson (1983) Gas-phase reactions of the vinoxy radical with  $O_2$  and NO. J. Phys. Chem., 87, 3902-3905.
- Klöpffer, W., R. Frank, E.-G. Kohl and F. Haag (1986) Quantitative Erfassung der photochemischen transformationprozesse in der troposphäre. Chemiker-Zeitung, 110, 57-61; Methods of the Ecotoxicological evaluation of chemicals. Photochemical degradation in the gas phase. Volume 6: OH reaction rate constants and tropospheric lifetimes of selected environmental chemicals. Report 1980-1983. Eds. K. H. Becker, H. M. Biehl, P. Bruckmann, E. H. Fink, F. Führ, W. Klöpffer, R. Zellner and C. Zetzsch, Kernforschungsanlage, Jülich GmbH, November 1984.
- Logan, J. A. (1985) Tropospheric ozone: Seasonal behavior, trends, and anthropogenic influence. J. Geophys. Res., 90, 10463-10482.
- Lorenz, K. and R. Zellner (1984) Rate constants and vinoxy product yield in the reaction OH + ethylene oxide. Ber. Bunsenges. Phys. Chem., 88, 1228-1231; Methods of the Ecotoxicological Evaluation of Chemicals: Photochemical Degradation in the Gas Phase. Volume 6: OH reaction rate constants and tropospheric lifetimes of selected environmental chemicals. Report 1980-1983. Eds. K. H. Becker, H. M. Biehl, P. Bruckmann, E. H. Fink, F. Führ, W. Klöpffer, R. Zellner and C. Zetzsch, Kernforschungsanlage, Jülich GmbH, November, 1984.
- Lorenz, K., D. Rhäsa, R. Zellner and B. Fritz (1985) Laser-photolysis - LIF kinetic studies of the reactions of  $CH_3O$  and  $CH_2CHO$  with  $O_2$  between 300 and 500 K. Ber. Bunsenges. Phys. Chem., 89, 341-342.
- Platt, U. F., A. M. Winer, H. W. Biermann, R. Atkinson and J. N. Pitts, Jr. (1984) Measurement of nitrate radical concentrations in continental air. Environ. Sci. Technol., 18, 365-369.
- Prinn, R., D. Cunnold, R. Rasmussen, P. Simmonds, F. Alyea, A. Crawford, P. Fraser and R. Rosen (1987) Atmospheric trends in methylchloroform and the global average for the hydroxyl radical. Science, 238, 945-950.
- Wallington, T. J., R. Atkinson, A. M. Winer and J. N. Pitts, Jr. (1986a) Absolute rate constants for the gas-phase reactions of the  $NO_2$  radical with  $CH_3SCH_3$ ,  $NO_2$ ,  $CO_2$ , and a series of alkanes at  $298 \pm 2$  K. J. Phys. Chem., 90, 4640-4644.

- Wallington, T. J., R. Atkinson, A. M. Winer and J. N. Pitts, Jr. (1986b) Absolute rate constants for the gas-phase reactions of the  $\text{NO}_3$  radical with  $\text{CH}_3\text{SH}$ ,  $\text{CH}_3\text{SCH}_3$ ,  $\text{CH}_3\text{SSCH}_3$ ,  $\text{H}_2\text{S}$ ,  $\text{SO}_2$ , and  $\text{CH}_3\text{OCH}_3$  over the temperature range 280-350 K. *J. Phys. Chem.*, 90, 5393-5396.
- Wallington, T. J., R. Atkinson, A. M. Winer and J. N. Pitts, Jr. (1987) A study of the reaction  $\text{NO}_3 + \text{NO}_2 + \text{M} \rightarrow \text{N}_2\text{O}_5 + \text{M}$  ( $\text{M} = \text{N}_2, \text{O}_2$ ). *Int. J. Chem. Kinet.*, 19, 243-249.
- Wallington, T. J., R. Liu, P. Dagaut and M. J. Kurylo (1988) The gas phase reactions of hydroxyl radicals with a series of aliphatic ethers over the temperature range 240-440 K. *Int. J. Chem. Kinet.*, 20, 41-49.
- Winer, A. M., R. Atkinson, J. Arey, S. M. Aschmann, M. A. Goodman and E. C. Tuazon (1987) Lifetimes and fates of toxic chemicals in California's atmosphere. Final Report to California Air Resources Board Contract No. A5-104-32, Sacramento, CA, August.
- Zetzsch, C. (1980) "Neuere Methoden zur Untersuchung von Radikalreaktionen in der Gasphase." Presented at Bunsenkolloquium, Göttingen, West Germany, September 10.





APPENDIX E

ATMOSPHERIC LIFETIME AND FATE OF FORMALDEHYDE

A Report Submitted in Partial Fulfillment of  
California Air Resources Board Contract No. A732-107

"Lifetimes and Fates of Toxic Air Contaminants in  
California's Atmosphere"

Roger Atkinson

August 1989

Statewide Air Pollution Research Center  
University of California  
Riverside, CA 92521



## TABLE OF CONTENTS

	<u>Page</u>
A. Introduction.....	E-3
B. Chemical Loss Processes.....	E-5
1. Reaction with Ozone.....	E-5
2. Reaction with the Hydroxyl Radical.....	E-5
3. Reaction with the NO <sub>3</sub> Radical.....	E-11
4. Reaction with the HO <sub>2</sub> Radical.....	E-13
5. Photolysis.....	E-14
C. Physical Loss Processes.....	E-16
D. Tropospheric Lifetime of Formaldehyde.....	E-18
E. Atmospheric Formation of HCHO.....	E-19
F. Ambient Atmospheric Concentrations of HCHO.....	E-21
G. Conclusions.....	E-23
H. References.....	E-23



#### A. Introduction

The atmospheric fates of chemical compounds emitted into the troposphere are governed by a number of chemical and physical removal processes. As the result of laboratory studies carried out over the past 20 years, the potentially important gas-phase chemical reaction pathways which must be considered includes:

- Photolysis during daylight hours.
- Reaction with the hydroxyl (OH) radical during daylight hours.
- Reaction with the hydroperoxyl ( $\text{HO}_2$ ) radical, mainly during afternoon/evening hours.
- Reaction with the nitrate ( $\text{NO}_3$ ) radical during nighttime hours.
- Reaction with nitrogen dioxide ( $\text{NO}_2$ ).
- Reaction with ozone ( $\text{O}_3$ ).
- Reaction with gaseous nitric acid ( $\text{HNO}_3$ ).

For the great majority of organic compounds, the most important of these remain photolysis and reaction with hydroxyl (OH) and nitrate ( $\text{NO}_3$ ) radicals and with ozone ( $\text{O}_3$ ).

However, for a limited number of organic compounds, one or more of the other reactive chemical species in the troposphere may react with organic compounds at significant rates. For example,  $\text{HO}_2$  radicals react with formaldehyde (see Section B.4, below) and, more slowly, with acetaldehyde and glyoxal;  $\text{NO}_2$  reacts with dialkenes; and gaseous  $\text{HNO}_3$  reacts with basic compounds such as the amines.

Additionally, for chemical compounds present in the adsorbed phase, photolysis and reaction with  $\text{O}_3$ ,  $\text{N}_2\text{O}_5$ ,  $\text{NO}_2$ ,  $\text{HNO}_3$ ,  $\text{HONO}$ ,  $\text{H}_2\text{SO}_4$  and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) may contribute to their degradation. For certain of these reactions involving adsorbed organics, synergistic interactions may be of importance.

The physical removal processes can be defined in a simplistic manner as accretion (or coagulation) of particles, and dry and wet deposition of gases and particles. Removal of gases and particles at ground surfaces (including snow and other moist surfaces) is referred to as dry deposition, while removal of these species by precipitation is referred to as wet deposition. Our current understanding of these processes, which are

dynamic in nature, must be considered to be semi-quantitative at the present time, and in general there is a lack of experimental data concerning these processes for most organic compounds (Atkinson, 1988; Bidleman, 1988).

The atmospheric lifetime  $\tau$  of a chemical is defined as the time to decay to a concentration of  $1/e$  ( $\approx 0.368$ ) of that initially present. Also,

$$\frac{1}{\tau} = \frac{1}{\tau_{\text{chemical}}} + \frac{1}{\tau_{\text{physical}}} \quad (1)$$

where  $\tau_{\text{chemical}}$  and  $\tau_{\text{physical}}$  are the lifetimes of the chemical with respect to only chemical reaction and only physical loss processes, respectively. These chemical and physical lifetimes are composites of those for the individual loss processes, for example

$$\frac{1}{\tau_{\text{chemical}}} = \frac{1}{\tau_{\text{photolysis}}} + \frac{1}{\tau_{\text{OH}}} + \frac{1}{\tau_{\text{NO}_3}} + \frac{1}{\tau_{\text{O}_3}} \dots \quad (2)$$

where  $\tau_{\text{photolysis}}$ ,  $\tau_{\text{OH}}$ ,  $\tau_{\text{NO}_3}$  and  $\tau_{\text{O}_3}$  are the lifetimes with respect to photolysis, reaction with the OH radical, reaction with the  $\text{NO}_3$  radical, and reaction with  $\text{O}_3$ , respectively. In turn, these reactive loss processes are determined by the rate constants,  $k_x$ , for reaction and the ambient atmospheric concentrations,  $[X]$ , of the reactive intermediates. For example

$$\frac{1}{\tau_{\text{OH}}} = k_{\text{OH}}[\text{OH}]$$

For photolysis, the photolysis rate depends on the absorption cross section ( $\sigma$ ), the photolysis quantum yield ( $\phi$ ), and the radiation intensity ( $J$ ), all of which are wavelength dependent:

$$\frac{1}{\tau_{\text{photolysis}}} = k_{\text{photolysis}} = \int_{\sim 290 \text{ nm}}^{\sim 800 \text{ nm}} \sigma_{\lambda} \phi_{\lambda} J_{\lambda} d\lambda$$

In this report, the available literature data concerning the chemical and physical removal processes for formaldehyde (HCHO) are presented and its resulting atmospheric lifetime is estimated. The atmospheric chemistry of this compound is also discussed. Although formaldehyde is gaseous at room temperature, its dissolution into the aqueous phase and subsequent aqueous-phase reactions must also be considered in addition to the gas-phase processes.

## B. Chemical Loss Processes

### 1. Reaction with Ozone

The only kinetic data for the gas-phase reaction of  $O_3$  with HCHO is the upper limit to the rate constant of  $k_1 \leq 2.1 \times 10^{-24} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  obtained by Braslavsky and Heicklen (1976) at 298 K. This upper limit to the rate constant is consistent with the small amount of rate constant data for the other saturated aldehydes  $CH_3CHO$ ,  $(CHO)_2$  and  $CH_3COCHO$  (Atkinson and Carter, 1984).

### 2. Reaction with the Hydroxyl Radical

The available literature data concerning the kinetics and mechanism of the reaction of the OH radical with HCHO have been reviewed and evaluated by Atkinson (1989). The available rate constant data are given in Table 1. The rate constants obtained by Hoare (1962, 1966), Baldwin and Cowe (1962), Blundell et al. (1965), Westenberg and Fristrom (1966), Hoare and Peacock (1966), Morris and Niki (1971a,b), Peeters and Mahnen (1973), Vandooren and Van Tiggelen (1977), Niki et al. (1978), Atkinson and Pitts (1978), Stief et al. (1980), Temps and Wagner (1984) and Zabarnick et al. (1988) for  $^{12}CH_2O$  and of Niki et al. (1984) for  $^{13}CH_2O$  are plotted in Arrhenius form in Figure 1. A significant amount of scatter in these data is evident. Since the rate constant for the self reaction of OH radicals is subject to significant uncertainties (DeMore et al., 1987), the rate constants derived from the study of Smith (1978) are not plotted in Figure 1 and were not used by Atkinson (1989) in the evaluation of the rate constant for this reaction.

It can be seen from Figure 1 that the rate constant for this reaction appears to be approximately independent of temperature over the range -230-500 K, but that at temperatures >500 K the rate constant increases with increasing temperature. At around room temperature, absolute rate

Table 1. Rate Constants  $k_2$  and Temperature-Dependent Parameters ( $k_2 = Ae^{-B/T}$ ) for the Gas-Phase Reaction of the OH Radical with Formaldehyde

$10^{12} \times A$ ( $\text{cm}^3 \text{ mole}^{-1} \text{ s}^{-1}$ )	B (K)	$10^{12} \times k_2$ ( $\text{cm}^3$ molecule $^{-1} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
<u>Formaldehyde</u>					
		$28 \pm 8$	773	RR [relative to $k(\text{CH}_4) = 6.95 \times 10^{-18}$ $\text{T}^2 \text{e}^{-1282/\text{T}}$ ]	Hoare (1962)
		29	798		
		33	873		
		33	923		
		47	813	RR [relative to $k(\text{H}_2) =$ $1.12 \times 10^{-12} \text{J}^a$ ]	Baldwin and Cowe (1962)
		$26 \pm 3$	773	RR [relative to $k(\text{CH}_4) =$ $7.91 \times 10^{-13} \text{J}^a$ ]	Blundell et al. (1965)
-880	-4265		1250-1400	RR [relative to $k(\text{CO}) =$ $1.12 \times 10^{-13} \text{e}^{0.000907\text{T}} \text{J}^a$ ]	Westenberg and Fristrom (1965)
		25	723	RR [relative to $k(\text{CH}_4) =$ $6.95 \times 10^{-18} \text{T}^2 \text{e}^{-1282/\text{T}} \text{J}^a$ ]	Hoare (1966)
		29	798		
		33	923		
		25	798	RR [relative to $k(\text{CH}_4) =$ $8.88 \times 10^{-13} \text{J}^a$ ]	Hoare and Peacock (1966)
		14	798	RR [relative to $k(\text{C}_2\text{H}_6) =$ $5.07 \times 10^{-12} \text{J}^a$ ]	Hoare and Peacock (1966)



Table 1 (continued) - 2

$10^{12} \times A$ ( $\text{cm}^3 \text{ mole}^{-1} \text{ s}^{-1}$ )	B (K)	$10^{12} \times k_2$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
		$\geq 6.6$	300	DF-MS	Herron and Penzhorn (1969)
		$14 \pm 3.5$	298	DF-MS	Morris and Niki (1971a)
		15.3	298	DF-MS	Morris and Niki (1971b)
	-500	42	1600	Flame-MS	Peeters and Mahnen (1973)
		17	485	Flame-MS	Vandooren and Van Tiggelen (1977)
		22	570		
		$15.8 \pm 0.9$	$298 \pm 2$	RR [relative to $k(\text{ethene-d}_4) =$ $8.78 \times 10^{-12} \text{ J}^a$	Niki et al. (1978)
		$9.4 \pm 1.0$	299.3	FP-RF	Atkinson and Pitts (1978)
		$9.4 \pm 1.0$	356.5		
	$88 \pm 151$	$10.3 \pm 1.1$	426.4		
		$5.5 \pm 0.7$	268	DF-MS [relative to $k(\text{OH} + \text{OH})_2 =$ $4.2 \times 10^{-12} \text{ e}^{-240/T} \text{ J}^b$	Smith (1978)
		$5.5 \pm 0.7$	298		
		$7.4 \pm 1.0$	334		
		$11.22 \pm 0.98$	228	FP-RF	Stief et al. (1980)
		$10.28 \pm 0.90$	257		
		$9.86 \pm 1.13$	298		
$10.5 \pm 1.1$	0	$10.46 \pm 1.50$	362		
		$8.1 \pm 1.7$	296	DF-LMR	Temps and Wagner (1984)

Table 1 (continued) - 3

$10^{12} \times A$ ( $\text{cm}^3 \text{ mole}^{-1} \text{ s}^{-1}$ )	B (K)	$10^{12} \times k_2$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
				LP-LIF	Zabarnick et al. (1988)
		$10.7 \pm 1.3$	296		
		$12.5 \pm 0.5$	297		
		$11.4 \pm 0.6$	297		
		$13.6 \pm 1.0$	297		
		$13.2 \pm 0.4$	298		
		$13.1 \pm 0.4$	298		
		$13.7 \pm 0.5$	299		
		$12.0 \pm 0.3$	301		
		$13.9 \pm 0.4$	378		
		$13.3 \pm 0.3$	473		
		$11.4 \pm 0.5$	567		
		$15.8 \pm 1.2$	572		
		$14.1 \pm 0.5$	574		
$16.6 \pm 2.0$	$86 \pm 40$	$16.7 \pm 0.8$	576		
<u>Formaldehyde-<math>^{13}\text{C}</math></u>		$8.40 \pm 0.51$	$299 \pm 2$	RR [relative to $k(\text{ethene}) - \bar{\nu}_{12}^a$ $8.48 \times 10^{-12} \text{ J a}$	Niki et al. (1984)
<u>Formaldehyde-<math>\text{d}_1</math></u>		$\sim 14$	298	DF-MS	Morris and Niki (1971a)

<sup>a</sup>From Atkinson (1989).<sup>b</sup>From DeMore et al. (1987).




Figure 1. Arrhenius plot of rate constants for the reaction of the OH radical with formaldehyde. ( $\diamond$ ) Hoare (1962, 1966), Blundell et al. (1965), Hoare and Peacock (1966); ( $\blacklozenge$ ) Baldwin and Cowe (1962); (— — —) Westenberg and Fristrom (1965); (x) Morris and Niki (1971a); ( $\nabla$ ) Morris and Niki (1971b); (—|—) Peeters and Mahnen (1973); (— · — · —) Vandooren and Van Tiggelen (1977); ( $\blacksquare$ ) Niki et al. (1978); ( $\bullet$ ) Atkinson and Pitts (1978); ( $\triangle$ ) Stief et al. (1980); ( $\blacktriangle$ ) Temps and Wagner (1984); ( $\circ$ ) Zabarnick et al. (1988); ( $\square$ ) Niki et al. (1984) [for reaction with formaldehyde- $^{13}\text{C}$ ]; (—) recommendation (see text).

constants have been determined by Morris and Niki (1971a), Atkinson and Pitts (1978), Stief et al. (1980), Temps and Wagner (1984) and Zabarnick et al. (1988). Again, a significant amount of scatter is observed, with Morris and Niki (1971a) and Zabarnick et al. (1988) obtaining rate constants of  $(1.2-1.4) \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , Atkinson and Pitts (1978) and Stief et al. (1980) rate constants of  $(9.4-9.9) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , and Temps and Wagner (1984) a rate constant of  $8.1 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ .

Based upon the data shown in Figure 1, the Arrhenius plot exhibits significant curvature. A unit-weighted least-squares analysis of the absolute rate constants determined by Atkinson and Pitts (1978) and Stief et al. (1980) (which are in excellent agreement) was carried out by Atkinson (1989), using the expression  $k_2 = CT^2e^{-D/T}$ , to obtain the recommendation of

$$k_2(\text{HCHO}) = (1.25^{+0.20}_{-0.18}) \times 10^{-17} T^2 e^{(648 \pm 45)/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

over the temperature range 228-426 K, where the indicated errors are two least-squares standard deviations, and

$$k_2(\text{HCHO}) = 9.77 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$


at 298 K, with an estimated overall uncertainty at 298 K of  $\pm 30\%$ . This recommendation is  $\sim 10\%$  higher than that recommended by Atkinson (1986) of  $k_2(\text{HCHO}) = 9.0 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , independent of temperature over the same range of 228-426 K. At elevated temperatures, the recommended expression yields calculated rate constants in good agreement with those obtained from the flame studies of Westenberg and Fristrom (1965) and Peeters and Mahnen (1973) [Figure 1].

As expected, the rate constant for the reaction of OH radicals with formaldehyde- $^{13}\text{C}$  is, within the likely experimental errors, essentially identical to that for formaldehyde- $^{12}\text{C}$  (Niki et al., 1984).

This OH radical reaction with formaldehyde can proceed by the pathways



Morrison and Heicklen (1980), Temps and Wagner (1984) and Niki et al. (1984) have shown from product studies that reaction pathway (2b) is negligible, accounting for  $\leq 2\%$  of the overall reaction (Niki et al., 1984). Morrison and Heicklen (1980) did not observe any formation ( $< 10\%$ ) of HCOOH, and concluded that reaction pathways (2a) and (2c) occur with approximately equal probability. More recently, Temps and Wagner (1984), using a discharge flow technique with LMR detection to monitor both OH and HCO radicals, have shown that reaction pathway (2a) accounts for  $100 \pm 5\%$  of the overall reaction. Thus, at room temperature the OH radical reaction with formaldehyde proceeds essentially entirely by the H-atom abstraction process.




The resulting HCO radical reacts rapidly with  $\text{O}_2$  to yield the  $\text{HO}_2$  radical (DeMore et al., 1987; Atkinson et al., 1989):



### 3. Reaction with the $\text{NO}_3$ Radical

Kinetic data for the gas-phase reaction of the  $\text{NO}_3$  radical with HCHO have been obtained by Atkinson et al. (1984), Cantrell et al. (1985) and Hjorth et al. (1988). All of these studies, except for five experiments in the study of Cantrell et al. (1985), were relative rate measurements carried out at room temperature, with the rate constants for the  $\text{NO}_3$  radical reaction with HCHO being relative to the equilibrium constant K for the reactions



While the reported rate constants  $k_3$  range from  $(3.2-6.3) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , at least part of these differences are due to the differing values of  $K$  used. The available rate data are given in Table 2, with the relative rate data from Atkinson et al. (1984), Cantrell et al. (1985) and Hjorth et al. (1988) being re-evaluated by use of an equilibrium constant of  $K = 1.26 \times 10^{-27} e^{11275/T} \text{ cm}^3 \text{ molecule}^{-1}$  ( $K = [\text{N}_2\text{O}_5]/[\text{NO}_2][\text{NO}_3]$ ). The (re-evaluated) relative rate constant of Atkinson et al. (1984) agrees very well with the rate constant obtained from the absolute rate measurements of Cantrell et al. (1985), while the re-evaluated rate constants derived from the relative rate data of Cantrell et al. (1985) and Hjorth et al. (1988) are 30-50% higher.


Table 2. Room Temperature Rate Constants,  $k_3$ , for the Gas-Phase Reaction of the  $\text{NO}_3$  Radical with Formaldehyde

$10^{16} \times k_3$ ( $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ )	at T (K)	Reference
$5.9 \pm 0.5^a$	$298 \pm 1$	Atkinson et al. (1984)
$\left. \begin{array}{l} 5.6^b \\ 8.9 \pm 0.7^a \end{array} \right\}$	$298 \pm 2$	Cantrell et al. (1985)
$7.9 \pm 1.6^a$	$295 \pm 2$	Hjorth et al. (1988)

<sup>a</sup>Relative to an equilibrium constant of  $K = 1.26 \times 10^{-27} e^{11275/T} \text{ cm}^3 \text{ molecule}^{-1}$ , derived from the experimental data of Burrows et al. (1985), Tuazon et al. (1984), Kircher et al. (1984) and Cantrell et al. (1988).


<sup>b</sup> $\text{NO}_3$  radicals monitored directly, leading to an absolute rate measurement.

Both of the recent NASA (DeMore et al., 1987) and IUPAC (Atkinson et al., 1989) data evaluations recommend a rate constant for the reaction of the  $\text{NO}_3$  radical with  $\text{HCHO}$  of  $6.0 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K, with an estimated uncertainty of  $\pm$  a factor of 1.5-2. Moreover, the room temperature rate constant for the reaction of the  $\text{NO}_3$  radical with  $\text{CH}_3\text{CHO}$  also determined by Atkinson et al. (1984), when re-evaluated as discussed above, is in excellent agreement with the recent absolute rate constant of Dlugokencky and Howard (1989). Accordingly, the rate constant for the reaction of the  $\text{NO}_3$  radical with  $\text{HCHO}$  is taken to be  $k_3 = 6 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K. As for the OH radical reaction, this reaction proceeds by H-atom abstraction




#### 4. Reaction with the $\text{HO}_2$ Radical


The  $\text{HO}_2$  radical has been shown to react with  $\text{HCHO}$  (Su et al., 1979a,b; Niki et al., 1980a,b; Veyret et al., 1982, 1989; Barnes et al., 1985), with this reaction proceeding by initial  $\text{HO}_2$  radical addition to form the  $\text{HOOCH}_2\dot{\text{O}}$  species



This alkoxy radical then rapidly undergoes isomerization via a 5-membered transition state to the peroxy radical  $\text{HOCH}_2\text{OO}\dot{\text{O}}$



Furthermore, the available data show that this sequence of reactions is reversible, and hence that the reactions to be considered are



The rate constants for the forward [reaction (4)] and back [reaction (-4)] reactions are given in Table 3. Clearly, the reported rate data span a significant range, especially for reaction (-4). The most recent IUPAC data evaluation (Atkinson et al., 1989) utilizes the rate constants of Barnes et al. (1985) and Veyret et al. (1989) to recommend that

$$k_4 = 9.7 \times 10^{-15} e^{625/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$


$$= 7.9 \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } 298 \text{ K}$$

and

$$k_{-4} = 2.4 \times 10^{12} e^{-7000/T} \text{ s}^{-1}$$

$$= 150 \text{ s}^{-1} \text{ at } 298 \text{ K.}$$

Since  $\text{HO}_2$  radicals react rapidly with NO (DeMore et al., 1987; Atkinson et al., 1989), and the other reaction of the  $\text{HOCH}_2\text{OO}$  radical [in addition to reaction (-4)] is expected to be with NO,



the magnitude of the rate constant for decomposition of the  $\text{HOCH}_2\text{OO}$  radical [reaction (-4)] means that the reaction of the  $\text{HO}_2$  radical with HCHO is of minor importance as a tropospheric loss process for HCHO.

#### 5. Photolysis


The absorption cross sections and quantum yields for the photolysis of HCHO have most recently been reviewed and evaluated by DeMore et al. (1987) and Atkinson et al. (1989). The most recent IUPAC evaluation (Atkinson et al., 1989) accepts the absorption cross-section data of Moortgat et al. (1983). At the longer wavelengths, these absorption cross-sections of Moortgat et al. (1983) are higher than those measured by Bass et al. (1980), but are substantiated by recent unpublished data of H. W. Biermann (University of California, Riverside). The recommended absorption cross-sections and quantum yields (Atkinson et al., 1989) for the processes



are tabulated in Table 4.



Table 3. Rate Constants  $k_f$  and  $k_{-f}$  for the Gas-Phase Reaction of the  $\text{HO}_2$  Radical with



$10^{14} \times k_f$ ( $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ )	$k_{-f}$ ( $\text{s}^{-1}$ )	at T (K)	Reference
1.0	1.5	$298 \pm 2$	Su et al. (1979a,b)
$7.5 \pm 3.5$	30	298	Veyret et al. (1982)
$11 \pm 4$	$20^{+20}_{-10}$	273	Barnes et al. (1985)
$0.77 e^{(625 \pm 550)/T}$	$2.0 \times 10^{12} e^{-(7000 \pm 2000)/T}$	$275-333$	Veyret et al. (1989)
$6.0 \pm 0.7$	$100 \pm 50$	295	

Table 4. Absorption Cross Sections,  $\sigma$ , and Quantum Yields,  $\phi$ , for the Photolysis of HCHO (from Atkinson et al., 1989)


$\lambda$ (nm)	$10^{20} \sigma(\text{cm}^2)^a$		
	298 K	$\phi_{5a}$	$\phi_{5b}$
270	0.95	0.38	0.43
280	1.80	0.57	0.32
290	2.93	0.73	0.24
300	4.06	0.78	0.21
310	4.60	0.78	0.22
320	4.15	0.62	0.38
330	3.21	0.27	0.66
340	2.22	0	0.56 <sup>b</sup>
350	1.25	0	0.21 <sup>b</sup>
360	0.18	0	0.03 <sup>b</sup>

<sup>a</sup>The values are averaged for 10 nm intervals centered on the indicated wavelength.

<sup>b</sup>760 torr total pressure of air.

### C. Physical Loss Processes

In addition to the tropospheric chemical loss processes of HCHO discussed above, HCHO is soluble in water and is hence incorporated into cloud, rain and fog water, with subsequent aqueous phase reactions (see, for example, Graedel and Weschler, 1981; Jacob and Hoffmann, 1983; Munger et al., 1983, 1986; Adewuyi et al., 1984; Jacob, 1986; McElroy, 1986; Graedel et al., 1986; Pandis and Seinfeld, 1989). In the aqueous phase, formaldehyde is present as the glycol through the reaction



with this equilibrium almost totally favoring the glycol (Jacob, 1986), with  $[\text{H}_2\text{C(OH)}_2\text{aq}]/[\text{HCHO(aq)}] = 2.3 \times 10^3$  at 298 K (Betterton and Hoffmann, 1988).



Betterton and Hoffmann (1988) measured the apparent Henry's law coefficient,  $H^*$ , where

$$H^* = \{[\text{HCHO(aq)}] + [\text{H}_2\text{C(OH)}_2\text{(aq)}]\}/[\text{HCHO(gas)}]$$



as a function of temperature over the range 288-318 K. At 298 K the value of  $H^*$  determined was  $H^* = (3.0 \pm 0.7) \times 10^3 \text{ mol atm}^{-1}$ .

For gas-phase organic compounds which are very efficiently rained out (and for particles, which are also very efficiently rained out), the washout ratio  $W$ , where  $W = \text{concentration in rain}/\text{concentration in air}$  ( $W = RTH^*$  with the definition of  $H^*$  given above), is  $10^5$  to  $10^6$  (Eisenreich et al., 1981; Atkinson, 1988). The value of  $H^*$  of Betterton and Hoffmann (1988) corresponds to a washout ratio of  $W = 7.3 \times 10^4$  at 298 K. With a washout ratio of this magnitude, wet deposition of HCHO is expected to be significant as a tropospheric loss process for HCHO during rain or fog events.


Incorporation of HCHO into rain, cloud and fog water is followed by a series of aqueous-phase reactions, including the oxidation of formaldehyde to formic acid (see, for example, Jacob, 1986; Pandis and Seinfeld, 1989).



Furthermore, the complexation of sulfur [S(IV)] to hydroxymethanesulfonate



leads to stabilization of S(IV) in clouds under nighttime conditions. In the presence of light, the reaction of aqueous OH radicals with hydroxymethanesulfonate



is expected to be reasonably rapid, leading to a lifetime of hydroxymethanesulfonate in cloudwater of the order of 30 min (Jacob, 1986).

The solubility of HCHO in water leads to efficient wet deposition of HCHO, with gas-phase HCHO being expected to be efficiently scavenged into rain, cloud and fog water. This wet deposition process is, of course, episodic, and will, on average, lead to a tropospheric lifetime of HCHO somewhat shorter than calculated solely from the gas-phase loss processes discussed above (see below).

#### D. Tropospheric Lifetime of Formaldehyde



As discussed above, the tropospheric lifetime of HCHO is determined from the loss rates of both the chemical and physical loss processes. Based upon the rate constants given above for the gas-phase reactions of HCHO with OH, NO<sub>3</sub> and HO<sub>2</sub> radicals and with O<sub>3</sub>, and measured or estimated tropospheric concentrations of these reactive species of: OH radicals,  $1.5 \times 10^6$  molecule cm<sup>-3</sup> during daylight hours (Prinn et al., 1987); NO<sub>3</sub> radicals,  $2.4 \times 10^8$  molecule cm<sup>-3</sup> during nighttime hours (Platt et al., 1984; Atkinson et al., 1986); HO<sub>2</sub> radicals,  $\sim 10^7$  molecule cm<sup>-3</sup> (Hard et al., 1984; Zellner and Weibring, 1989); and O<sub>3</sub>,  $7 \times 10^{11}$  molecule cm<sup>-3</sup> throughout a 24-hr period (Logan, 1985), the tropospheric loss rates of HCHO due to these chemical processes can be calculated. The calculated lifetimes due to these individual reactions are then: reaction with the OH radical, 1.6 days; reaction with the NO<sub>3</sub> radical, 160 days; reaction with the HO<sub>2</sub> radical (neglecting the back-decomposition reaction of the HOOCH<sub>2</sub>O radical),  $\sim 15$  days; and reaction with O<sub>3</sub>,  $> 2 \times 10^4$  years. Since the back-decomposition of the HOCH<sub>2</sub>O radical formed from the HO<sub>2</sub> radical reaction significantly decreases the importance of the HO<sub>2</sub> radical reaction as an HCHO loss process [by around two orders of magnitude for  $2 \times 10^{11}$  molecule cm<sup>-3</sup> (10 ppb) of NO], the above calculated lifetime of HCHO due to reaction with the HO<sub>2</sub> radical reaction is likely to be of minor importance as a tropospheric loss process.

Clearly, the daytime OH radical reaction is the dominant of these chemical reaction loss processes. Photolysis of HCHO is estimated to lead to a lifetime (due to photolysis) of 0.4 day in the lower troposphere for a zenith angle of 40° (Atkinson, 1988; W. P. L. Carter, University of California, Riverside, private communication, 1989). Thus, photolysis and


reaction with the OH radical are the major tropospheric chemical loss processes, with an overall lifetime of HCHO of ~0.3 days and with photolysis being the dominant removal route. Wet deposition will, on average, lead to a shorter tropospheric lifetime, but since wet deposition is episodic in nature, the above lifetime calculated from OH radical reaction and photolysis is best regarded as being the tropospheric lifetime, with faster removal of HCHO from the lower troposphere during rain and/or fog events. It must also be noted that photolysis and reaction with the OH radical only occur during daytime hours.

#### E. Atmospheric Formation of HCHO



Formaldehyde is formed in the troposphere from the tropospheric degradation reactions of many organic compounds. In the "clean" troposphere, HCHO is formed as a product of the atmospheric reactions of methane:




[involving rapid formation of  $\text{CH}_3\text{O}_2$ , followed by reactions of the methyl peroxy radical with  $\text{NO}$ ,  $\text{NO}_2$ ,  $\text{HO}_2$  radicals and other organic peroxy radicals to ultimately lead to formation of the methoxy ( $\text{CH}_3\dot{\text{O}}$  radical (Ravishankara, 1988; Atkinson, 1990)], followed by the reaction




The formation of  $\text{CH}_3\dot{\text{O}}$  and/or  $\dot{\text{C}}\text{H}_3$  radicals during the atmospheric degradation reactions of organic compounds thus leads to the formation of HCHO. Sources of  $\dot{\text{C}}\text{H}_3$  radicals include the important tropospheric reactions of acetaldehyde




and the decomposition reactions of the more complex alkoxy radicals; for example




Analogous decompositions of alkoxy radicals can also lead directly to HCHO. For example, the  $\beta$ -hydroxyalkoxy radicals formed after OH radical addition to the terminal alkenes (such as the  $\text{CH}_3\text{CHOHCH}_2\text{O}\cdot$  radical formed from propene) decompose, at least in part, as follows:



The simplest  $\alpha$ -hydroxy radical,  $\dot{\text{C}}\text{H}_2\text{OH}$ , which is formed from the decomposition reactions of  $\text{R}_1\text{R}_2\text{C}(\dot{\text{O}})\text{CH}_2\text{OH}$   $\beta$ -hydroxyalkoxy radicals, reacts rapidly with  $\text{O}_2$  to yield HCHO and the  $\text{HO}_2$  radical (Atkinson and Lloyd, 1984; DeMore et al., 1987; Atkinson et al., 1989)



In addition, the gas-phase reactions of  $\text{O}_3$  with alkenes containing terminal  $=\text{CH}_2$  groups lead to the formation of HCHO. For example, for the reaction of  $\text{O}_3$  with ethene at room temperature and atmospheric pressure of air (Atkinson and Lloyd, 1984; Atkinson and Carter, 1984; Atkinson, 1990)



with similar HCHO yields for other 1-alkenes (Atkinson, 1990).

Thus, in polluted airmasses characteristic of urban areas, the atmospheric formation of HCHO from organic precursors occurs. Grosjean et al. (1983) concluded from ambient measurements that in many cases the atmospheric formation of HCHO dominates over direct emission of HCHO from combustion sources. More recent ambient air monitoring data in the Los Angeles air basin (Lawson et al. 1990) provide clear evidence for directly emitted HCHO as well as HCHO formed in the atmosphere by photochemical reactions of organic precursors. However, Lawson et al. (1990) also conclude that the atmospheric formation of HCHO dominates over direct emissions of HCHO under conditions of high photochemical activity in the Los Angeles air basin.

#### F. Ambient Atmospheric Concentrations of HCHO

The measured HCHO mixing ratio in the clean lower troposphere is around 0.2 parts-per-billion (ppb) [Lowe et al., 1981; Lowe and Schmidt, 1983], with these measured mixing ratios being in good agreement with the predicted values of 0.25-0.35 ppb presented by Logan et al. (1981) for the lower troposphere in the northern hemisphere. Reported data for the ambient concentrations of HCHO in California since 1980 are given in Table 5. These particular data were all collected in the Los Angeles air basin, and vary from <1 ppb up to 86 ppb, with the most recent concentrations measured in the California Air Resources Board-funded intercomparison studies at Claremont and Glendora in the Los Angeles air basin being significantly lower,  $\leq 25$  ppb (Grosjean, 1988; Lawson et al. 1990). Further, as yet unpublished, ambient HCHO data are available from the 1987 ARB-funded South Coast Air Quality Study (SCAQS).

It may be expected that the ambient atmospheric HCHO levels encountered in the Los Angeles air basin are among the highest to be expected in California, and hence the ambient atmospheric HCHO mixing ratios in California range from a few tenths of a part-per-billion (for clean tropospheric air) up to a few tens of ppb, with the actual values depending to a large extent on the organic precursor emissions, the extent of photochemical activity and the meteorological conditions pertaining.

Table 5. Recent Measurements of Formaldehyde Concentrations in Ambient Air in California

HCHO Mixing Ratio (ppb)	Measurement Date and Place	Reference
2-40	5-6/1980, Los Angeles	Grosjean (1982)
3-48	9-10/1980, Claremont	Grosjean (1982)
10-41	7/1980, Riverside	Singh et al. (1982)
18-60	7-10/1980, Los Angeles	Grosjean et al. (1983)
65-70	7-8/1980, Burbank	Grosjean et al. (1983)
53	7/1980, Pasadena	Grosjean et al. (1983)
5	8/1980, Pacoima	Grosjean et al. (1983)
33	8/1980, Newhall	Grosjean et al. (1983)
38-47	9-10/1980, Rosemead	Grosjean et al. (1983)
27	9/1980, Covina	Grosjean et al. (1983)
42	9/1980, Cucamonga	Grosjean et al. (1983)
59-66	10/1980, El Monte	Grosjean et al. (1983)
34	10/1980, San Dimas	Grosjean et al. (1983)
34	10/1980, Upland	Grosjean et al. (1983)
0.7-35	10/1980, Azusa	Grosjean et al. (1983)
0.5-40	10/1980, Lennox	Grosjean et al. (1983)
4-86	9-11/1981, Los Angeles	Grosjean and Fung (1984)
1.5-11	9/1985, Claremont	Grosjean (1988)
3-22	8/1986, Glendora	Lawson et al. (1990)



#### G. Conclusions

Formaldehyde is both directly emitted into the atmosphere as well as being formed in the atmosphere from the photochemical degradation of more methane and other non-methane organic compounds. HCHO is removed from the lower troposphere mainly by photolysis and reaction with the OH radical, and by wet deposition (leading to incorporation of formaldehyde into rain, cloud and fog water). The estimated lifetime of HCHO due to photolysis and OH radical reaction is ~0.3 days, with episodic wet deposition events leading to shorter lifetimes. Reported ambient atmospheric HCHO concentrations in the Los Angeles air basin (expected to have among the higher levels encountered in California) range up to 86 ppb, with mixing ratios measured during two recent intercomparison studies in 1985 and 1986 ranging up to 25 ppb.

#### H. References

- Adewuyi, Y. G., S.-Y. Cho, R.-P. Tsay and G. R. Carmichael (1984): Importance of formaldehyde in cloud chemistry. *Atmos. Environ.*, 18, 2413-2420.
- Atkinson, R. and J. N. Pitts, Jr. (1978): Kinetics of the reactions of the OH radical with HCHO and CH<sub>3</sub>CHO over the temperature range 299-426°K. *J. Chem. Phys.*, 68, 3581-3584.
- Atkinson, R. and W. P. L. Carter (1984): Kinetics and mechanisms of the gas-phase reactions of ozone with organic compounds under atmospheric conditions. *Chem. Rev.*, 84, 437-470.
- Atkinson, R. and A. C. Lloyd (1984): Evaluation of kinetic and mechanistic data for modeling of photochemical smog. *J. Phys. Chem. Ref. Data*, 13, 315-444.
- Atkinson, R., C. N. Plum, W. P. L. Carter, A. M. Winer, and J. N. Pitts, Jr. (1984) Rate constants for the gas-phase reactions of nitrate radicals with a series of organics in air at 298 ± 1 K. *J. Phys. Chem.*, 88, 1210-1215.
- Atkinson, R. (1986): Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. *Chem. Rev.*, 86, 69-201.
- Atkinson, R., A. M. Winer and J. N. Pitts, Jr. (1986): Estimation of night-time N<sub>2</sub>O<sub>5</sub> concentrations from ambient NO<sub>2</sub> and NO<sub>3</sub> radical concentrations and the role of N<sub>2</sub>O<sub>5</sub> in night-time chemistry. *Atmos. Environ.*, 20, 331-339.

- Atkinson, R. (1988): Atmospheric transformations of automotive emissions. In: Air Pollution, the Automobile, and Public Health, A. Y. Watson, R. R. Bates and D. Kennedy (eds.), National Academy Press, Washington, D.C., pp. 99-132.
- Atkinson, R. (1989): Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. J. Phys. Chem. Ref. Data, Monograph. 1, 1-246.
- Atkinson, R., D. L. Baulch, R. A. Cox, R. F. Hampson, Jr., J. A. Kerr and J. Troe (1989): Evaluated kinetic and photochemical data for atmospheric chemistry: Supplement III. J. Phys. Chem. Ref. Data, 18, 881-1097.
- Atkinson, R. (1990): Gas-phase tropospheric chemistry of organic compounds: a review. Atmos. Environ., 24A, 1-41.
- Baldwin, R. R. and D. W. Cowe (1962): The inhibition of the hydrogen + oxygen reaction by formaldehyde. Trans. Faraday Soc., 58, 1768-1781.
- Barnes, I., K. H. Becker, E. H. Fink, A. Reimer, F. Zabel and H. Niki (1985): FT-IR spectroscopic study of the gas-phase reaction of HO<sub>2</sub> with H<sub>2</sub>CO. Chem. Phys. Lett., 115, 1-8.
- Bass, A. M., L. C. Glasgow, C. Miller, J. P. Jesson and D. L. Filkin (1980): Temperature dependent absorption cross sections for formaldehyde (CH<sub>2</sub>O): the effect of formaldehyde on stratospheric chlorine chemistry. Planet. Space Sci., 28, 675-679.
- Betterton, E. A. and M. R. Hoffmann (1988): Henry's law constants of some environmentally important aldehydes. Environ. Sci. Technol., 22, 1415-1418.
- Bidleman, T. F. (1988): Atmospheric processes. Environ. Sci. Technol., 22, 361-367.
- Blundell, R. V., W. G. A. Cook, D. E. Hoare and G. S. Milne (1965): Rates of radical reactions in methane oxidation. 10th International Symposium on Combustion, 1964; The Combustion Institute, Pittsburgh, PA, pp. 445-452.
- Braslavsky, S. and J. Heicklen (1976): The gas-phase reaction of O<sub>3</sub> with H<sub>2</sub>CO. Int. J. Chem. Kinet., 8, 801-808.
- Burrows, J. P., G. S. Tyndall and G. K. Moortgat (1985): A study of the N<sub>2</sub>O<sub>5</sub> equilibrium between 275 and 315 K and determination of the heat of formation of NO<sub>3</sub>. Chem. Phys. Lett., 119, 193-198.
- Cantrell, C. A., W. R. Stockwell, L. G. Anderson, K. L. Busarow, D. Perner, A. Schmeltekopf, J. G. Calvert and H. S. Johnston (1985): Kinetic study of the NO<sub>3</sub>-CH<sub>2</sub>O reaction and its possible role in nighttime tropospheric chemistry. J. Phys. Chem., 89, 139-146.

- Cantrell, C. A., J. A. Davidson, A. H. McDaniel, R. E. Shetter and J. G. Calvert (1988): The equilibrium constant for  $\text{N}_2\text{O}_5 \rightleftharpoons \text{NO}_2 + \text{NO}_3$ : absolute determination by direct measurement from 243 to 397 K. J. Chem. Phys., 88, 4997-5006.
- DeMore, W. B., M. J. Molina, S. P. Sander, D. M. Golden, R. F. Hampson, M. J. Kurylo, C. J. Howard and A. R. Ravishankara (1987): Chemical Kinetics and photochemical data for use in stratospheric modeling. NASA Panel for data evaluation, Evaluation No. 8; Jet Propulsion Laboratory Publication 87-41, September 15.
- Dlugokencky, E. J. and C. J. Howard (1989): Studies of  $\text{NO}_3$  radical reactions with some atmospheric organic compounds at low pressures. J. Phys. Chem., 93, 1091-1096.
- Eisenreich, S. J., B. B. Looney and J. D. Thornton (1981): Airborne organic contaminants in the Great Lakes ecosystem. Environ. Sci. Technol., 15, 30-38.
- Graedel, T. E. and C. J. Weschler (1981): Chemistry within aqueous atmospheric aerosols and raindrops. Rev. Geophys. Space Phys., 19, 505-539.
- Graedel, T. E., M. L. Mandich and C. J. Weschler (1986): Kinetic model studies of atmospheric droplet chemistry. 2. Homogeneous transition state chemistry in raindrops. J. Geophys. Res., 91, 5205-5221.
- Grosjean, D. (1982): Formaldehyde and other carbonyls in Los Angeles ambient air. Environ. Sci. Technol., 16, 254-262.
- Grosjean, D., R. D. Swanson and C. Ellis (1983): Carbonyls in Los Angeles air: contribution of direct emissions and photochemistry. Sci. Total Environ., 29, 65-85.
- Grosjean, D. and K. Fung (1984): Hydrocarbons and carbonyls in Los Angeles air. J. Air Poll. Contr. Assoc., 34, 537-543.
- Grosjean, D. (1988): Aldehydes, carboxylic acids and inorganic nitrate during NSMCS. Atmos. Environ., 22, 1637-1648.
- Hard, T. M., R. J. O'Brien, C. Y. Chan and A. A. Mehrabzadeh (1984): Tropospheric free radical determination by FAGE. Environ. Sci. Technol., 18, 768-777.
- Herron, J. T. and R. D. Penzhorn (1969): Mass spectrometric study of the reactions of atomic oxygen with ethylene and formaldehyde. J. Phys. Chem., 73, 191-196.
- Hjorth, J., G. Ottobriani and G. Restelli (1988): Reaction between  $\text{NO}_3$  and  $\text{CH}_2\text{O}$  in air: determination of the rate constant at  $295 \pm 2$  K. J. Phys. Chem., 92, 2669-2672.
- Hoare, D. E. (1962): Reactions of gaseous hydroxyl radicals. Nature, 194, 283.

- Hoare, D. E. (1966): Studies of the reactions of hydroxyl radicals. I., Proc. Roy. Soc. (London), A291, 73-84.
- Hoare, D. E. and G. B. Peacock (1966): Studies of the reactions of hydroxyl radicals. II. Proc. Roy. Soc. (London), A291, 85-93.
- Jacob, D. J. and M. R. Hoffmann (1983) A dynamic model for the production of  $\text{H}^+$ ,  $\text{NO}_3^-$ , and  $\text{SO}_4^{2-}$  in urban fog. J. Geophys. Res., 88, 6611-6621.
- Jacob, D. J. (1986): Chemistry of OH in remote clouds and its role in the production of formic acid and peroxymonosulfate. J. Geophys. Res., 91, 9807-9826.
- Kircher, C. C., J. J. Margitan and S. P. Sander (1984): Pressure and temperature dependence of the reaction  $\text{NO}_2 + \text{NO}_3 + \text{M} \rightarrow \text{N}_2\text{O}_5 + \text{M}$ . J. Phys. Chem., 88, 4370-4375.
- Lawson, D. R., H. W. Biermann, E. C. Tuazon, A. M. Winer, G. I. Mackay, H. I. Schiff, G. L. Kok, P. K. Dasgupta and K. Fung (1990): Formaldehyde measurement methods evaluation and ambient concentrations during the Carbonaceous Species Methods Comparison Study, Aerosol. Sci. Technol., 12, 64-76.
- Logan, J. A., M. J. Prather, S. C. Wofsy and M. B. McElroy (1981): Tropospheric chemistry: a global perspective. J. Geophys. Res., 86, 7210-7254.
- Logan, J. A. (1985): Tropospheric ozone: seasonal behavior, trends, and anthropogenic influence. J. Geophys. Res., 90, 10463-10482.
- Lowe, D. C., U. Schmidt, D. H. Ehhalt, C. G. B. Frischkorn and K. W. Nümborg (1981): Determination of formaldehyde in clean air. Environ. Sci. Technol., 15, 819-823.
- Lowe, D. C. and U. Schmidt (1983): Formaldehyde (HCHO) measurements in the nonurban atmosphere. J. Geophys. Res., 88, 10844-10858.
- McElroy, W. J. (1986): Sources of hydrogen peroxide in cloudwater. Atmos. Environ., 20, 427-438.
- Moortgat, G. K., W. Seiler and P. Warneck (1983): Photodissociation of HCHO in air: CO and  $\text{H}_2$  quantum yields. J. Chem. Phys., 78, 1185-1190.
- Morris, E. D., Jr. and H. Niki (1971a): Mass spectrometric study of the reaction of hydroxyl radical with formaldehyde. J. Chem. Phys., 55, 1991-1992.
- Morris, E. D., Jr. and H. Niki (1971b): Reactivity of hydroxyl radicals with olefins. J. Phys. Chem., 75, 3640-3641.
- Morrison, B. M., Jr. and J. Heicklen (1980): The reactions of HO with  $\text{CH}_2\text{O}$  and of HCO with  $\text{NO}_2$ . J. Photochem., 13, 189-199.

- Munger, J. W., D. J. Jacob, J. M. Waldman and M. R. Hoffmann (1983): Fogwater chemistry in an urban atmosphere. *J. Geophys. Res.*, 88, 5109-5121.
- Munger, J. W., C. Tillier and M. R. Hoffmann (1986): Identification of hydroxymethanesulfonate in fog water. *Science*, 231, 247-249.
- Niki, H., P. D. Maker, C. M. Savage and L. P. Breitenbach (1978): Relative rate constants for the reaction of hydroxyl radical with aldehydes. *J. Phys. Chem.*, 82, 132-134.
- Niki, H., P. D. Maker, C. M. Savage and L. P. Breitenbach (1980a): FT-IR studies of the Cl-atom initiated oxidation of formaldehyde: detection of a new metastable species in the presence of NO<sub>2</sub>. *Chem. Phys. Lett.*, 72, 71-73.
- Niki, H., P. D. Maker, C. M. Savage and L. P. Breitenbach (1980b): Further IR spectroscopic evidence for the formation of CH<sub>2</sub>(OH)OOH in the gas-phase reaction of HO<sub>2</sub> with CH<sub>2</sub>O. *Chem. Phys. Lett.*, 75, 533-535.
- Niki, H., P. D. Maker, C. M. Savage and L. P. Breitenbach (1984): Fourier transform infrared study of the kinetics and mechanism for the reaction of the hydroxyl radical with formaldehyde. *J. Phys. Chem.*, 88, 5342-5344.
- Pandis, S. N. and J. H. Seinfeld (1989): Sensitivity analysis of a chemical mechanism for aqueous-phase atmospheric chemistry. *J. Geophys. Res.*, 94, 1105-1126.
- Peeters, J. and G. Mahnen (1973): Reaction mechanisms and rate constants of elementary steps in methane-oxygen flames. 14th International Symposium on Combustion, 1972; The Combustion Institute, Pittsburgh, PA; pp. 133-146.
- Platt, U. F., A. M. Winer, H. W. Biermann, R. Atkinson and J. N. Pitts, Jr. (1984): Measurement of nitrate radical concentrations in continental air. *Environ. Sci. Technol.*, 18, 365-369.
- Prinn, R., D. Cunnold, R. Rasmussen, P. Simmonds, F. Alyea, A. Crawford, P. Fraser and R. Rosen (1987): Atmospheric trends in methylchloroform and the global average for the hydroxyl radical. *Science*, 238, 945-950.
- Ravishankara, A. R. (1988): Kinetics of radical reactions in the atmospheric oxidation of CH<sub>4</sub>. *Ann. Rev. Phys. Chem.*, 39, 367-394.
- Singh, H. B., L. J. Salas and R. E. Stiles (1982): Distribution of selected gaseous organic mutagens and suspect carcinogens in ambient air. *Environ. Sci. Technol.*, 16, 872-880.
- Smith, R. H. (1978): Rate constant and activation energy for the gaseous reaction between hydroxyl and formaldehyde. *Int. J. Chem. Kinet.*, 10, 519-528.

- Stief, L. J., D. F. Nava, W. A. Payne and J. V. Michael (1980): Rate constant for the reaction of hydroxyl radical with formaldehyde over the temperature range 228-362 K. *J. Chem. Phys.*, 73, 2254-2258.
- Su, F., J. G. Calvert, J. H. Shaw, H. Niki, P. D. Maker, C. M. Savage and L. P. Breitenbach (1979a): Spectroscopic and kinetic studies of a new metastable species in the photooxidation of gaseous formaldehyde. *Chem. Phys. Lett.*, 65, 221-225.
- Su, F., J. G. Calvert and J. H. Shaw (1979b): Mechanism of the photo-oxidation of gaseous formaldehyde. *J. Phys. Chem.*, 83, 3185-3191.
- Temps, F. and H. Gg. Wagner (1984): Rate constants for the reactions of OH radicals with  $\text{CH}_2\text{O}$  and  $\text{HCO}$ . *Ber. Bunsenges. Phys. Chem.*, 88, 415-418.
- Tuazon, E. C., E. Sanhueza, R. Atkinson, W. P. L. Carter, A. M. Winer, and J. N. Pitts, Jr. (1984): Direct determination of the equilibrium constant at 298 K for the  $\text{NO}_2 + \text{NO}_3 \rightleftharpoons \text{N}_2\text{O}_5$  reactions. *J. Phys. Chem.*, 88, 3095-3098.
- Vandooren, J. and P. J. Van Tiggelen (1977): Reaction mechanisms of combustion in low pressure acetylene-oxygen flames. 16th International Symposium on Combustion, 1976; The Combustion Institute, Pittsburgh, PA; pp. 1133-1144.
- Veyret, B., J.-C. Rayez and R. Lesclaux (1982): Mechanism of the photo-oxidation of formaldehyde studied by flash photolysis of  $\text{CH}_2\text{O}-\text{O}_2-\text{NO}$  mixtures. *J. Phys. Chem.*, 86, 3424-3430.
- Veyret, B., R. Lesclaux, M.-T. Rayez, J.-C. Rayez, R. A. Cox and G. K. Moortgat (1989): Kinetics and mechanism of the photooxidation of formaldehyde. 1. Flash photolysis study. *J. Phys. Chem.*, 93, 2368-2374.
- Westenberg, A. A. and R. M. Fristrom (1965): H and O atom profiles measured by ESR in  $\text{C}_2$  hydrocarbon- $\text{O}_2$  flames. 10th International Symposium on Combustion 1964; The Combustion Institute, Pittsburgh, PA; pp. 473-487.
- Zabarnick, S., J. W. Fleming and M. C. Lin (1988): Kinetics of hydroxyl radical reactions with formaldehyde and 1,3,5-trioxane between 290 and 600 K. *Int. J. Chem. Kinet.*, 20, 117-129.
- Zellner, R. and G. Weibring (1989): Tropospheric  $\text{HO}_x$  and its response to increased hydrocarbon emissions. *Zet. Phys. Chem. Neue Folge*, 161, S. 167-188.

APPENDIX F

ATMOSPHERIC LIFETIME AND FATE OF ACETALDEHYDE

A Report Submitted in Partial Fulfillment of  
California Air Resources Board Contract No. A732-107

"Lifetimes and Fates of Toxic Air Contaminants in  
California's Atmosphere"

Roger Atkinson

September 1989

Statewide Air Pollution Research Center  
University of California  
Riverside, CA 92521





## TABLE OF CONTENTS

	<u>Page</u>
A. Introduction.....	F-3
B. Chemical Loss Processes.....	F-5
1. Reaction with Ozone.....	F-5
2. Reaction with the Hydroxyl Radical.....	F-5
3. Reaction with the NO <sub>3</sub> Radical.....	F-9
4. Reaction with the HO <sub>2</sub> Radical.....	F-11
5. Photolysis.....	F-12
C. Physical Loss Processes.....	F-13
D. Tropospheric Lifetime and Fate of Acetaldehyde.....	F-14
E. Atmospheric Formation of Acetaldehyde.....	F-19
F. Ambient Atmospheric Concentrations of CH <sub>3</sub> CHO.....	F-21
G. Conclusions.....	F-22
H. References.....	F-23



#### A. Introduction

The atmospheric fates of chemical compounds emitted into the troposphere are governed by a number of chemical and physical removal processes. As the result of laboratory studies carried out over the past 20 years, the potentially important gas-phase chemical reaction pathways which must be considered includes:

- Photolysis during daylight hours.
- Reaction with the hydroxyl (OH) radical during daylight hours.
- Reaction with the hydroperoxyl (HO<sub>2</sub>) radical, mainly during afternoon/evening hours.
- Reaction with the nitrate (NO<sub>3</sub>) radical during nighttime hours.
- Reaction with nitrogen dioxide (NO<sub>2</sub>).
- Reaction with ozone (O<sub>3</sub>).
- Reaction with gaseous nitric acid (HNO<sub>3</sub>).

For the great majority of organic compounds, the most important of these remain photolysis and reaction with hydroxyl (OH) and nitrate (NO<sub>3</sub>) radicals and with ozone (O<sub>3</sub>).

However, for a limited number of organic compounds, one or more of the other reactive chemical species in the troposphere may react with organic compounds at significant rates. For example, HO<sub>2</sub> radicals react with formaldehyde and, more slowly, with acetaldehyde and glyoxal; NO<sub>2</sub> reacts with dialkenes; and gaseous HNO<sub>3</sub> reacts with basic compounds such as the amines.

Additionally, for chemical compounds present in the adsorbed phase, photolysis and reaction with O<sub>3</sub>, N<sub>2</sub>O<sub>5</sub>, NO<sub>2</sub>, HNO<sub>3</sub>, HONO, H<sub>2</sub>SO<sub>4</sub> and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) may contribute to their degradation. For certain of these reactions involving adsorbed organics, synergistic interactions may be of importance.

The physical removal processes can be defined in a simplistic manner as accretion (or coagulation) of particles, and dry and wet deposition of gases and particles. Removal of gases and particles at ground surfaces (including snow and other moist surfaces) is referred to as dry deposition, while removal of these species by precipitation is referred to as wet deposition. Our current understanding of these processes, which are

dynamic in nature, must be considered to be semi-quantitative at the present time, and in general there is a lack of experimental data concerning these processes for most organic compounds (Atkinson, 1988; Bidleman, 1988).

The atmospheric lifetime  $\tau$  of a chemical is defined as the time to decay to a concentration of  $1/e$  ( $\approx 0.368$ ) of that initially present. Also,

$$\frac{1}{\tau} = \frac{1}{\tau_{\text{chemical}}} + \frac{1}{\tau_{\text{physical}}} \quad (1)$$

where  $\tau_{\text{chemical}}$  and  $\tau_{\text{physical}}$  are the lifetimes of the chemical with respect to only chemical reaction and only physical loss processes, respectively. These chemical and physical lifetimes are composites of those for the individual loss processes, for example

$$\frac{1}{\tau_{\text{chemical}}} = \frac{1}{\tau_{\text{photolysis}}} + \frac{1}{\tau_{\text{OH}}} + \frac{1}{\tau_{\text{NO}_3}} + \frac{1}{\tau_{\text{O}_3}} \dots \quad (2)$$

where  $\tau_{\text{photolysis}}$ ,  $\tau_{\text{OH}}$ ,  $\tau_{\text{NO}_3}$  and  $\tau_{\text{O}_3}$  are the lifetimes with respect to photolysis, reaction with the OH radical, reaction with the  $\text{NO}_3$  radical, and reaction with  $\text{O}_3$ , respectively. In turn, these reactive loss processes are determined by the rate constants,  $k_x$ , for reaction and the ambient atmospheric concentrations,  $[X]$ , of the reactive intermediates. For example

$$\frac{1}{\tau_{\text{OH}}} = k_{\text{OH}}[\text{OH}]$$

For photolysis, the photolysis rate depends on the absorption cross section ( $\sigma$ ), the photolysis quantum yield ( $\phi$ ), and the radiation intensity ( $J$ ), all of which are wavelength dependent:

$$\frac{1}{\tau_{\text{photolysis}}} = k_{\text{photolysis}} = \int_{\sim 290 \text{ nm}}^{\sim 800 \text{ nm}} \sigma_{\lambda} \phi_{\lambda} J_{\lambda} d\lambda$$

In this report, the available literature data concerning the chemical and physical removal processes for acetaldehyde ( $\text{CH}_3\text{CHO}$ ) are presented and its resulting atmospheric lifetime is estimated. The atmospheric chemistry of this compound is also discussed. Although acetaldehyde is gaseous at room temperature, its dissolution into the aqueous phase and subsequent aqueous-phase reactions must also be considered in addition to the gas-phase processes.

## B. Chemical Loss Processes

### 1. Reaction with Ozone

Rate constants for the gas-phase reaction of  $\text{O}_3$  with acetaldehyde have been reported by Stedman and Niki (1973) and Atkinson et al. (1981). Both studies were carried out at room temperature using essentially the same experimental technique. However, the rate constants obtained differ substantially, with Stedman and Niki (1973) reporting a rate constant of  $k_1 = (3.4 \pm 0.5) \times 10^{-20} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at  $298 \pm 2 \text{ K}$  and Atkinson et al. (1981) an upper limit to the rate constant of  $k_1 \leq 6 \times 10^{-21} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at  $296 \pm 2 \text{ K}$ . Since only upper limits to the room temperature rate constants of  $k < 10^{-20} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  have been obtained for the aldehydes  $\text{HCHO}$ ,  $(\text{CHO})_2$  and  $\text{CH}_3\text{COCHO}$  (Atkinson and Carter, 1984), Atkinson and Carter (1984) recommended that

$$k_1(\text{CH}_3\text{CHO}) < 1 \times 10^{-20} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } 298 \text{ K}$$

### 2. Reaction with the Hydroxyl Radical

The available literature data concerning the kinetics and mechanism of the reaction of the OH radical with  $\text{HCHO}$  have been reviewed and evaluated by Atkinson (1989). The available kinetic data are given in Table 1, and those of Morris et al. (1971), Morris and Niki (1971), Niki et al. (1978), Atkinson and Pitts (1978), Kerr and Sheppard (1981), Semmes et al. (1985) and Michael et al. (1985) are plotted in Arrhenius form in Figure 1. Within the cited experimental errors, the room

Table 1. Rate Constants  $k_2$  and Temperature-Dependent Parameters ( $k_2 = Ae^{-B/T}$ ) for the Gas-Phase Reaction of the OH Radical with Acetaldehyde

$10^{12} \times A$ ( $\text{cm}^3 \text{ mole}^{-1} \text{ s}^{-1}$ )	B (K)	$10^{12} \times k_2$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
6.87	-257 $\pm$ 151	15 $\pm$ 3.8	300	DF-MS	Morris et al. (1971)
		15.3	298	DF-MS	Morris and Miki (1971)
		$\leq 18$	295 $\pm$ 2	RR [relative to k(HOMO) $= 4.80 \times 10^{-12} \text{ s}^{-1}$ ]	Cox et al. (1976)
		23.0	1100	RR [relative to k(CO) $= 3.04 \times 10^{-13} \text{ s}^{-1}$ ]	Colket et al. (1977)
		16.2 $\pm$ 1.8	298 $\pm$ 2	RR [relative to k(ethene) $\cdot 10^{-12} \text{ s}^{-1}$ $8.52 \times 10^{-12} \text{ s}^{-1}$ ]	Miki et al. (1978)
7.1 $\pm$ 0.2	-165 $\pm$ 91	16.0 $\pm$ 1.6	299.4	FP-RF	Atkinson and Pitts (1978)
		14.4 $\pm$ 1.5	355.0		
		12.4 $\pm$ 1.3	426.1		
		12.8 $\pm$ 4.3	298 $\pm$ 4	RR [relative to k(ethene) $\cdot 10^{-12} \text{ s}^{-1}$ $8.52 \times 10^{-12} \text{ s}^{-1}$ ]	Kerr and Sheppard (1981)
				FP-RF	Somay et al. (1985)
5.52 $\pm$ 0.80	-307 $\pm$ 52	14.0 $\pm$ 3.1	253	DF-RF	Michael et al. (1985)
		12.2 $\pm$ 2.7	298		
		10.7 $\pm$ 2.3	356		
		11.0 $\pm$ 2.3	424		
		21.0 $\pm$ 1.4	244		
		19.2 $\pm$ 0.6	244		
		18.9 $\pm$ 1.4	259		
		17.9 $\pm$ 1.2	259		
		15.6 $\pm$ 0.8	273		
		16.3 $\pm$ 1.2	273		
		17.8 $\pm$ 0.6	273		
		19.6 $\pm$ 1.2	273		
		14.2 $\pm$ 1.0	298		
		14.7 $\pm$ 2.8	298		
		13.0 $\pm$ 1.2	333		
		14.0 $\pm$ 0.8	355		
		14.0 $\pm$ 0.4	367		
		14.3 $\pm$ 1.0	367		
		15.0 $\pm$ 1.0	373		
		11.6 $\pm$ 0.6	393		
		11.7 $\pm$ 0.8	420		
		10.6 $\pm$ 0.6	424		
		11.0 $\pm$ 0.6	433		
		11.5 $\pm$ 0.8	466		
		10.4 $\pm$ 0.4	468		
		10.4 $\pm$ 0.4	492		
		9.2 $\pm$ 1.4	499		
		9.9 $\pm$ 0.4	528		

<sup>a</sup>From Atkinson (1989).




Figure 1. Arrhenius plot of rate constants for the reaction of the OH radical with acetaldehyde. (x) Morris et al., (1971) and Morris and Niki (1971); ( □ ) Niki et al. (1978); ( ● ) Atkinson and Pitts (1978); ( ▽ ) Kerr and Sheppard, (1981); ( ▲ ) Semmes et al., (1985); ( ○ ) Michael et al., (1985); (—) recommendation (see text).

temperature rate constants of Morris et al. (1971), Morris and Niki (1971), Niki et al. (1978), Atkinson and Pitts (1978), Kerr and Sheppard (1981), Semmes et al. (1985) and Michael et al. (1985) are in reasonably good agreement, although those of Kerr and Sheppard (1981) and Semmes et al. (1985) are somewhat lower than the remaining data. The Arrhenius plot (Fig. 1) does not show any definitive evidence of curvature, and hence the experimental data were fitted by Atkinson (1989) to the Arrhenius expression  $k = Ae^{-B/T}$ . A unit-weighted least-squares analysis of the rate constants of Niki et al. (1978), Atkinson and Pitts (1978) and Michael et al. (1985) leads to the recommendation of

$$k_2(\text{CH}_3\text{CHO}) = (5.55^{+0.76}_{-0.66}) \times 10^{-12} e^{(311 \pm 42)/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

over the temperature range 244-528 K, where the indicated errors are two least-squares standard deviations, and

$$k_2(\text{CH}_3\text{CHO}) = 1.58 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

at 298 K, with an estimated overall uncertainty at 298 K of  $\pm 20\%$ .

Using the three-parameter expression  $k = CT^2e^{-D/T}$ , a unit-weighted least-squares analysis of these same kinetic data (Niki et al., 1978; Atkinson and Pitts, 1978; Michael et al., 1985) yields

$$k_2(\text{CH}_3\text{CHO}) = (6.03^{+1.07}_{-0.92}) \times 10^{-18} T^2 e^{(999 \pm 54)/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

over the temperature range 244-528 K, where the indicated errors are again the two least-squares standard deviations, and


$$k_2(\text{CH}_3\text{CHO}) = 1.53 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$



at 298 K. Over a wider temperature range extending to temperatures  $\geq 600$  K, curvature in the Arrhenius plot is expected, and the above three-parameter expression should probably then be used.

The recent rate constants of Semmes et al. (1985) were not included in the evaluation of this rate constant since they reported difficulties in adequately determining the acetaldehyde concentrations in their reactant mixtures.

While definite product and mechanistic data are not available for the OH radical reaction with acetaldehyde, the observation of peroxyacetyl nitrate (PAN) from the reaction of the OH radical with  $\text{CH}_3\text{CHO}$  in air in the presence of  $\text{NO}_x$  (Atkinson and Lloyd, 1984) shows that at room temperature this reaction must proceed via overall H-atom abstraction from the  $-\text{CHO}$  group.




This is consistent with the observation that the room temperature rate constants for the  $\geq \text{C}_2$  aldehydes are reasonably similar, increasing only slightly with the length of the alkyl side chain (Atkinson, 1989) and showing that the alkyl substituent group has only a minimal effect on the OH radical rate constant. As for formaldehyde, the observed negative temperature dependence suggests that, although the reaction proceeds by overall H-atom abstraction, the reaction involves initial OH radical addition followed by rapid decomposition of the adduct to the observed products. H-atom abstraction from the  $-\text{CH}_3$  group is expected to be of minimal importance at room temperature, (Atkinson, 1987).

### 3. Reaction with the $\text{NO}_3$ Radical

Kinetic data for the gas-phase reaction of the  $\text{NO}_3$  radical with  $\text{CH}_3\text{CHO}$  have been obtained by Morris and Niki (1974), Atkinson et al. (1984), Cantrell et al. (1986) and Dlugokencky and Howard (1989). The studies of Morris and Niki (1974), Atkinson et al. (1984) and Cantrell et

al. (1986) were relative rate measurements carried out at room temperature, with the rate constants for the  $\text{NO}_3$  radical reaction with  $\text{CH}_3\text{CHO}$  being relative to the equilibrium constant  $K$  for the reactions



The study of Dlugokencky and Howard (1989) was an absolute measurement carried out using a flow system with laser-induced fluorescence detection of the  $\text{NO}_3$  radical. The available rate data are given in Table 2, with the relative rate data from Morris and Niki (1974), Atkinson et al. (1984) and Cantrell et al. (1986) being re-evaluated by use of an equilibrium constant of  $K = 1.26 \times 10^{-27} e^{11275/T} \text{ cm}^3 \text{ molecule}^{-1}$  ( $K = [\text{N}_2\text{O}_5]/\{[\text{NO}_2][\text{NO}_3]\}$ ).

These re-evaluated room temperature rate constants of Morris and Niki (1974), Atkinson et al. (1984) and Cantrell et al. (1986) are in good agreement and are in excellent agreement with the absolute room temperature rate constant determined by Dlugokencky and Howard (1989). Since the room temperature rate constants derived from the relative rate studies (Morris and Niki, 1974; Atkinson et al., 1984; Cantrell et al., 1986) have significant uncertainties associated with them because of the uncertainties in the equilibrium constant  $K$  for the  $\text{NO}_3 + \text{NO}_2 \rightleftharpoons \text{N}_2\text{O}_5$  reactions, the absolute data of Dlugokencky and Howard (1989) are used to obtain the recommendation of

$$k_3(\text{CH}_3\text{CHO}) = 1.4 \times 10^{-12} e^{-1860/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

over the temperature range 264 - 274 K, and

$$k_3(\text{CH}_3\text{CHO}) = 2.7 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } 298 \text{ K.}$$

The recent IUPAC evaluation (Atkinson et al., 1989) recommends an identical rate expression.

As is the case for the OH radical reaction, this  $\text{NO}_3$  radical reaction proceeds by H atom abstraction from the  $-\text{CHO}$  group (Cantrell et al., 1986)




Table 2. Room Temperature Rate Constants,  $k_3$ , for the Gas-Phase Reaction of the  $\text{NO}_3$  Radical with Acetaldehyde


$10^{15} \times k_3$ ( $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ )	at T (K)	Reference
$2.54 \pm 0.64^a$	300	Morris and Niki (1974)
$2.44 \pm 0.52^a$	$298 \pm 1$	Atkinson et al. (1984)
$3.06 \pm 0.59^a$	$299 \pm 1$	Cantrell et al. (1986)
$1.26 \pm 0.15$	264	Dlugokencky and Howard (1989) <sup>b</sup>
$2.74 \pm 0.33$	298	
$5.27 \pm 0.63$	332	
$10.0 \pm 1.2$	374	

<sup>a</sup>Relative to an equilibrium constant of  $K = 1.26 \times 10^{-27} e^{11275/T} \text{ cm}^3 \text{ molecule}^{-1}$ , which is derived from the experimental data of Tuazon et al. (1984), Kircher et al. (1984), Burrows et al. (1985) and Cantrell et al. (1988).



<sup>b</sup> $\text{NO}_3$  radicals monitored directly by laser induced fluorescence, with a temperature dependence of  $k_3 = 1.44 \times 10^{-12} e^{-(1860 \pm 300)/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ .

#### 4. Reaction with the $\text{HO}_2$ Radical

While the gas-phase reaction of the  $\text{HO}_2$  radical with formaldehyde is well documented (see, for example, Atkinson et al., 1989), less definitive data are available concerning the rate constant and reaction products of the analogous  $\text{HO}_2$  radical reaction with  $\text{CH}_3\text{CHO}$ . Moortgat and McQuigg (1984) used Fourier transform infrared absorption spectroscopy to study the photooxidation of  $\text{CH}_3\text{CHO}$  in air, and ascribed the formation of  $\text{CH}_3\text{COOH}$  to the reaction of the  $\text{HO}_2$  radical with  $\text{CH}_3\text{CHO}$ . The initially formed  $\text{CH}_3\text{CH}(\text{OOH})\dot{\text{O}}$  radical was expected to rapidly isomerize (Moortgat and McQuigg, 1984),



followed by



Based upon the time-concentration profiles of the reactants and products and unpublished rate data of Barnes and coworkers for a series of  $\alpha$ -dicarbonyls, Moortgat and McQuigg (1984) derived a rate constant for the reaction of  $\text{HO}_2$  with  $\text{CH}_3\text{CHO}$  at 298 K of  $k_{14} = 1 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , and a rate constant for the unimolecular decomposition of the  $\text{CH}_3\text{CH}(\text{OOH})\dot{\text{O}}$  radical back to reactants of  $k_{-14} = 1.5 \text{ s}^{-1}$ . Moortgat and McQuigg (1984) also cited an unpublished rate constant of Barnes and coworkers of  $k_{14} = 1 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  for the  $\text{HO}_2$  radical reaction with  $\text{CH}_3\text{CHO}$ , based upon the decay rate of  $\text{HO}_2$  in the presence of  $\text{CH}_3\text{CHO}$ . However, reformation of the reactants from the initially formed  $\text{CH}_3\text{CH}(\text{OOH})\dot{\text{O}}$  radical [reaction (-4)] would lead to this measured rate constant being a lower limit to that for the initial addition reaction.

Moortgat and coworkers (Moortgat et al., 1987) have carried out further studies of the photolysis of  $\text{CH}_3\text{CHO}$ , and derived rate constants at 298 K of  $k_{14} = 1 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  and  $k_{-14} = 100 \text{ s}^{-1}$ . With these rate constants, the reaction of the  $\text{HO}_2$  radical with  $\text{CH}_3\text{CHO}$  is not expected to be a significant loss process for  $\text{CH}_3\text{CHO}$  under tropospheric conditions (see Section D below).

### 5. Photolysis

The absorption cross-sections and quantum yields for the photolysis of  $\text{CH}_3\text{CHO}$  have most recently been reviewed and evaluated by the IUPAC data panel (Atkinson et al., 1989). This evaluation accepts the earlier evaluation of Baulch et al. (1984). The recommended absorption cross-sections were obtained from the experimental measurements of Calvert and Pitts (1966) and Weaver et al. (1976/77). The recommended quantum yields for the processes,




derived from the studies of Horowitz and Calvert (1982), Horowitz et al. (1982) and Meyrahn et al. (1982), are given in Table 3 together with the recommended absorption cross-sections. At wavelengths >290 nm the other possible photodissociation processes



are of negligible importance (Atkinson and Lloyd, 1984; Baulch et al., 1984).

### C. Physical Loss Processes

Acetaldehyde can also dissolve in aqueous solutions, and hence the possibility that  $\text{CH}_3\text{CHO}$  is incorporated into cloud, rain and fog water, with subsequent aqueous-phase reactions, needs to be considered. As discussed by Betterton and Hoffmann (1988),  $\text{CH}_3\text{CHO}$  can exist in aqueous solution as the hydrated, gem-diol, form



Betterton and Hoffmann (1988) measured the apparent Henry's law coefficient,  $H^*$ , where

$$H^* = \{[\text{CH}_3\text{CHO}(\text{aq})] + [\text{CH}_3\text{CH}(\text{OH})_2(\text{aq})]\} / [\text{CH}_3\text{CHO}(\text{gas})]$$

as a function of temperature over the range 278-308 K. At 298 K the value of  $H^*$  determined was  $H^* = 11.4 \pm 0.4 \text{ mol atm}^{-1}$ . The degree of hydration for  $\text{CH}_3\text{CHO}$  is fairly small, with (Betterton and Hoffmann, 1988)  $K_{\text{hyd}} = [\text{CH}_3\text{CH}(\text{OH})_2(\text{aq})] / [\text{CH}_3\text{CHO}(\text{aq})] = 1.4$  at 298 K (which can be contrasted to formaldehyde, for which  $K_{\text{hyd}} = 2.3 \times 10^3$  at 298 K).

For gas-phase organic compounds which are very efficiently rained out (and for particles, which are also very efficiently rained out), the washout ratio  $W$ , where  $W = \text{concentration in rain} / \text{concentration in air}$  ( $W = RTH^*$  with the definition of  $H^*$  given above), is  $10^5$  to  $10^6$  (Eisenreich et al., 1981; Atkinson, 1988). The value of  $H^*$  of Betterton and Hoffmann

Table 3. Absorption Cross Sections,  $\sigma$ , and Quantum Yields  $\Phi$  (for 1 atm air) for the Photolysis of  $\text{CH}_3\text{CHO}$  (from Atkinson et al., 1989)

Wavelength $\lambda$ (nm)	$10^{20} \sigma$ ( $\text{cm}^2$ )	$\Phi_{5a}$	$\Phi_{5b}$
260	2.0	0.46	0.31
270	3.4	0.31	0.39
280	4.5	0.05	0.58
290	4.9	0.01	0.53
295	4.5	0.00	0.48
300	4.3		0.43
305	3.4		0.37
315	2.1		0.17
320	1.8		0.10
325	1.1		0.04
330	0.69		0.00
335	0.38		
340	0.15		
345	0.08		

(1988) corresponds to a washout ratio of  $W = 280$  at 298 K, in good agreement with the value of 370 reported earlier by Buttery et al. (1969). With a washout ratio of this magnitude, wet deposition of  $\text{CH}_3\text{CHO}$  is expected to be of minor significance as a tropospheric loss process for  $\text{CH}_3\text{CHO}$ , especially compared to the gas-phase processes.


#### D. Tropospheric Lifetime and Fate of Acetaldehyde

As discussed above, the tropospheric lifetime of  $\text{CH}_3\text{CHO}$  is determined from the loss rates of both the chemical and physical loss processes. Based upon the rate constants given above for the gas-phase reactions of  $\text{CH}_3\text{CHO}$  with OH,  $\text{NO}_3$  and  $\text{HO}_2$  radicals and with  $\text{O}_3$ , and measured or estimated tropospheric concentrations of these reactive species of: OH radicals,  $1.5 \times 10^6$  molecule  $\text{cm}^{-3}$  during daylight hours (Prinn et al., 1987);  $\text{NO}_3$  radicals,  $2.4 \times 10^8$  molecule  $\text{cm}^{-3}$  during nighttime hours (Platt et al., 1984; Atkinson et al., 1986);  $\text{HO}_2$  radicals,  $\sim 10^7$  molecule  $\text{cm}^{-3}$  for conditions applicable to polluted areas (Hard et al., 1984; Zellner and Weibring, 1989); and  $\text{O}_3$ ,  $7 \times 10^{11}$  molecule  $\text{cm}^{-3}$  throughout a 24-hr period


(Logan, 1985), the tropospheric loss rates of  $\text{CH}_3\text{CHO}$  due to these chemical processes can be calculated. The calculated lifetimes due to these individual reactions are then: reaction with the OH radical, 12 hours (i.e., 1 day since the OH radical reaction occurs only during daylight); reaction with the  $\text{NO}_3$  radical, 35 days; reaction with the  $\text{HO}_2$  radical (neglecting the back-decomposition reaction of the  $\text{CH}_3\text{CH}(\text{OOH})\text{O}^\cdot$  radical), ~3 years; and reaction with  $\text{O}_3$ , >4.5 years. Since the back-decomposition of the  $\text{CH}_3\text{CH}(\text{OOH})\text{O}^\cdot$  radical formed from the  $\text{HO}_2$  radical reaction significantly decreases the importance of the  $\text{HO}_2$  radical reaction as a  $\text{CH}_3\text{CHO}$  loss process, the above calculated lifetimes of  $\text{CH}_3\text{CHO}$  due to reaction with the  $\text{HO}_2$  radical and with  $\text{O}_3$  show that these reactions will be totally negligible as a tropospheric loss process for  $\text{CH}_3\text{CHO}$ . Clearly, the daytime OH radical reaction is the dominant of these chemical loss processes.

Photolysis of  $\text{CH}_3\text{CHO}$  is estimated to lead to a lifetime (due to photolysis) of ~6.6 day (~80 hrs) in the lower troposphere for a zenith angle of  $0^\circ$  (Horowitz and Calvert, 1982; Meyrahn et al., 1982). Thus, reaction with the OH radical and, to a lesser extent, photolysis are the major tropospheric chemical loss processes, with an overall lifetime of  $\text{CH}_3\text{CHO}$  of ~12 hours (or 1 day) and with the OH radical reaction being the dominant removal route. Wet deposition will, on average, lead to a shorter tropospheric lifetime, but since wet deposition is episodic in nature, and expected to be of minor importance due to the relatively low washout ratio, the above lifetime calculated from the OH radical reaction can be regarded as being the tropospheric lifetime, with a somewhat faster removal of  $\text{CH}_3\text{CHO}$  from the lower troposphere during rain and/or fog events (see Grosjean and Wright, 1983). It must also be noted that reaction with the OH radical and photolysis only occur during daytime hours.



The reactions subsequent to the OH radical reaction are as discussed below. The initial H atom abstraction reaction







forms the acetyl radical, which rapidly (and exclusively under tropospheric conditions) adds  $\text{O}_2$  to yield the acetyl peroxy radical (Atkinson, 1990).



The acetyl peroxy radical reacts with NO and NO<sub>2</sub>




with rate constants of  $\sim 5.1 \times 10^{-12} e^{200/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  ( $1.0 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K) and  $\sim 7.2 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  (at 298 K and 760 Torr total pressure of air), respectively (Atkinson, 1990). Additionally, in the "clean" troposphere, the reactions of the CH<sub>3</sub>C(O)OO· radical with HO<sub>2</sub> and CH<sub>3</sub>Ö<sub>2</sub> radicals may be important,




with  $k_a/k_b \sim 3$  (Niki et al., 1985) and  $k_c \sim k_d \sim 5.5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K (Moortgat et al., 1989). The  $\dot{\text{C}}\text{H}_3$  radical produced in reaction pathway (c) and from the reaction of the CH<sub>3</sub>C(O)OO· radical with NO ultimately forms HCHO (Ravishankara, 1988; Atkinson, 1990) by the




following series of reactions




Peroxyacetyl nitrate (PAN) acts as a temporary reservoir of the  $\text{CH}_3\text{C}(\text{O})\text{OO}\dot{\text{O}}$  radical, with a lifetime with respect to thermal decomposition of ~33 min at 298 K and 760 Torr total pressure of air (Atkinson et al., 1989; Atkinson, 1990).




As noted above, PAN formation occurs by the reaction of the acetyl peroxy radical with  $\text{NO}_2$ , and the organic precursor to PAN formation is the acetyl radical,  $\text{CH}_3\dot{\text{C}}\text{O}$ . The  $\text{CH}_3\dot{\text{C}}\text{O}$  radical is formed from the OH and  $\text{NO}_3$  radical reactions with acetaldehyde,




from the photolysis of  $\alpha$ -dicarbonyls such as methylglyoxal and biacetyl [which are formed during the  $\text{NO}_x$ -air photooxidations of toluene, the xylenes and the trimethylbenzenes (Atkinson, 1990)],



and from the decomposition reactions of certain  $\beta$ -carboxyalkoxy radicals of the structure  $\text{CH}_3\text{COC}(\text{O}^\cdot)\text{R}_1\text{R}_2$ . These  $\text{CH}_3\text{COC}(\text{O}^\cdot)\text{R}_1\text{R}_2$  radicals are formed subsequent to OH radical reaction with 2-ketones and certain  $\alpha,\beta$ -unsaturated ketones; for example





(which also forms acetaldehyde)




Thus, acetaldehyde photooxidation is not the only route to PAN formation, although it is expected from the ambient air data of Grosjean (1982) and Grosjean et al. (1983) that the major precursors to PAN formation will be acetaldehyde and the methyl-substituted benzenes.

#### E. Atmospheric Formation of Acetaldehyde

Acetaldehyde is formed in the troposphere from the tropospheric degradation reactions of many organic compounds. In the "clean" troposphere,  $\text{CH}_3\text{CHO}$  is formed as a product of the atmospheric reactions of ethane



[involving rapid formation of  $\text{CH}_3\text{CH}_2\text{O}_2^\cdot$ , followed by reactions of the ethyl peroxy radical with  $\text{NO}$ ,  $\text{NO}_2$ ,  $\text{HO}_2$  radicals and other organic peroxy radicals to ultimately lead to the formation of the ethoxy ( $\text{CH}_3\text{CH}_2\text{O}^\cdot$ ) radical (Atkinson, 1990)], followed by the reaction




The steady state concentration of acetaldehyde is then approximately given by


$$[\text{CH}_3\text{CHO}] = \frac{k_{\text{OH} + \text{C}_2\text{H}_6}}{k_{\text{OH} + \text{CH}_3\text{CHO}}} [\text{C}_2\text{H}_6]$$

and for a northern hemisphere ethane mixing ratio of ~1 ppb (Blake and Rowland, 1986; Singh et al., 1988) the acetaldehyde mixing ratio will be ~0.02 ppb.


The formation of  $\text{CH}_3\text{CH}_2\text{O}^\cdot$  and/or  $\text{C}_2\text{H}_5^\cdot$  radicals during the atmospheric degradation reactions of organic compounds thus lead to the formation of  $\text{CH}_3\text{CHO}$ . Sources of  $\text{C}_2\text{H}_5^\cdot$  radicals include the reaction of propionaldehyde with the OH radical




and the decomposition reactions of the more complex alkoxy radicals; for example, of the 2-butoxy radical formed from the OH radical-initiated reaction with n-butane




This and analogous alkoxy radical decompositions lead directly to  $\text{CH}_3\text{CHO}$ . The  $\beta$ -hydroxyalkoxy radicals formed subsequent to OH radical addition to propene and certain 2-alkenes decompose to form  $\text{CH}_3\text{CHO}$  or the  $\alpha$ -hydroxy radical  $\text{CH}_3\dot{\text{C}}\text{HOH}$  (Atkinson and Lloyd, 1984; Atkinson, 1990).



The  $\alpha$ -hydroxy radical  $\text{CH}_3\dot{\text{C}}\text{HOH}$  forms acetaldehyde by reaction with  $\text{O}_2$  (Atkinson and Lloyd, 1984; Atkinson, 1990)



In addition, the gas-phase reactions of  $\text{O}_3$  with propene and certain 2-alkenes leads to the formation of  $\text{CH}_3\text{CHO}$ . For example, the reaction of  $\text{O}_3$  with propene at room temperature and atmospheric pressure is expected to lead to the products (Atkinson, 1990):



Thus, the atmospheric formation of  $\text{CH}_3\text{CHO}$  from organic precursors occurs in polluted areas characteristic of urban areas, and in many cases this atmospheric formation of  $\text{CH}_3\text{CHO}$  dominates over direct emission of  $\text{CH}_3\text{CHO}$  from combustion sources (Grosjean et al., 1983).

#### F. Ambient Atmospheric Concentrations of $\text{CH}_3\text{CHO}$

The  $\text{CH}_3\text{CHO}$  mixing ratio in the clean lower troposphere is expected to be low, and using a  $\text{C}_2\text{H}_6$  mixing ratio of ~1 ppb the  $\text{CH}_3\text{CHO}$  mixing ratio is estimated to be ~0.02 ppb (see Section E above). Reported data for the ambient concentrations of  $\text{CH}_3\text{CHO}$  in California since 1980 are given in Table 4. These data were all collected in the Los Angeles air basin, and vary from ~0.1 ppb up to ~50 ppb, with the most recent concentrations measured in the California Air Resources Board-funded intercomparison study at Claremont in the Los Angeles air basin being significantly lower at <10 ppb. Further, as yet unpublished, ambient  $\text{CH}_3\text{CHO}$  data are available from the 1987 ARB-funded South Coast Air Quality Study (SCAQS).

It may be expected that the ambient atmospheric  $\text{CH}_3\text{CHO}$  levels encountered in the Los Angeles air basin are among the highest to be encountered in California, and hence the ambient atmospheric  $\text{CH}_3\text{CHO}$  mixing ratios in California range from <0.1 ppb (for clean tropospheric air) up to a few tens of ppb, with the actual values depending on the organic precursor emissions, the extent of photochemical activity and the meteorological conditions pertaining.

Table 4. Recent Measurements of Acetaldehyde Concentrations in Ambient Air in California

CH <sub>3</sub> CHO Mixing Ratio (ppb)	Measurement Date and Place	Reference
0-32	5-6/1980, Los Angeles	Grosjean (1982)
3-35	9-10/1980, Claremont	Grosjean (1982)
13-38	7-10/1980, Los Angeles	Grosjean et al. (1983)
8-16	7-8/1980, Burbank	Grosjean et al. (1983)
13	7/1980, Pasadena	Grosjean et al. (1983)
3	8/1980, Pacoima	Grosjean et al. (1983)
8	8/1980, Newhall	Grosjean et al. (1983)
16-41	9-10/1980, Rosemead	Grosjean et al. (1983)
56	9/1980, Covina	Grosjean et al. (1983)
38	9/1980, Cucamonga	Grosjean et al. (1983)
25-34	10/1980, El Monte	Grosjean et al. (1983)
15	10/1980, San Dimas	Grosjean et al. (1983)
19	10/1980, Upland	Grosjean et al. (1983)
1.5-10	10/1980, Azusa	Grosjean et al. (1983)
0.1-2.7	10/1980, Lennox	Grosjean et al. (1983)
2-39	9-11/1981, Los Angeles	Grosjean and Fung (1984)
1.0-9.0	9/1985, Claremont	Grosjean (1988)

#### G. Conclusions

Acetaldehyde is both directly emitted into the atmosphere as well as being formed in the atmosphere from the photochemical degradation of non-methane organic compounds. CH<sub>3</sub>CHO is removed from the lower troposphere by reaction with the OH radical and, to a lesser extent, by photolysis, with wet deposition (leading to incorporation of acetaldehyde into rain, cloud and fog water) being expected to be of minor importance. The estimated lifetime of CH<sub>3</sub>CHO due to OH radical reaction is ~1 day with this OH radical reaction leading to the formation of (together with other products) peroxyacetyl nitrate (PAN). Reported ambient atmospheric HCHO concentrations in the Los Angeles air basin (expected to have among the higher levels encountered in California) range up ~50 ppb.

## H. References

- Atkinson, R. and J. N. Pitts, Jr. (1978): Kinetics of the reactions of the OH radical with HCHO and CH<sub>3</sub>CHO over the temperature range 299-426°K. J. Chem. Phys., 68, 3581-3584.
- Atkinson, R., S. M. Aschmann, A. M. Winer and J. N. Pitts, Jr. (1981): Rate constants for the gas-phase reactions of O<sub>3</sub> with a series of carbonyls at 296 K. Int. J. Chem. Kinet., 13, 1133-1142.
- Atkinson, R. and W. P. L. Carter (1984): Kinetics and mechanisms of the gas-phase reactions of ozone with organic compounds under atmospheric conditions. Chem. Rev., 84, 437-470.
- Atkinson, R. and A. C. Lloyd (1984): Evaluation of kinetic and mechanistic data for modeling of photochemical smog. J. Phys. Chem. Ref. Data, 13, 315-444.
- Atkinson, R., C. N. Plum, W. P. L. Carter, A. M. Winer, and J. N. Pitts, Jr. (1984): Rate constants for the gas-phase reactions of nitrate radicals with a series of organics in air at 298 ± 1 K. J. Phys. Chem., 88, 1210-1215.
- Atkinson, R., A. M. Winer and J. N. Pitts, Jr. (1986): Estimation of night-time N<sub>2</sub>O<sub>5</sub> concentrations from ambient NO<sub>2</sub> and NO<sub>3</sub> radical concentrations and the role of N<sub>2</sub>O<sub>5</sub> in night-time chemistry. Atmos. Environ., 20, 331-339.
- Atkinson, R. (1987): A structure-activity relationship for the estimation of rate constants for the gas-phase reactions of OH radicals with organic compounds. Int. J. Chem. Kinet., 19, 799-828.
- Atkinson, R. (1988): Atmospheric transformations of automotive emissions. In: Air Pollution, the Automobile, and Public Health, A. Y. Watson, R. R. Bates and D. Kennedy (eds.), National Academy Press, Washington, D.C., pp. 99-132.
- Atkinson, R. (1989): Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. J. Phys. Chem. Ref. Data, Monograph 1, 1-246.
- Atkinson, R., D. L. Baulch, R. A. Cox, R. F. Hampson, Jr., J. A. Kerr and J. Troe (1989): Evaluated kinetic and photochemical data for atmospheric chemistry: Supplement III. J. Phys. Chem. Ref. Data, 18, 881-1097.
- Atkinson, R. (1990): Gas-phase tropospheric chemistry of organic compounds: a review. Atmos. Environ., 24A, 1-41.
- Baulch, D. L., R. A. Cox, R. F. Hampson, Jr., J. A. Kerr, J. Troe and R. T. Watson (1984): Evaluated kinetics and photochemical data for atmospheric chemistry: Supplement II. J. Phys. Chem. Ref. Data, 13, 1259-1380.

- Betterton, E. A. and M. R. Hoffmann (1988): Henry's Law Constants of some environmentally important aldehydes. *Environ. Sci. Technol.*, 22, 1415-1418.
- Bidleman, T. F. (1988): Atmospheric processes. *Environ. Sci. Technol.* 22, 361-367.
- Blake, D. R. and F. S. Rowland (1986): Global atmospheric concentrations and source strength of ethane. *Nature*, 321, 231-233.
- Burrows, J. P., G. S. Tyndall and G. K. Moortgat (1985): A study of the  $N_2O_5$  equilibrium between 275 and 315 K and determination of the heat of formation of  $NO_3$ . *Chem. Phys. Lett.*, 119, 193-198.
- Buttery, R. G., L. C. Ling and D. G. Guadagni (1969): Volatilities of aldehydes, ketones, and esters in dilute water solution. *J. Agr. Food Chem.*, 17, 385-389.
- Calvert, J. G. and J. N. Pitts, Jr. (1966) Photochemistry, Wiley, New York, N.Y.
- Cantrell, C. A., J. A. Davidson, K. L. Busarow and J. G. Calvert (1986): The  $CH_3CHO-NO_3$  reaction and possible nighttime PAN generation. *J. Geophys. Res.*, 91, 5347-5353.
- Cantrell, C. A., J. A. Davidson, A. H. McDaniel, R. E. Shetter and J. G. Calvert (1988): The equilibrium constant for  $N_2O_5 \rightleftharpoons NO_2 + NO_3$ : absolute determination by direct measurement from 243 to 397 K. *J. Chem. Phys.*, 88, 4997-5006.
- Colket, M. B. III, D. W. Naegeli and I. Glassman (1977): High temperature oxidation of acetaldehyde. 16th Int. Symposium on combustion, 1976; The Combustion Institute, Pittsburgh, PA, pp 1023-1039.
- Cox, R. A., R. G. Derwent, P. M. Holt and J. A. Kerr (1976): Photolysis of nitrous acid in the presence of acetaldehyde. *J. Chem. Soc., Faraday Trans. 1*, 72, 2061-2075.
- Dlugokencky, E. J. and C. J. Howard (1989): Studies of  $NO_3$  radical reactions with some atmospheric organic compounds at low pressures. *J. Phys. Chem.*, 93, 1091-1096.
- Eisenreich, S. J., B. B. Looney and J. D. Thornton, (1981): Airborne organic contaminants in the Great Lakes ecosystem. *Environ. Sci. Technol.*, 15, 30-38.
- Grosjean, D. (1982): Formaldehyde and other carbonyls in Los Angeles ambient air. *Environ. Sci. Technol.*, 16, 254-262.
- Grosjean, D. and B. Wright (1983): Carbonyls in urban fog, ice fog, cloudwater and rainwater. *Atmos. Environ.*, 17, 2093-2096.



- Grosjean, D., R. D. Swanson and C. Ellis (1983): Carbonyls in Los Angeles air: contribution of direct emissions and photochemistry. *Sci. Total Environ.*, 29, 65-85.
- Grosjean, D. and K. Fung (1984): Hydrocarbons and carbonyls in Los Angeles air. *J. Air Poll. Contr. Assoc.*, 34, 537-543.
- Grosjean, D. (1988): Aldehydes, carboxylic acids and inorganic nitrate during NSMCS. *Atmos. Environ.*, 22, 1637-1648.
- Hard, T. M., R. J. O'Brien, C. Y. Chan and A. A. Mehrabzadeh (1984): Tropospheric free radical determination by FAGE. *Environ. Sci. Technol.*, 18, 768-777.
- Horowitz, A. and J. G. Calvert (1982): Wavelength dependence of the primary processes in acetaldehyde photolysis. *J. Phys. Chem.*, 86, 3105-3114.
- Horowitz, A., C. J. Kershner and J. G. Calvert (1982): Primary processes in the photolysis of acetaldehyde at 3000 Å and 25°C. *J. Phys. Chem.*, 86, 3094-3105.
- Kerr, J. A. and D. W. Sheppard (1981): Kinetics of the reactions of hydroxyl radicals with aldehydes studied under atmospheric conditions. *Environ. Sci. Technol.*, 15, 960-963.
- Kircher, C. C., J. J. Margitan and S. P. Sander (1984): Pressure and temperature dependence of the reaction  $\text{NO}_2 + \text{NO}_3 + \text{M} \rightarrow \text{N}_2\text{O}_5 + \text{M}$ . *J. Phys. Chem.*, 88, 4370-4375.
- Logan, J. A. (1985): Tropospheric ozone: seasonal behavior, trends, and anthropogenic influence. *J. Geophys. Res.*, 90, 10463-10482.
- Meyrahn, H., G. K. Moortgat and P. Warneck (1982): The photolysis of acetaldehyde under atmospheric conditions. 15th Informal Conference on Photochemistry, Stanford Univ., Stanford, CA, June 27-July 1.
- Michael, J. V., D. G. Keil and R. B. Klemm (1985): Rate constants for the reaction of hydroxyl radicals with acetaldehyde from 244-528 K. *J. Chem. Phys.*, 83, 1630-1636.
- Moortgat, G. K. and R. D. McQuigg (1984): A FTIR spectroscopic study of the photooxidation of acetaldehyde in air. *Proc. 3rd European Symposium on the Physico-Chemical Behavior of Atmospheric Pollutants*, pp. 194-204, D. Reidel Publ. Co., Dordrecht, Holland.
- Moortgat, G. K., J. P. Burrows, W. Schneider, G. S. Tyndall and R. A. Cox (1987): A study of the  $\text{HO}_2 + \text{CH}_3\text{CHO}$  reaction in the photolysis of  $\text{CH}_3\text{CHO}$ , and its consequences for atmospheric chemistry. *Proc. 4th European Symposium on the Physico-Chemical Behavior of Atmospheric Pollutants*, pp. 271-281, D. Reidel Publ. Co., Dordrecht, Holland.
- Moortgat, G. K., B. Veyret and R. Lesclaux (1989): Absorption spectrum and kinetics of reactions of the acetylperoxy radical. *J. Phys. Chem.*, 93, 2362-2368.

- Morris, E. D., Jr. and H. Niki (1971): Reactivity of hydroxyl radicals with olefins. *J. Phys. Chem.*, 75, 3640-3641.
- Morris, E. D., Jr., D. H. Stedman and H. Niki (1971): Mass spectrometric study of the reactions of the hydroxyl radical with ethylene, propylene, and acetaldehyde in a discharge-flow system. *J. Am. Chem. Soc.*, 93, 3570-3572.
- Morris, E. D., Jr. and H. Niki (1974): Reaction of the nitrate radical with acetaldehyde and propylene. *J. Phys. Chem.*, 78, 1337-1338.
- Niki, H., P. D. Maker, C. M. Savage and L. P. Breitenbach (1978): Relative rate constants for the reaction of hydroxyl radical with aldehydes. *J. Phys. Chem.*, 82, 132-134.
- Niki, H., P. D. Maker, C. M. Savage and L. P. Breitenbach (1985): FTIR study of the kinetics and mechanism for Cl-atom-initiated reactions of acetaldehyde. *J. Phys. Chem.*, 89, 588-591.
- Platt, U. F., A. M. Winer, H. W. Biermann, R. Atkinson and J. N. Pitts, Jr. (1984): Measurement of nitrate radical concentrations in continental air. *Environ. Sci. Technol.*, 18, 365-369.
- Prinn, R., D. Cunnold, R. Rasmussen, P. Simmonds, F. Alyea, A. Crawford, P. Fraser and R. Rosen (1987): Atmospheric trends in methylchloroform and the global average for the hydroxyl radical. *Science*, 238, 945-950.
- Ravishankara, A. R. (1988): Kinetics of radical reactions in the atmospheric oxidation of  $\text{CH}_4$ . *Ann. Rev. Phys. Chem.*, 39, 367-394.
- Semmes, D. H., A. R. Ravishankara, C. A. Gump-Perkins and P. H. Wine (1985): Kinetics of the reactions of hydroxyl radical with aliphatic aldehydes. *Int. J. Chem. Kinet.*, 17, 303-313.
- Singh, H. B., W. Viezee and L. J. Salas (1988): Measurements of selected  $\text{C}_2\text{-C}_5$  hydrocarbons in the troposphere: latitudinal, vertical and temporal variations. *J. Geophys. Res.*, 93, 15861-15878.
- Stedman, D. H. and H. Niki (1973): Ozonolysis rates of some atmospheric gases. *Environ. Lett.*, 4, 303-310.
- Tuazon, E. C., E. Sanhueza, R. Atkinson, W. P. L. Carter, A. M. Winer, and J. N. Pitts, Jr. (1984): Direct determination of the equilibrium constant at 298 K for the  $\text{NO}_2 + \text{NO}_3 \rightleftharpoons \text{N}_2\text{O}_5$  reactions. *J. Phys. Chem.*, 88, 3095-3098.
- Weaver, J., J. Meagher and J. Heicklen (1976/77) Photooxidation of  $\text{CH}_3\text{CHO}$  vapor at 3130 Å, *J. Photochem.*, 6, 111-126.
- Zellner, R. and G. Weibring (1989): Tropospheric  $\text{HO}_x$  and its response to increased hydrocarbon emissions. *Zeit. Phys. Chem. Neue Folge*, 161, S.167-188.

APPENDIX G

ATMOSPHERIC LIFETIME AND FATE OF STYRENE

A Report Submitted in Partial Fulfillment of  
California Air Resources Board Contract No. A732-107

"Lifetimes and Fates of Toxic Air Contaminants in  
California's Atmosphere"

Roger Atkinson

November 1989

Statewide Air Pollution Research Center  
University of California  
Riverside, CA 92521



## TABLE OF CONTENTS

	<u>Page</u>
A. Introduction.....	G-3
B. Chemical Loss Processes.....	G-5
1. Reaction with Ozone.....	G-5
2. Reaction with the OH Radical.....	G-7
3. Reaction with the NO <sub>3</sub> Radical.....	G-9
4. Photolysis.....	G-11
C. Physical Loss Processes.....	G-11
D. Tropospheric Lifetime and Fate of Styrene.....	G-11
E. Atmospheric Formation of Styrene.....	G-12
F. Ambient Concentrations of Styrene.....	G-12
G. Conclusions.....	G-13
H. References.....	G-13



#### A. Introduction

The atmospheric fates of chemical compounds emitted into the troposphere are governed by a number of chemical and physical removal processes. As the result of laboratory studies carried out over the past 20 years, the potentially important gas-phase chemical reaction pathways which must be considered includes:

- Photolysis during daylight hours.
- Reaction with the hydroxyl (OH) radical during daylight hours.
- Reaction with the hydroperoxyl (HO<sub>2</sub>) radical, mainly during afternoon/evening hours.
- Reaction with the nitrate (NO<sub>3</sub>) radical during nighttime hours.
- Reaction with nitrogen dioxide (NO<sub>2</sub>).
- Reaction with ozone (O<sub>3</sub>).
- Reaction with gaseous nitric acid (HNO<sub>3</sub>).

For the great majority of organic compounds, the most important of these remain photolysis and reaction with hydroxyl (OH) and nitrate (NO<sub>3</sub>) radicals and with ozone (O<sub>3</sub>).

However, for a limited number of organic compounds, one or more of the other reactive chemical species in the troposphere may react with organic compounds at significant rates. For example, HO<sub>2</sub> radicals react with formaldehyde and, more slowly, with acetaldehyde and glyoxal; NO<sub>2</sub> reacts with dialkenes; and gaseous HNO<sub>3</sub> reacts with basic compounds such as the amines.

Additionally, for chemical compounds present in the adsorbed phase, photolysis and reaction with O<sub>3</sub>, N<sub>2</sub>O<sub>5</sub>, NO<sub>2</sub>, HNO<sub>3</sub>, HONO, H<sub>2</sub>SO<sub>4</sub> and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) may contribute to their degradation. For certain of these reactions involving adsorbed organics, synergistic interactions may be of importance.

The physical removal processes can be defined in a simplistic manner as accretion (or coagulation) of particles, and dry and wet deposition of gases and particles. Removal of gases and particles at ground surfaces (including snow and other moist surfaces) is referred to as dry deposition, while removal of these species by precipitation is referred to as wet deposition. Our current understanding of these processes, which are

dynamic in nature, must be considered to be semi-quantitative at the present time, and in general there is a lack of experimental data concerning these processes for most organic compounds (Atkinson, 1988; Bidleman, 1988).

The atmospheric lifetime  $\tau$  of a chemical is defined as the time to decay to a concentration of  $1/e$  ( $\approx 0.368$ ) of that initially present. Also,

$$\frac{1}{\tau} = \frac{1}{\tau_{\text{chemical}}} + \frac{1}{\tau_{\text{physical}}} \quad (1)$$

where  $\tau_{\text{chemical}}$  and  $\tau_{\text{physical}}$  are the lifetimes of the chemical with respect to only chemical reaction and only physical loss processes, respectively. These chemical and physical lifetimes are composites of those for the individual loss processes, for example

$$\frac{1}{\tau_{\text{chemical}}} = \frac{1}{\tau_{\text{photolysis}}} + \frac{1}{\tau_{\text{OH}}} + \frac{1}{\tau_{\text{NO}_3}} + \frac{1}{\tau_{\text{O}_3}} \dots \quad (2)$$

where  $\tau_{\text{photolysis}}$ ,  $\tau_{\text{OH}}$ ,  $\tau_{\text{NO}_3}$  and  $\tau_{\text{O}_3}$  are the lifetimes with respect to photolysis, reaction with the OH radical, reaction with the  $\text{NO}_3$  radical, and reaction with  $\text{O}_3$ , respectively. In turn, these reactive loss processes are determined by the rate constants,  $k_x$ , for reaction and the ambient atmospheric concentrations,  $[X]$ , of the reactive intermediates. For example

$$\frac{1}{\tau_{\text{OH}}} = k_{\text{OH}}[\text{OH}]$$

For photolysis, the photolysis rate depends on the absorption cross section ( $\sigma$ ), the photolysis quantum yield ( $\phi$ ), and the radiation intensity ( $J$ ), all of which are wavelength dependent:



$$\frac{1}{\tau_{\text{photolysis}}} = k_{\text{photolysis}} = \int_{290 \text{ nm}}^{800 \text{ nm}} \sigma_{\lambda} \phi_{\lambda} J_{\lambda} d\lambda$$

In this report, the available literature data concerning the chemical and physical removal processes for styrene ( $\text{C}_6\text{H}_5\text{CH}=\text{CH}_2$ ) are presented and its resulting atmospheric lifetime is estimated. The atmospheric chemistry of this compound is also discussed. Styrene is gaseous at room temperature, and hence this report focusses on the potential gas-phase atmospheric loss processes.

## B. Chemical Loss Processes

### 1. Reaction with Ozone

The rate constants determined for the gas-phase reaction of  $\text{O}_3$  with styrene are given in Table 1. All three rate constant determinations have been carried out at room temperature, and the measured rate constants are in reasonable agreement. In particular, the two most recent absolute rate constant determinations of Atkinson et al. (1982, 1989) are in good agreement, and a weighted average of these two rate constants leads to the recommended rate constant of

$$k_1 = 1.8 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

at 298 K, with an estimated overall uncertainty of  $\pm 40\%$ .

The magnitude of this rate constant, when compared to the literature rate constants at room temperature for the gas-phase reactions of  $\text{O}_3$  with benzene and the methyl-substituted benzenes of  $k < 10^{-20} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  (Atkinson and Carter, 1984), shows that the reaction of  $\text{O}_3$  with styrene proceeds by initial addition of  $\text{O}_3$  to the  $-\text{CH}=\text{CH}_2$  substituent group. Product studies using in situ long pathlength Fourier transform infrared (FT-IR) absorption spectroscopy to monitor the reactants and products (Atkinson et al., 1989) showed that the major products of the gas-phase reaction of  $\text{O}_3$  with styrene in one atmosphere of air were formaldehyde ( $\text{HCHO}$ ) and benzaldehyde ( $\text{C}_6\text{H}_5\text{CHO}$ ), together with a minor amount of formic acid ( $\text{HCOOH}$ ). The observed product yields were:  $\text{HCHO}$  and  $\text{C}_6\text{H}_5\text{CHO}$ , each  $\sim 40\%$ ; and  $\text{HCOOH}$ ,  $\sim 1\text{-}2\%$ . Analyses of reaction samples from analogous experiments by gas chromatography (GC) and combined gas chromatography-


Table 1. Rate Constants,  $k_1$  for the Gas-Phase Reaction of  $O_3$  with Styrene

$10^{17} \times k_1$ ( $cm^3$ molecule $^{-1}$ s $^{-1}$ )	at T (K)	Technique <sup>a</sup>	Reference
3.0	303	F-CA	Bufalini and Altshuller (1965)
$2.16 \pm 0.46$	$296 \pm 2$	S-CL	Atkinson et al. (1982)
$1.71 \pm 0.18$	$296 \pm 2$	S-CL	Atkinson et al. (1989)


<sup>a</sup>Symbols: F, flow system; S, static system; CA, chemical analysis by wet chemical methods; CL, ozone monitored by chemiluminescence.

mass spectrometry (GC-MS) yielded the same major products as identified from the FT-IR data.

The available kinetic and product data suggest that the reaction then proceeds by



where  $[ ]^\ddagger$  denotes an initially energy-rich biradical species. The subsequent reactions of the  $[\dot{C}H_2\ddot{O}\ddot{O}]^\ddagger$  radical formed from the  $O_3$  + ethene reaction have been discussed in detail previously (Atkinson and Lloyd, 1984; Atkinson and Carter, 1984; Atkinson, 1990). At atmospheric pressure of air and room temperature




with the thermalized  $\dot{\text{C}}\text{H}_2\text{O}\dot{\text{O}}$  biradical being expected to react with water vapor to form HCOOH under atmospheric conditions. The reactions of the  $[\text{C}_6\text{H}_5\dot{\text{C}}\text{HO}\dot{\text{O}}]^\ddagger$  radical are not presently known. The product data obtained by Atkinson et al. (1989) are reasonably consistent with the above reaction scheme, providing that the  $[\text{C}_6\text{H}_5\dot{\text{C}}\text{HO}\dot{\text{O}}]^\ddagger$  biradical does not produce  $\text{C}_6\text{H}_5\text{CHO}$  in high yield.

## 2. Reaction with the OH Radical

Rate constants for the gas-phase reactions of the OH radical with styrene have been determined at room temperature by Bignozzi et al. (1981) and Atkinson and Aschmann (1988) using relative rate techniques. The rate constants from these two studies, as reevaluated by Atkinson (1989) to be consistent with the most recent data for the rate constants for the reference reactions, are given in Table 2. The agreement is excellent, and a rate constant of

$$k_2 = 5.8 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

was recommended by Atkinson (1989). Analogous to the  $\text{O}_3$  reaction, the magnitude of the rate constant  $k_2$  suggests that the reaction of the OH radical with styrene proceeds by initial OH radical addition to the substituent  $-\text{CH}=\text{CH}_2$  group



The products of the OH radical reaction with styrene in air have been investigated by Bignozzi et al. (1981) and Atkinson et al. (1989). Bignozzi et al. (1981) observed benzaldehyde as the major product, and derived a formation yield of benzaldehyde of  $1.03 \pm 0.15$  from the time-concentration profiles of styrene and benzaldehyde in an irradiated  $\text{NO}_x$  - styrene - air mixture. More recently, Atkinson et al. (1989) have used long pathlength FT-IR absorption spectroscopy to monitor the reactants and

Table 2. Literature Rate Constants  $k_2$  for the Gas-Phase Reaction of the OH Radical with Styrene


$10^{11} \times k_2$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{ s}^{-1}$ )	at T (K)	Technique <sup>a</sup>	Reference
$52 \pm 5$	$298 \pm 2$	RR [relative to $k(\text{OH} + 2,2,4\text{-}$ $\text{trimethylpentane})$ $= 3.68 \times 10^{-12}]^b$	Bignozzi et al. (1981)
$58.1 \pm 1.5$	$296 \pm 2$	RR [relative to $k(\text{OH} + 2\text{-methyl-}$ $1,3\text{-butadiene}) =$ $1.01 \times 10^{-10}]^b$	Atkinson and Aschmann (1988)



<sup>a</sup>Symbols: RR, relative rate.

<sup>b</sup>Recommendations of Atkinson (1989).





products in irradiated  $\text{C}_2\text{H}_5\text{ONO} - \text{NO} - \text{styrene} - \text{air}$  mixtures, and observed  $\text{HCHO}$  and  $\text{C}_6\text{H}_5\text{CHO}$  as the major products, with formation yields of 0.90 and ~0.75, respectively. A further product was observed by Atkinson et al. (1989) which contained a nitrate ( $-\text{ONO}_2$ ) group. Heuss and Glasson (1968) also observed  $\text{HCHO}$  and  $\text{C}_6\text{H}_5\text{CHO}$  as the major products of the  $\text{NO}_x - \text{air}$  photooxidation of styrene.

The products observed in these studies are those expected to arise from the reactions subsequent to the initial OH radical addition to the  $-\text{CH}=\text{CH}_2$  substituent group. Thus, in the presence of NO, and neglecting the formation of organic nitrates from the reaction of the peroxy radicals with NO,





and



### 3. Reaction with the NO<sub>3</sub> Radical

To date there has only been one determination of the rate constant for the gas-phase reaction of the NO<sub>3</sub> radical with styrene. Using a relative rate technique, Atkinson and Aschmann (1988) determined the rate constant  $k_3$  relative to the rate constant for trans-2-butene at  $296 \pm 2$  K and, using a rate constant for the reaction of the NO<sub>3</sub> radical with trans-2-butene of  $(3.87 \pm 0.45) \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  (Ravishankara and Mauldin, 1985; Dlugokencky and Howard, 1989), this leads to


$$k_3 = (1.51 \pm 0.18) \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

at  $296 \pm 2$  K.



The products of the gas-phase reaction of styrene with the NO<sub>3</sub> radical have been investigated by Atkinson et al. (1989) using long path-length FT-IR absorption spectroscopy to monitor the reactants and products. Formaldehyde (HCHO) and benzaldehyde (C<sub>6</sub>H<sub>5</sub>CHO) were observed as products, with equal formation yields of ~0.10-0.12.

Three as yet unidentified nitrogen-containing compounds were the most conspicuous products of these  $\text{NO}_3$  radical reactions with styrene (Atkinson et al., 1989). These products (A, B and C) had infrared absorption bands which indicated the presence of the following groups: A,  $-\text{ONO}_2$  and  $-\text{OONO}_2$ ; B and C,  $-\text{ONO}_2$  and  $>\text{C}=\text{O}$ . Product A was formed in the initial stages of the reaction, and then disappeared rapidly (as expected from the presence of the thermally unstable peroxyxynitrate,  $-\text{OONO}_2$ , group). Product B was more stable than A, but a steady decrease in its concentration with time was also observed (Atkinson et al., 1989). Product C was the most stable of these three nitrogen-containing compounds.


The  $\text{NO}_3$  radical reaction with styrene is expected to proceed by initial  $\text{NO}_3$  radical addition to the  $-\text{CH}=\text{CH}_2$  substituent group:




followed by the reactions (taking the  $\text{C}_6\text{H}_5\dot{\text{C}}\text{HCH}_2\text{ONO}_2$  radical as an example)




(possibly product A)



(where the  $\text{RO}_2\cdot$  radical can include the  $\text{C}_6\text{H}_5\overset{\text{OO}\cdot}{\underset{|}{\text{C}}}\text{HCH}_2\text{ONO}_2$  or  $\text{C}_6\text{H}_5\text{CH}(\text{ONO}_2)\text{CH}_2\text{OO}\cdot$  radical)



(possibly product C)



Analogous intermediate or first-generation products ( $\text{C}_6\text{H}_5\text{CH}(\text{ONO}_2)\text{CH}_2\text{OONO}_2$  and  $\text{C}_6\text{H}_5\text{CH}(\text{ONO}_2)\text{CHO}$ ) are expected to be formed from the  $\text{C}_6\text{H}_5\text{CH}(\text{ONO}_2)\dot{\text{C}}\text{H}_2$  radical.

#### 4. Photolysis

According to Calvert and Pitts (1966), in solution styrene has its largest wavelength absorption cross-section of  $\sim 2 \times 10^{-20} \text{ cm}^2$ . There is no evidence for photolysis of styrene being of importance in the studies of Bignozzi et al. (1981) or Atkinson and Aschmann (1988). In particular, the good agreement between the studies of Bignozzi et al. (1981) and Atkinson and Aschmann (1988) concerning the OH radical reaction rate constant indicates that photolysis was of minor or negligible significance in the  $\text{NO}_x$ -styrene-air irradiations carried out by Bignozzi et al. (1981), in which the light intensity and spectral distribution and OH radical concentration approximated those in ambient air.

#### C. Physical Loss Processes

No data are available for styrene concerning dry and/or wet deposition. Since styrene is a hydrocarbon present in the gas phase, it is expected that dry and wet deposition will be of negligible importance as tropospheric loss processes for styrene, especially when compared to the relatively short tropospheric lifetime calculated from the gas-phase chemical loss process (see Section D below).

#### D. Tropospheric Lifetime and Fate of Styrene

As discussed in Section A, the tropospheric lifetime of styrene is determined from the loss rates of both the chemical and physical loss processes. The physical loss processes are expected to be of no significance, and the tropospheric lifetime of styrene is determined by its reactions with OH and  $\text{NO}_3$  radicals and  $\text{O}_3$ . Using measured or estimated tropospheric concentrations of these species of: OH radicals,  $1.5 \times 10^6$

molecule  $\text{cm}^{-3}$  during daylight hours (Prinn et al., 1987);  $\text{NO}_3$  radicals,  $2.4 \times 10^8$  molecule  $\text{cm}^{-3}$  during nighttime hours (Platt et al., 1984; Atkinson et al., 1986); and  $\text{O}_3$ ,  $7 \times 10^{11}$  molecule  $\text{cm}^{-3}$  throughout a 24-hr period (Logan, 1985), the tropospheric loss rates of styrene due to these chemical reactions can be assessed. The calculated lifetimes for these individual reactions are then: reaction with the OH radical, 3.2 hrs; reaction with the  $\text{NO}_3$  radical, 7.7 hrs; and reaction with  $\text{O}_3$ , 22 hr. The daytime OH radical reaction is the dominant of the reaction loss processes, leading to a tropospheric lifetime of styrene during daylight hours of ~3 hrs.

As discussed in Section B.2 above, the tropospheric degradation of styrene will then lead (by reaction with the OH radical) to the formation of HCHO and  $\text{C}_6\text{H}_5\text{CHO}$  in high yield, and the tropospheric degradation of styrene will lead to the formation in the atmosphere of both formaldehyde and benzaldehyde. It should also be noted that the tropospheric degradation of benzaldehyde will lead to the production in the atmosphere of peroxybenzoyl nitrate [PBzN;  $\text{C}_6\text{H}_5\text{C}(\text{O})\text{OONO}_2$ ] (Atkinson and Lloyd, 1984; Atkinson, 1990).

#### E. Atmospheric Formation of Styrene

As presently understood, there are no in situ atmospheric formation processes leading to styrene from other organics emitted into the troposphere from anthropogenic or biogenic sources.

#### F. Ambient Concentrations of Styrene

Styrene has not been identified in most ambient atmospheric monitoring studies nor in emission studies. However, Grosjean and Fung (1984) report the presence of styrene in 16 ambient air samples collected in downtown Los Angeles in the fall of 1981, with the styrene mixing ratios being in the range 0.5-3 part-per-billion (ppb). Ambient air samples collected in August 1986 in Glendora during the Carbonaceous Species Measurement Comparison Study did not show the presence of styrene (D. Lawson, California Air Resources Board; private communication, 1989).



#### G. Conclusions

Styrene is removed from the troposphere by chemical reaction with OH and NO<sub>3</sub> radicals and O<sub>3</sub>, with the daytime OH radical reaction being expected to dominate and to lead to a styrene lifetime of ~3 hrs during daylight hours. The photooxidation of styrene leads to the production in the atmosphere of formaldehyde (HCHO) and benzaldehyde (C<sub>6</sub>H<sub>5</sub>CHO) in close to unit yield.

Styrene is not formed in the atmosphere from either anthropogenic or biogenic emissions, and ambient air measurement data show that the styrene mixing ratios in the Los Angeles air basin are a few ppb or less.

#### H. References

- Atkinson, R., S. M. Aschmann, D. R. Fitz, A. M. Winer and J. N. Pitts, Jr. (1982): Rate constants for the gas-phase reactions of O<sub>3</sub> with selected organics at 296 K. Int. J. Chem. Kinet., 14, 13-18.
- Atkinson, R. and W. P. L. Carter (1984) Kinetics and mechanisms of the gas-phase reactions of ozone with organic compounds under atmospheric conditions. Chem. Rev., 84, 437-470.
- Atkinson, R. and A. C. Lloyd (1984) Evaluation of kinetic and mechanistic data for modeling of photochemical smog. J. Phys. Chem. Ref. Data, 13, 315-444.
- Atkinson, R., A. M. Winer and J. N. Pitts, Jr. (1986): Estimation of night-time N<sub>2</sub>O<sub>5</sub> concentrations from ambient NO<sub>2</sub> and NO<sub>3</sub> radical concentrations and the role of N<sub>2</sub>O<sub>5</sub> in night-time chemistry. Atmos. Environ., 20, 331-339.
- Atkinson, R. (1988): Atmospheric transformations of automotive emissions. In: Air Pollution, the Automobile, and Public Health. A. Y. Watson, R. R. Bates and D. Kennedy (eds.), National Academy Press, Washington, D.C., pp. 99-132.
- Atkinson, R. and S. M. Aschmann (1988): Kinetics of the reactions of acenaphthene and acenaphthylene and structurally-related aromatic compounds with OH and NO<sub>3</sub> radicals, N<sub>2</sub>O<sub>5</sub> and O<sub>3</sub> at 296 ± 2 K. Int. J. Chem. Kinet., 20, 513-539.
- Atkinson, R., S. M. Aschmann, J. Arey and E. C. Tuazon (1989): Unpublished data, Quarterly report to California Air Resources Board Contract No. A732-107 "Lifetimes and Fates of Toxic Air Contaminants in California's Atmosphere."

- Atkinson, R. (1989): Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. J. Phys. Chem. Ref. Data, Monograph. 1, 1-246.
- Atkinson, R. (1990): Gas-phase tropospheric chemistry of organic compounds: a review. Atmos. Environ., 24A, 1-41.
- Bidleman, T. F. (1988): Atmospheric processes. Environ. Sci. Technol., 22, 361-367.
- Bignozzi, C. A., A. Maldotti, C. Chiorboli, C. Bartocci and V. Carassiti (1981): Kinetics and mechanisms of reactions between aromatic olefins and hydroxyl radicals. Int. J. Chem. Kinet., 13, 1235-1242.
- Bufalini, J. J. and A. P. Altshuller (1965): Kinetics of vapor-phase hydrocarbon-ozone reactions. Can. J. Chem., 43, 2243-2250.
- Calvert, J. G. and J. N. Pitts, Jr. (1966): "Photochemistry," Wiley, New York, NY.
- Dlugokencky, E. J. and C. J. Howard (1989): Studies of NO<sub>3</sub> radical reactions with some atmospheric organic compounds at low pressures. J. Phys. Chem., 93, 1091-1096.
- Grosjean, D. and K. Fung (1984): Hydrocarbons and carbonyls in Los Angeles air. J. Air Pollut. Control Assoc., 34, 537-543.
- Heuss, J. M. and W. A. Glasson (1968): Hydrocarbon reactivity and eye irritation. Environ. Sci. Technol., 2, 1109-1116.
- Logan, J. A. (1985): Tropospheric ozone: seasonal behavior, trends, and anthropogenic influence. J. Geophys. Res., 90, 10463-10482.
- Platt, U. F., A. M. Winer, R. Atkinson and J. N. Pitts, Jr. (1984): Measurement of nitrate radical concentrations in continental air. Environ. Sci. Technol., 18, 365-369.
- Prinn, R., D. Cunnold, R. Rasmussen, P. Simmonds, F. Alyea, A. Crawford, P. Fraser and R. Rosen (1987): Atmospheric trends in methylchloroform and the global average for the hydroxyl radical. Science, 238, 945-950.
- Ravishankara, A. R. and R. L. Mauldin, III (1985): Rate coefficient for the reaction of NO<sub>3</sub> with trans-2-butene. J. Phys. Chem., 89, 3144-3147.

APPENDIX H

**ATMOSPHERIC LIFETIME AND FATE OF DIESEL EXHAUST PARTICLES  
AND GAS- AND PARTICLE-PHASE PAH AND PAH-DERIVATIVES**

A Report Submitted in Partial Fulfillment of  
California Air Resources Board Contract No. A732-107

"Lifetimes and Fates of Toxic Air Contaminants in  
California's Atmosphere"

Roger Atkinson and Janet Arey

March 1990

Statewide Air Pollution Research Center  
University of California  
Riverside, CA 92521



## TABLE OF CONTENTS

	<u>Page</u>
A. Introduction.....	H-3
B. Physical and Chemical Characterization of Diesel Particles and PAH and PAH-Derivatives.....	H-6
1. Particle-Size Distribution.....	H-6
2. PAH and PAH-Derivatives.....	H-6
3. Nitro-PAH.....	H-12
C. Physical Loss Processes.....	H-13
D. Gas-Phase Loss Processes of PAH and PAH-Derivatives.....	H-14
1. Wet Deposition of Gas-Phase PAH and PAH-Derivatives.....	H-14
2. Reaction with the OH Radical.....	H-16
3. NO <sub>3</sub> Radical Reactions.....	H-30
4. O <sub>3</sub> Reactions.....	H-37
5. Photolysis.....	H-37
6. Atmospheric Lifetimes of Gas-Phase PAH and PAH-Derivatives....	H-37
7. Evidence for Atmospheric Transformations of PAH.....	H-41
E. Adsorbed-Phase Loss Processes of PAH and PAH-Derivatives.....	H-44
1. Reaction with O <sub>3</sub> .....	H-45
2. Reaction with Oxides of Nitrogen.....	H-47
3. Photolysis.....	H-49
4. Ambient Air Data.....	H-50
F. Atmospheric Formation Processes.....	H-51
G. References.....	H-52



#### A. Introduction

The atmospheric fates of chemical compounds emitted into the troposphere are governed by a number of chemical and physical removal processes. As the result of laboratory studies carried out over the past 20 years, it is recognized that the potentially important gas-phase chemical reaction pathways include:

- Photolysis during daylight hours
- Reaction with the hydroxyl (OH) radical during daylight hours
- Reaction with the hydroperoxyl ( $\text{HO}_2$ ) radical, mainly during afternoon/evening hours
- Reaction with the nitrate ( $\text{NO}_3$ ) radical during nighttime hours
- Reaction with nitrogen dioxide ( $\text{NO}_2$ )
- Reaction with ozone ( $\text{O}_3$ )
- Reaction with gaseous nitric acid ( $\text{HNO}_3$ )

For the great majority of organic compounds which are present in the gas phase, the most important of these loss processes are photolysis and reaction with hydroxyl (OH) and nitrate ( $\text{NO}_3$ ) radicals and ozone ( $\text{O}_3$ ).

However, for a limited number of gaseous organic compounds, one or more of the other reactive chemical species in the troposphere may react with these organic compounds at significant rates. For example,  $\text{HO}_2$  radicals react with formaldehyde and, more slowly, with acetaldehyde and glyoxal;  $\text{NO}_2$  reacts with conjugated dienes; and gaseous  $\text{HNO}_3$  reacts with basic compounds such as the amines.

For chemical compounds present in the adsorbed phase, photolysis and reaction with  $\text{O}_3$ ,  $\text{N}_2\text{O}_5$ ,  $\text{NO}_2$ ,  $\text{HNO}_3$ ,  $\text{HONO}$ ,  $\text{H}_2\text{SO}_4$  and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) may contribute to their degradation. For certain of these reactions involving adsorbed organics, synergistic interactions may be of importance.

The physical removal processes can be defined in a simplistic manner as accretion (or coagulation) of particles, and dry and wet deposition of gases and particles. Removal of gases and particles at ground surfaces (including snow and other moist surfaces) is referred to as dry deposition, while removal of these species by precipitation is referred to as wet deposition. Our understanding of these processes, which are dynamic

in nature, must be considered to be semi-quantitative at the present time, and in general there is a lack of experimental data concerning these processes for most organic compounds (Atkinson, 1988; Bidleman, 1988).

The atmospheric lifetime  $\tau$  of a chemical is defined as the time to decay to a concentration of  $1/e$  ( $\approx 0.368$ ) of that initially present. Also,

$$\frac{1}{\tau} = \frac{1}{\tau_{\text{chemical}}} + \frac{1}{\tau_{\text{physical}}} \quad (1)$$

where  $\tau_{\text{chemical}}$  and  $\tau_{\text{physical}}$  are the lifetimes of the chemical with respect to only chemical reaction and only physical loss processes, respectively. These chemical and physical lifetimes are composites of those for the individual loss processes, for example

$$\frac{1}{\tau_{\text{chemical}}} = \frac{1}{\tau_{\text{photolysis}}} + \frac{1}{\tau_{\text{OH}}} + \frac{1}{\tau_{\text{NO}_3}} + \frac{1}{\tau_{\text{O}_3}} \dots \quad (2)$$

where  $\tau_{\text{photolysis}}$ ,  $\tau_{\text{OH}}$ ,  $\tau_{\text{NO}_3}$  and  $\tau_{\text{O}_3}$  are the lifetimes with respect to photolysis, reaction with the OH radical, reaction with the  $\text{NO}_3$  radical, and reaction with  $\text{O}_3$ , respectively. In turn, these reactive loss processes are determined by the rate constants,  $k_x$ , for reaction and the ambient atmospheric concentrations,  $[X]$ , of the reactive intermediates. For example

$$\frac{1}{\tau_{\text{OH}}} = k_{\text{OH}}[\text{OH}]$$

For photolysis, the photolysis rate depends on the absorption cross section ( $\sigma$ ), the photolysis quantum yield ( $\phi$ ), and the radiation intensity ( $J$ ), all of which are wavelength dependent:



$$\frac{1}{\tau_{\text{photolysis}}} = k_{\text{photolysis}} = \int_{\sim 290 \text{ nm}}^{\sim 800 \text{ nm}} \sigma_{\lambda} \phi_{\lambda} J_{\lambda} d\lambda$$

In this report, we describe the available literature data concerning the physical and chemical removal processes for the particulate matter and the gas- and particle-phase polycyclic aromatic hydrocarbons (PAH) and PAH-derivatives emitted in diesel exhaust. In addition to particles and gas- and particle-phase PAH and PAH-derivatives, the exhaust from diesel-fueled vehicles is comprised of a wide spectrum of inorganic and organic compounds, including carbon dioxide (CO<sub>2</sub>), carbon monoxide (CO), water vapor (H<sub>2</sub>O), oxides of nitrogen (NO<sub>x</sub>, consisting mainly of NO + NO<sub>2</sub>), formaldehyde (HCHO) and other saturated and unsaturated aldehydes and ketones, alkanes, alkenes and monocyclic aromatic hydrocarbons (see, for example, National Research Council, 1983; Jonsson et al., 1985; Hayano et al., 1985; Schuetzle and Frazier, 1986; Obuchi et al., 1987; Johnston, 1988). Discussion of the lifetimes and fates of all of these emitted species is beyond the scope of this document, which focuses on the following:

- The tropospheric lifetime and fate of particle matter emitted from diesel-fueled vehicles.
- The atmospheric lifetimes and fates of PAH and PAH-derivatives which are present in the atmosphere at least partially in the gas phase.
- The atmospheric lifetimes and fates of PAH and PAH-derivatives which are particle-associated in the atmosphere.

The processes to be considered are then

- Physical loss processes for the particles on which, or in which, the PAH and PAH-derivatives reside.

- The gas-phase tropospheric removal and/or transformation processes for PAH and PAH-derivatives present in the gas-phase.
- Adsorbed-phase reactions of PAH and PAH-derivatives.

B. Physical and Chemical Characterization of Diesel Particles and PAH and PAH-Derivatives

1. Particle-Size Distribution

The particles emitted from diesel-fueled vehicles have mass-median aerodynamic diameters in the range 0.1-0.25  $\mu\text{m}$  (Groblicki and Begeman, 1979; Dolan et al., 1980; National Research Council, 1982; Williams, 1982; Pierson et al., 1983), with more than 75% of the mass being associated with particles of diameter  $<1 \mu\text{m}$  (Pierson et al., 1983). The composition of diesel exhaust particulate is mainly elemental carbon, which is then a good carrier for organic compounds of low volatility which can reside on the particle surface or be included inside the particles. The position of attachment of organic compounds on or in the particles can profoundly affect the atmospheric chemistry and lifetimes of these compounds, since those organic compounds present inside the largely elemental carbon particles are prevented from volatilizing into the gas phase and are protected against photolysis and chemical reaction, while those organic compounds present on the surface of the particles can undergo volatilization into the gas phase and be susceptible to photolysis and/or chemical reaction. As noted by the National Research Council (1983), diesel exhaust particulate is typically 25% extractable into organic solvents, although there is a wide variability in the percent extractable, from 10-90%, depending on the vehicle and operating conditions (see, for example, Williams et al., 1989).

2. PAH and PAH-Derivatives

A chemical analysis of the non-polar and moderately polar fractions of a diesel exhaust sample collected on filters from an exhaust dilution tube is given in Table 1 [adapted from Schuetzle (1983)]. A large proportion of diesel particulate extracts are heavy aliphatic hydrocarbons and it seems generally accepted that these are mainly unburnt fuel or lubricant oil components (Obuchi et al., 1987; Barbella et al.,

Table 1. Analysis of Nonpolar and Moderately Polar Fractions of an Extract of Diesel Particles Collected on Filters using an Exhaust Dilution Tube (from Schuetzle, 1983)

Compound	Approximate Concentration in Oldsmobile Extract (ppm)
<u>Nonpolar fractions</u>	
Phenanthrenes and anthracene	600
Methylphenanthrenes and methylanthracenes	1,400
Dimethylphenanthrenes and dimethylanthracenes	3,200
Pyrene	1,700
Fluoranthene	1,400
Methylpyrenes and methylfluoranthenes	800
Chrysene	100
Cyclopenta[cd]pyrene	20
Benzo[ghi]fluoranthene	100
Benz[a]anthracene	500
Benzo[a]pyrene	40
Other PAHs, heterocyclics	34,000
HCs and alkylbenzenes	510,000
Total nonpolar fractions	556,000
<u>Moderately polar fractions</u>	
<u>PAH ketones</u>	
Fluorenones	4,000
Methylfluorenones	400
Dimethylfluorenones	200
Anthrones and phenanthrones	1,600
Methylanthrones and methylphenanthrones	1,600
Dimethylanthrones and dimethylphenanthrones	1,300
Fluoranthones and pyrones	1,200
Benzanthrones	200
Xanthoness	300
Methylxanthoness	200
Thioxanthoness	1,600
Methylthioxanthoness	900
Total	13,500
<u>PAH carboxaldehydes</u>	
Fluorene carboxaldehydes	1,600
Methyl fluorene carboxaldehydes	400
Phenanthrene and anthracene carboxaldehydes	2,600
Methylanthracene and methylphenanthrene carboxaldehydes	1,600
Dimethylanthracene and dimethylphenanthrene carboxaldehydes	400
Benz[a]anthracene, chrysene, and triphenylene carboxaldehydes	400
Naphthalene dicarboxaldehydes	300
Dimethylnaphthalene carboxaldehydes	300
Trimethylnaphthalene carboxaldehydes	1,000
Pyrene and fluoranthene carboxaldehydes	1,600

Table 1 (continued) - 2

Compound	Approximate Concentration in Oldsmobile Extract (ppm)
Xanthene carboxaldehydes	600
Dibenzofuran carboxaldehydes	400
Total	11,200
<u>PAH acid anhydrides</u>	
Naphthalene dicarboxylic acid anhydrides	2,900
Methylnaphthalene dicarboxylic acid anhydrides	1,000
Dimethylnaphthalene dicarboxylic acid anhydrides	500
Anthracene and phenanthrene dicarboxylic acid anhydrides	600
Total	5,000
<u>Hydroxy PAHs</u>	
Hydroxyfluorene	1,400
Methylhydroxyfluorene	400
Dimethylhydroxyfluorene	1,500
Hydroxyanthracenes and hydroxyphenanthrenes	600
Hydroxymethylanthracenes and hydroxymethylphenanthrenes	900
Hydroxydimethylanthracenes and hydroxydimethylphenanthrenes	1,300
Hydroxyfluorenone	2,100
Hydroxyxanthone	1,300
Hydroxyxanthene	1,000
Total	10,400
<u>PAH quinones</u>	
Fluorene quinones	700
Methylfluorene quinones	600
Dimethylfluorene quinones	500
Anthracene and phenanthrene quinones	1,900
Methylanthracene and methylphenanthrene quinones	2,100
Fluoranthene and pyrene quinones	200
Naphtho[1,8-cd]pyrene 1,3-dione	600
Total	6,500
<u>Nitro-PAHs</u>	
Nitrofluorenes	30
Nitroanthracenes and nitrophenanthrenes	70
Nitrofluoranthenes	5
Nitropyrenes	150
Methylnitropyrenes and methylnitrofluoranthenes	20
Total	270
Other oxygenated PAHs	7,700
PAH carryover from nonpolar fraction	6,000
Phthalates, HC contaminants	31,000
Total, moderately polar fractions	92,000

1989). In addition to a wide range of PAH, with methylated PAH being the most abundant, over one hundred species were identified in the moderately polar fractions of a diesel exhaust extract (Schuetzle et al., 1981). The moderately polar fractions contain mainly PAH derivatives including hydroxy, ketone, quinone, carboxaldehyde, acid anhydride and nitro derivatives of PAH (Schuetzle et al., 1981).

The PAH present in diesel exhaust may be either unburned fuel components or combustion-formed PAH. Some researchers have found the major PAH emissions to be unburned fuel components such as naphthalene, fluorene, phenanthrene and their alkyl derivatives (Williams et al., 1986), while other researchers have found the light naphthalenes present in the fuel to be absent from the exhaust (Barbella et al., 1989). Certain PAH, when added to a hexadecane fuel, have been found to promote soot formation and the emissions of other PAH, while the addition of pyrene or phenanthrene increased the emission of these PAH, but caused no increased soot formation (Henderson et al., 1984). There is some evidence that benzo[a]pyrene and other PAH may be formed during the combustion process (Henderson et al., 1984; Kittelson et al., 1985; Obuchi et al., 1987; see also below) and, consistent with PAH formation, diesel combustion of fuels containing no PAH still results in PAH emissions (Barbella et al., 1989).

Kittelson et al. (1985) measured selected PAH and 1-nitropyrene concentrations in the cylinder and the exhaust manifold of an operating 5.7-liter V-8 diesel engine and observed that the PAH concentrations were higher in the cylinder than in the exhaust manifold, in contrast to higher 1-nitropyrene concentrations in the exhaust manifold. These data suggest (Kittelson et al., 1985) that the PAH are formed early in the combustion process and subsequently decay in the high temperature environment of the expansion stroke and the exhaust process. In contrast, 1-nitropyrene appears to be formed during the expansion and exhaust processes (Kittelson et al., 1985).

Laboratory investigations of the formation of PAH and soot in flames have shown that PAH are formed from non-PAH containing fuels, and that both PAH formation and destruction occur under combustion conditions (see, for example, Bockhorn et al. 1981; Ciajolo et al., 1982; Prado et al., 1985; Toqan et al., 1985; Frenklach et al., 1988; Frenklach, 1989). These

studies conclude that hydrogen atoms are involved in the formation/destruction reactions of PAH (Bockhorn et al., 1981; Frenklach et al., 1988; Frenklach, 1989) and suggest that the PAH are precursors to soot formation (Bockhorn et al., 1981; Ciajolo et al., 1982; Prado et al., 1985; Toqan et al., 1985; Frenklach et al., 1988; Kroto, 1988; Frenklach, 1989).

As is the case for combustion sources in general, a wide spectrum of PAH and alkyl-PAH are emitted from diesel exhaust (see above and Schuetzle et al., 1981; National Research Council, 1983; Jensen and Hites, 1983; Williams et al., 1986; Barbella et al., 1988, 1989). These PAH exhibit a wide range of volatilities (Table 2) and in the atmosphere are distributed between the gas and particle phases (Coutant et al., 1988; Arey et al., 1987, 1989a; Atkinson et al., 1988). Naphthalene and the other two ring PAH (including biphenyl) are present in ambient air in the gas phase, while the  $\geq 5$ -ring PAH such as benzo[a]pyrene are particle associated. As discussed by Bidleman and Foreman (1987), Bidleman (1988) and Pankow (1989), the vapor pressure parameter determining the phase distribution of semivolatile organic compounds is the subcooled liquid vapor pressure and not the solid-phase vapor pressure.

Based upon the ambient air sampling measurements of Coutant et al. (1988), those PAH with subcooled liquid vapor pressures of  $<10^{-6}$  Torr are essentially totally particle associated, and those with higher subcooled liquid vapor pressures are present, at least partially, in the gas phase. Thus, Bidleman (1988) concludes from ambient air monitoring data that the majority of fluoranthene and pyrene (subcooled liquid vapor pressures  $\sim 4 \times 10^{-5}$  Torr) are present in the gas phase, that benz[a]anthracene (subcooled liquid vapor pressure  $\sim 2 \times 10^{-6}$  Torr) is approximately equally distributed between gas and particle phase, and that benzo[a]pyrene (subcooled liquid vapor pressure  $\sim 9 \times 10^{-8}$  Torr) is essentially totally particle associated. This translates into PAH with  $\geq 5$  rings being particle associated, and PAH with  $\leq 4$  rings being, at least partially, in the gas phase in ambient air. A similar distribution between gas- and particle-phase will occur for the PAH-derivatives, with, for example, the  $\geq 4$ -ring nitro-PAH being particle-associated.

Table 2. Literature Room Temperature Vapor Pressures (Torr) for PAH

PAH	Vapor pressure at 298 K (Torr)	
	Solid <sup>a</sup>	Subcooled Liquid <sup>b</sup>
Naphthalene	$7.8 \times 10^{-2}$	$1.3 \times 10^{-1}$
Biphenyl	$9.9 \times 10^{-3}$ <sup>c</sup>	$2.1 \times 10^{-2}$
Acenaphthylene	$6.7 \times 10^{-3}$	
Acenaphthene	$2.2 \times 10^{-3}$	$7.6 \times 10^{-3}$
Fluorene	$6.0 \times 10^{-4}$	$3.1 \times 10^{-3}$
Phenanthrene	$1.2 \times 10^{-4}$	$5.3 \times 10^{-4}$
Anthracene	$6.0 \times 10^{-6}$	$5.0 \times 10^{-4}$
Fluoranthene	$9.2 \times 10^{-6}$	$5.2 \times 10^{-5}$
Pyrene	$4.5 \times 10^{-6}$	$3.3 \times 10^{-5}$
Benzo[a]fluorene		$1.2 \times 10^{-5}$
Benzo[b]fluorene		$1.1 \times 10^{-5}$
Benz[a]anthracene	$2.1 \times 10^{-7}$	$1.9 \times 10^{-6}$
Chrysene		$1.7 \times 10^{-6}$
Triphenylene		$1.7 \times 10^{-6}$
Benzo[b]fluoranthene		$1.6 \times 10^{-7}$
Benzo[k]fluoranthene		$1.6 \times 10^{-7}$
Benzo[e]pyrene		$1.0 \times 10^{-7}$
Benzo[a]pyrene		$9.0 \times 10^{-8}$
1-Methylnaphthalene	$6.7 \times 10^{-2}$	
2-Methylnaphthalene	$5.4 \times 10^{-2}$ <sup>d</sup>	

<sup>a</sup>From Sonnefeld et al. (1983) unless indicated.

<sup>b</sup>From Yamasaki et al. (1984).

<sup>c</sup>From Mackay et al. (1982).

<sup>d</sup>From Macknick and Prausnitz (1979).

### 3. Nitro-PAH

The presence of nitroarenes in diesel exhaust particulate extracts has received much attention because of the strong mutagenic activity of certain nitro-PAH, and numerous studies of the health effects of nitro-PAH have been conducted over the past ten years (see for example, Rosenkranz and Mermelstein, 1983; Tokiwa and Ohnishi, 1986; King, 1988; and references therein). More than fifty nitro-PAH have been tentatively identified in diesel exhaust, including mononitro-PAH, mononitro-alkyl-PAH, di- and trinitro-PAH and oxygenated nitro-PAH, such as nitrofluorenone (Schuetzle et al., 1982; Xu et al., 1982; Paputa-Peck et al., 1983; Robbat et al., 1986).

When isomer-specific analyses have been conducted, the major nitroarenes observed in diesel exhaust have been the isomers formed by electrophilic nitration of the parent PAH and in amounts consistent with the abundance and reactivity (Schuetzle, 1983; Nielsen, 1984) of the parent PAH toward electrophilic nitration. For example, 1-nitropyrene, 2-nitrofluorene, 9-nitroanthracene (Schuetzle and Perez, 1983; Paputa-Peck et al., 1983; Robbat et al., 1986), 6-nitrobenzo[a]pyrene (Gibson 1982; 1983; Schuetzle and Perez, 1983; Paputa-Peck et al., 1983) and 3-nitrofluoranthene (Robbat et al., 1986) have been observed in diesel exhaust extracts, all nitro-PAH isomers that are the major electrophilic nitration products of the parent PAH (Ruehle et al., 1985). Thus, although the PAH isomer-pair of fluoranthene and pyrene are present in diesel exhaust in very similar amounts (see Table 1), 1-nitropyrene is found in diesel exhaust particles in large excess over 3-nitrofluoranthene (Schuetzle et al., 1982; Paputa-Peck et al., 1983; Robbat et al., 1986).

Adding pyrene to hexadecane fuel in a single cylinder diesel engine increased nitropyrene emissions (Henderson et al., 1984). Henderson et al. (1984) suggested that since the nitropyrene/pyrene ratio decreased with increasing pyrene emissions, the active NO<sub>x</sub> species in the exhaust stream could be the limiting factor in nitro-PAH formation. Schuetzle and Perez (1983) estimated that an average of 12% of the 1-nitropyrene in diesel emissions collected on filters and using dilution tubes was formed on the filter during sampling. As will be discussed in more detail below, the nitro-PAH isomers observed in ambient particulate samples have very different isomer distributions than those found in diesel particulate extracts (Arey et al., 1987; 1989b; 1990a; Atkinson et al., 1988).



It is interesting to note that recent work (Ross et al., 1987) suggests that dinitropyrenes observed in diesel exhaust (Pederson and Siak, 1981; Nakagawa et al., 1983; Schuetzle, 1983; Schuetzle and Perez, 1983) are the result of direct nitration of pyrene rather than further nitration of 1-nitropyrene. However, Lee et al. (1989) suggest that dinitropyrene formation may be a collection artifact, being formed from 1-nitropyrene nitration by  $\text{NO}_2$ .

### C. Physical Loss Processes

The particles with which the PAH and PAH-derivatives are associated undergo wet and dry deposition. The average atmospheric lifetimes for particles due to dry deposition are a function of the particle diameter, as shown in Table 3 (Graedel and Weschler, 1981). As discussed above, the particles emitted from diesel-fueled vehicles have mass-median aerodynamic diameters of  $\sim 0.1\text{--}0.3\ \mu\text{m}$ , and hence the atmospheric lifetimes of these particles due to dry deposition are expected to be of the order of  $\sim 5\text{--}15$  days.

In the case of wet deposition, a washout ratio of particles  $W_p$ , defined as,

$$W_p = C_{\text{rain}}/C_{\text{air}}$$

relates the particle concentration in the rain ( $C_{\text{rain}}$ ) to that in the ambient air ( $C_{\text{air}}$ ). The deposition rate,  $F$ , is given by,

$$F = W_p J C_{\text{air}}$$

where  $J$  is the precipitation rate. For particles of  $0.1\text{--}10\ \mu\text{m}$  diameter and for particle-associated organics such as the polychlorinated biphenyls (PCBs) and the  $\text{C}_{20}\text{--}\text{C}_{23}$  n-alkanes the washout ratio  $W_p$  is  $10^{5\pm 1}$  (Eisenreich et al., 1981; Ligocki et al., 1985a). With a washout ratio of this magnitude, essentially complete scavenging of the particles and their associated organics occurs during a rain event (Leuenberger et al., 1985). Of course, rain-out is an episodic event which, in California, will lead to generally shorter lifetimes of the particle-associated PAH and PAH-derivatives in the winter months than during the summertime.

Table 3. Average Atmospheric Lifetimes for Particles Due to Dry Deposition (From Graedel and Weschler, 1981)

Diameter ( $\mu\text{m}$ )	Lifetime (days)
0.002	0.01
0.02	1
0.2	10
2	10
20	1
200	0.01

Thus, considering only its physical atmospheric removal processes, the lifetime of particle-associated PAH and PAH-derivatives due to particle dry deposition is expected to be around 5-15 days, with an additional efficient, but episodic, removal process involving the rain-out of particles.

#### D. Gas-Phase Loss Processes of PAH and PAH-Derivatives

As noted in the Introduction, the PAH and PAH-derivatives present, at least partially, in the gas phase can undergo wet and dry deposition, photolysis and reaction with OH and NO<sub>3</sub> radicals and O<sub>3</sub>. To date, there is no evidence for the occurrence of other gas-phase reactions (for example, with NO<sub>2</sub>, HNO<sub>3</sub> or SO<sub>2</sub>) for the PAH or nitroarenes (the only PAH-derivatives for which experimental data are available). In this Section, the available kinetic and product data concerning photolysis and reactions with OH and NO<sub>3</sub> radicals and O<sub>3</sub> are presented and discussed.

##### 1. Wet Deposition of Gas-Phase PAH and PAH-Derivatives

Experimental data concerning the washout ratios  $W_g$ , of gas-phase PAH, where

$$W_g = [\text{PAH}]_{\text{rain}}/[\text{PAH}]_{\text{gas}}$$

have been obtained and reported by Ligoeki et al. (1985b). The values obtained are given in Table 4. From concurrent measurements of PAH and

Table 4. Washout Ratios for Gas-Phase PAH (from Ligoeki et al., 1985b)

PAH	Molecular Weight	Washout ratio W (at 281 K)
Naphthalene	128	250 ± 73
1-Methylnaphthalene	142	330 ± 100
2-Methylnaphthalene	142	250 ± 78
Acenaphthylene	152	1600 ± 500
Acenaphthene	154	1000 ± 310
Fluorene	166	1500 ± 390
Phenanthrene	178	3400 ± 740
Anthracene	178	1900 ± 600
Methylphenanthrenes	192	2500 ± 800
Fluoranthene	202	6300 ± 2000
Pyrene	202	5900 ± 1800
Benz[a]anthracene	228	12000 ± 4900
Chrysene	228	18000 ± 6500
Benzo[e]pyrene	252	5800
Benzo[b+j+k]fluoranthene	252	7400 ± 1300
9-Fluorenone	180	11000 ± 2200
9,10-Anthracenedione	208	27000 ± 7000

PAH-derivative concentrations in air and in rainwater, Ligocki et al. (1985a) concluded that the PAH of molecular weight  $<252$  underwent wet deposition primarily from the gas phase, while the  $\geq 252$  molecular weight (MW) PAH underwent wet deposition primarily from the particle phase. This conclusion is in reasonable agreement with the review of Bidleman (1988), which concludes that wet deposition from the gas phase dominates for PAH of MW  $\leq 202$  and wet deposition from the particle phase dominates for PAH of MW  $\geq 228$ . These conclusions of Ligocki et al. (1985b) and Bidleman (1988) are consistent with the PAH of MW  $\leq 202$  being predominantly in the gas phase with relatively low washout ratios ( $10^2$ - $10^4$ ), while the PAH of MW  $\geq 252$  reside on or in particles which generally have higher washout ratios ( $10^{5\pm 1}$ ). In comparison with the washout ratios of the very efficiently removed particles, the washout ratios of  $10^2$ - $10^4$  for the gas-phase PAH suggest that wet deposition will not be a major tropospheric removal process for the PAH.

Dry deposition of gas-phase PAH is expected to be of minor importance (see, for example, Eisenreich et al., 1981).

## 2. Reaction with the OH Radical

The available rate constant data for the gas-phase reactions of the OH radical with PAH and PAH-derivatives have been evaluated and reviewed by Atkinson (1989). The rate constant data are given in Table 5. For most of the PAH and PAH-derivatives for which data are available, only single studies have been carried out, and no firm recommendations were made by Atkinson (1989) for these compounds. For biphenyl, naphthalene and phenanthrene, sufficient data are available that recommendations were made.

Biphenyl. The rate constants of Zetzsch (1982), Atkinson et al. (1984), Atkinson and Aschmann (1985) and Klöpffer et al. (1986), all obtained at room temperature, are in generally good agreement, although the rate constant determined by Zetzsch (1982) (which required knowledge of the vapor pressure of biphenyl) is lower by ~25% than the other values. Since no details are available concerning the study of Klöpffer et al. (1986), the rate constant from that study is not used in the evaluation. A unit-weighted average of the room temperature rate constants of Zetzsch (1982), Atkinson et al. (1984) and Atkinson and Aschmann (1985) leads to the recommendation of

Table 5. Rate Constants  $k$  and Temperature-Dependent Parameters for the Gas-Phase Reactions of the OH Radical with PAH and PAH-Derivatives at the High Pressure Limit (from Atkinson, 1989)

Aromatic	$10^{12} \times A$ ( $\text{cm}^3 \text{mole}^{-1} \text{s}^{-1}$ )	B (K)	$10^{12} \times k$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{s}^{-1}$ )	at T (K)	Technique	Reference	Tempera- ture range covered (K)
Biphenyl			$5.8 \pm 0.8$	296	FP-RF	Zetzsch (1982)	
			$7.61 \pm 0.67$	$294 \pm 1$	RR [relative to $k(\text{n-nonane}) =$ $1.01 \times 10^{-11} \text{s}^{-1}$ ]	Atkinson et al. (1984)	
			$8.32 \pm 0.75$	$295 \pm 1$	RR [relative to $k(\text{cyclohexane}) =$ $7.43 \times 10^{-12} \text{s}^{-1}$ ]	Atkinson and Aschmann (1985)	
1,4-Naphtho- quinone			8.0	300	RR [relative to $k(\text{ethene}) =$ $8.44 \times 10^{-12} \text{s}^{-1}$ ]	Klopffer et al. (1986)	
			$3.1 \pm 1.2$	$298 \pm 2$	RR [relative to $k(\text{cyclohexane}) =$ $7.49 \times 10^{-12} \text{s}^{-1}$ ]	Atkinson et al. (1989a)	
			13.0	300	RR [relative to $k(\text{ethene}) =$ $8.44 \times 10^{-12} \text{s}^{-1}$ ]	Klopffer et al. (1986)	
Fluorene					LP-RF	Lorenz and Zellner (1983, 1984)	300-873
Naphthalene			$18.6 \pm 1.0$	300			
			$14.6 \pm 5.0$	337			
			$11.0 \pm 4.4$	358			
			$10.1 \pm 4.0$	$378 \pm 2$			
	$2.3 \pm 1.5$	$-640 \pm 300$ (300-407 K)	$11.6 \pm 3.0$ $10.5 \pm 4.0$	404 407			

Table 5 (continued) - 2

Aromatic	$10^{12} \times A$ ( $\text{cm}^3 \text{mole}^{-1} \text{s}^{-1}$ )	B (K)	$10^{12} \times k$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{s}^{-1}$ )	at T (K)	Technique	Reference	Temperature range covered (K)
1-Methyl- naphthalene	50	2500 (636-873 K)	6.3 $\pm$ 2.0	452	RR [relative to k(n-nonane) = 1.01 $\times 10^{-11}$ ] <sup>a</sup>	Atkinson et al. (1984)	
			4.3 $\pm$ 1.5	476			
			1.3 $\pm$ 0.5	502			
			1.2 $\pm$ 0.4	525 $\pm$ 1			
			0.7 $\pm$ 0.2	528			
			0.6 $\pm$ 0.1	531			
			1.1 $\pm$ 0.1	636			
			1.1 $\pm$ 0.2	665			
			1.4 $\pm$ 0.2	727			
			3.0 $\pm$ 0.5	873			
			22.8 $\pm$ 1.6	294 $\pm$ 1			
			23.5 $\pm$ 0.6	298 $\pm$ 1		Biermann et al. (1985)	
			25.9 $\pm$ 2.4	295 $\pm$ 1		Atkinson and Aschmann (1986)	
			21.6	300	RR [relative to k(ethene) $\frac{1}{12}$ ] 8.44 $\times 10^{-12}$ ] <sup>a</sup>	Klopffer et al. (1986)	
			53.0 $\pm$ 4.8	298 $\pm$ 2	RR [relative to k(2-methyl-1,3-butadiene) = 1.01 $\times 10^{-10}$ ] <sup>a</sup>	Atkinson and Aschmann (1987)	

Table 5 (continued) - 3

Aromatic	$10^{12} \times A$ ( $\text{cm}^3 \text{ mole}^{-1} \text{ s}^{-1}$ )	B (K)	$10^{12} \times k$ ( $\text{cm}^3$ $\text{molecule}^{-1} \text{ s}^{-1}$ )	at T (K)	Technique	Reference	Tempera- ture range covered (K)
2-Methyl- naphthalene			$52.3 \pm 4.2$	$295 \pm 1$	RR (relative to $k(2\text{-methyl-1,3-butadiene})$ $= 1.02 \times 10^{-10} \text{ s}^{-1}$ )	Atkinson and Aschmann (1986)	
1-Nitro- naphthalene			$5.4 \pm 1.8$	$298 \pm 2$	RR (relative to $k(\text{cyclohexane}) =$ $7.49 \times 10^{-12} \text{ s}^{-1}$ )	Atkinson et al. (1989a)	
2-Nitro- naphthalene			$5.6 \pm 0.9$	$298 \pm 2$	RR (relative to $k(\text{cyclohexane}) =$ $7.49 \times 10^{-12} \text{ s}^{-1}$ )	Atkinson et al. (1989a)	
2,3-Dimethyl- naphthalene			$76.8 \pm 4.8$	$295 \pm 1$	RR (relative to $k(2\text{-methyl-1,3-butadiene})$ $= 1.02 \times 10^{-10} \text{ s}^{-1}$ )	Atkinson and Aschmann (1986)	
2-Methyl-1- nitronaphthalene			$< 8.3$	$298 \pm 2$	RR (relative to $k(\text{cyclohexane}) =$ $7.49 \times 10^{-12} \text{ s}^{-1}$ )	Arey et al. (1990b)	
1,4-Dichloro- naphthalene			5.8	300	RR (relative to $k(\text{toluene}) =$ $5.91 \times 10^{-12} \text{ s}^{-1}$ )	Klopffer et al. (1986)	
Acenaphthene			58.4	300	RR (relative to $k(\text{ethene}) =$ $8.44 \times 10^{-12} \text{ s}^{-1}$ )	Klopffer et al. (1986)	
			$103 \pm 13$	$296 \pm 2$	RR (relative to $k(2,3\text{-dimethyl-2-butene})$ $= 1.11 \times 10^{-10} \text{ s}^{-1}$ )	Atkinson and Aschmann (1988)	

Table 5 (continued) - k

Aromatic	$10^{12} \times A$ ( $\text{cm}^3 \text{mole}^{-1} \text{s}^{-1}$ )	B (K)	$10^{12} \times k$ ( $\text{cm}^3$ molecule $^{-1} \text{s}^{-1}$ )	at T (K)	Technique	Reference	Temperature range covered (K)
Acenaph- thylene			110 $\pm$ 11	296 $\pm$ 2	RR [relative to k(2,3-dimethyl-2-butene) $= 1.11 \times 10^{-10}$ ] <sup>a</sup>	Atkinson and Aschmann (1988)	
Phenan- threne				338	LP-RF	Lorenz and Zellner (1984)	338-748
			15.6 $\pm$ 2.0	355			
			16.1 $\pm$ 2.0	387			
			19.1 $\pm$ 2.5	399			
			12.0 $\pm$ 1.7	431			
			8.3 $\pm$ 0.8	492			
			4.0 $\pm$ 0.7	526			
			2.8 $\pm$ 0.2	597			
Anthracene			1.2 $\pm$ 0.4	648		Biermann et al. (1985)	298-319
			2.2 $\pm$ 0.5	748			
			34 $\pm$ 12	298 $\pm$ 1	RR [relative to k(propene), $1.3504/T$ ] <sup>a</sup>		
			28 $\pm$ 6	319 $\pm$ 1	$4.85 \times 10^{-12} e^{504/T}$ <sup>a</sup>		
			112 $\pm$ 9	325 $\pm$ 1	RR [relative to k(propene), $1.1$ ] <sup>a</sup>		

<sup>a</sup>From the recommendations of Atkinson (1989).



$$k(\text{biphenyl}) = 7.2 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } 298 \text{ K},$$

with an estimated overall uncertainty of  $\pm 30\%$ . This rate constant is that for OH radical addition to the aromatic rings. At temperatures  $\geq 400 \text{ K}$  the OH-biphenyl adduct will rapidly thermally decompose and only H-atom abstraction from the C-H bonds of the aromatic rings will be observed.

Naphthalene. The available rate constants obtained at, or close to, the high-pressure limit are given in Table 5. Lorenz and Zellner (1983) have shown that at  $378 \pm 2 \text{ K}$  the rate constant for this reaction is in the fall-off region between second- and third-order kinetics below  $\sim 50$  Torr total pressure of helium, but that no such fall-off behavior is observed at  $525 \pm 1 \text{ K}$ .

At temperatures  $\leq 410 \text{ K}$  the rate constants obtained by Lorenz and Zellner (1983, 1984), Atkinson et al. (1984), Biermann et al. (1985), Atkinson and Aschmann (1986) and Klöpffer et al. (1986) are in good agreement (the rate constant of Klöpffer et al. (1986) was not used in the evaluation because of a lack of details). A unit-weighted least-squares analysis of the data of Lorenz and Zellner (1983, 1984), Atkinson et al. (1984), Biermann et al. (1985) and Atkinson and Aschmann (1986) yields the recommended Arrhenius expression of

$$k(\text{naphthalene}, T \leq 410 \text{ K}) = (1.07^{+1.14}_{-0.55}) \times 10^{-12} e^{(895 \pm 239)/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

over the temperature range  $294\text{--}407 \text{ K}$ , where the indicated errors are two least-squares standard deviations, and

$$k(\text{naphthalene}) = 2.16 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

at  $298 \text{ K}$ , with an estimated overall uncertainty of  $\pm 30\%$  at  $298 \text{ K}$ .

Phenanthrene. The available rate constants of Lorenz and Zellner (1984) and Biermann et al. (1985) are given in Table 5. The rate constants obtained by Biermann et al. (1985) from a relative rate study are consistent with the higher temperature ( $T \geq 338 \text{ K}$ ) data of Lorenz and Zellner (1984). For the temperature range  $\leq 410 \text{ K}$  (the same as for

naphthalene) a unit-weighted least-squares analysis of the data of Lorenz and Zellner (1984) and Biermann et al. (1985) yields the recommendation of


$$k(\text{phenanthrene}; T \leq 410 \text{ K}) = (1.02^{+5.41}_{-0.86}) \times 10^{-12} e^{(1021 \pm 634)/T} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

over the temperature range 298-399 K, where the indicated errors are two least-squares standard deviations, and


$$k(\text{phenanthrene}) = 3.1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } 298 \text{ K},$$

with an estimated overall uncertainty at 298 K of  $\pm$  a factor of 2.

Reaction Mechanisms. As discussed by Atkinson (1986, 1989) the OH radical reactions with the PAH and PAH-derivatives proceed by two reaction pathways; namely OH radical addition to the aromatic ring to form an initially energy-rich hydroxycyclohexadienyl-type radical and OH radical interaction with the substituent groups, either through H atom abstraction from C-H or O-H bonds or OH radical addition to  $>\text{C}=\text{C}<$  bonds. For example, for acenaphthene



and for acenaphthylene



These OH radical reactions are essentially at the high pressure limits at room temperature and atmospheric pressure. For the alkyl-substituted PAH such as 1- and 2-methylnaphthalene, 2,3-dimethylnaphthalene and (probably) acenaphthene, the reaction pathway involving OH radical addition to the aromatic ring dominates under atmospheric conditions (Atkinson, 1989). For those PAH containing unsaturated cyclopenta-fused rings (acenaphthylene and acephenanthrylene), OH radical addition to the cyclopenta-fused ring  $>C=C<$  bond may be significant.

The products of these OH radical-initiated reactions are not well understood. The observed products of the OH radical-initiated reactions (in the presence of  $NO_x$ ) of naphthalene and biphenyl are hydroxy- and nitro-arenes (Atkinson et al., 1987). The yields of the hydroxyarenes are significantly higher than those of the nitroarenes, being 7% and 4% for 1- and 2-naphthol, respectively, in comparison to ~0.3% each for 1- and 2-nitronaphthalene. Similarly, the yield of 2-hydroxybiphenyl from biphenyl is 20% (with much lower amounts of 3- and 4-hydroxybiphenyl also being produced), while the single nitro-derivative observed is 3-nitrobiphenyl in ~5% yield.

The specific nitroarene isomers formed, and their product yields, from the gas-phase OH radical-initiated reactions of the 2- to 4-ring PAH 1- and 2-methylnaphthalene (Zielinska et al., 1989a), biphenyl (Atkinson et al., 1987; Arey et al., 1989b), acenaphthene (Arey et al., 1989b), acenaphthylene (Arey et al., 1989b), phenanthrene (Arey et al., 1989b), anthracene (Arey et al., 1989b), acephenanthrylene (Zielinska et al.,

1988), fluoranthene (Arey et al., 1986; Atkinson et al., 1990a) and pyrene (Arey et al., 1986; Atkinson et al., 1990a) are given in Table 6. Preliminary results from a recent study of the gas-phase OH radical-initiated reaction of fluorene shows that the major nitroarene product from this PAH is 3-nitrofluorene with a yield of ~1% (Arey, Helmig and Atkinson, unpublished results).

The available product data for the monocyclic aromatic hydrocarbons and biphenyl (Atkinson et al., 1987, 1989b, 1990b) indicate that the nitroarene product yields do not extrapolate to zero at low  $\text{NO}_2$  concentrations, and hence the nitroarene formation yields determined under laboratory conditions (see Table 6) are believed to be applicable to ambient atmospheric conditions. The nitroarene product formation yields are in all cases low, ranging from ~0.5-5%, and as noted above, the hydroxy-PAH yields for naphthalene and biphenyl are a factor of ~5-10 higher than their nitroarene yields. It is important to note that the majority of the OH radical-initiated reaction products of the PAH remain unidentified. While there are uncertainties about the reaction mechanisms, a recently postulated mechanism (Atkinson et al., 1989b) for the reaction of the OH radical in the presence of  $\text{NO}_x$  with fluoranthene which is consistent with our recent product data is shown in Figure 1.

Note that the nitroarenes formed from the OH radical-initiated reactions of the PAH (Table 6) are often isomers distinct from those observed in diesel exhaust particles. Thus as noted above, the most abundant nitro-isomers of pyrene, fluorene and fluoranthene observed in diesel exhaust are 1-nitropyrene, 2-nitrofluorene and 3-nitrofluoranthene, respectively, while the isomers formed from the gas-phase OH radical-initiated reactions of these PAH are 2-nitropyrene, 3-nitrofluorene and 2-nitrofluoranthene, respectively.

Table 6. Nitroarene Product Data for the Gas-Phase OH and NO<sub>3</sub> Radical-Initiated Reactions of PAH  
(Adapted from Arey et al., 1989b; Atkinson et al., 1990a)





PAH	Structure	Nitroarene Formed (Yield) in Reaction with	
		OH	NO <sub>3</sub>
Naphthalene		1-Nitronaphthalene (0.3%) 2-Nitronaphthalene (0.3%)	1-Nitronaphthalene (17%) 2-Nitronaphthalene (7%)
1-Methylnaphthalene <sup>a</sup>		1-Methyl-5-nitronaphthalene <sup>b</sup> 1-Methyl-4-nitronaphthalene <sup>b</sup> 1-Methyl-6-nitronaphthalene 1-Methyl-3-nitronaphthalene 1-Methyl-7-nitronaphthalene <sup>b</sup> 1-Methyl-2-nitronaphthalene <sup>b</sup> 1-Methyl-8-nitronaphthalene <sup>b</sup>	1-Methyl-3-nitronaphthalene 1-Methyl-5-nitronaphthalene 1-Methyl-4-nitronaphthalene 1-Methyl-8-nitronaphthalene 1-Methyl-6-nitronaphthalene 1-Methyl-7-nitronaphthalene 1-Methyl-2-nitronaphthalene <sup>b</sup> Total Yield (-30%)
2-Methylnaphthalene <sup>a</sup>		2-Methyl-5-nitronaphthalene 2-Methyl-6-nitronaphthalene 2-Methyl-7-nitronaphthalene 2-Methyl-4-nitronaphthalene 2-Methyl-8-nitronaphthalene 2-Methyl-3-nitronaphthalene <sup>b</sup> 2-Methyl-1-nitronaphthalene <sup>b</sup>	2-Methyl-4-nitronaphthalene 2-Methyl-1-nitronaphthalene 2-Methyl-5-nitronaphthalene 2-Methyl-8-nitronaphthalene 2-Methyl-3-nitronaphthalene 2-Methyl-7-nitronaphthalene 2-Methyl-6-nitronaphthalene Total Yield (-30%)
Acenaphthylene		4-Nitroacenaphthylene <sup>c</sup> (2%)	None observed <sup>d</sup>

Table 6 (continued) - 2





PAH	Structure	Nitroarene Formed (Yield) in Reaction with	
		OH	NO <sub>3</sub>
Acenaphthene <sup>a</sup>		5-Nitroacenaphthene 3-Nitroacenaphthene 4-Nitroacenaphthene Total Yield (-0.2%)	4-Nitroacenaphthene (40%) <sup>e</sup> 3-Nitroacenaphthene (-2%) <sup>e</sup> 5-Nitroacenaphthene (-2%) <sup>e</sup>
Biphenyl		3-Nitrobiphenyl (5%)	No reaction observed
Phenanthrene		Two isomers (not 9-nitrophenanthrene) Total Yield (<0.1%)	Four isomers (including 9-nitrophenanthrene) Total Yield (<1%)
Anthracene		1-Nitroanthracene <sup>f</sup> 2-Nitroanthracene Total Yield (-0.2%)	1-Nitroanthracene <sup>f</sup> 2-Nitroanthracene Total Yield (<2%)

Table 6 (continued) - 3




PAH	Structure	Nitroarene Formed (Yield) in Reaction with	
		OH	NO <sub>3</sub>
Fluoranthene		2-Nitrofluoranthene (3%) 7-Nitrofluoranthene (1%) 8-Nitrofluoranthene (0.3%)	2-Nitrofluoranthene (24%)
Pyrene		2-Nitropyrene (0.5%) 4-Nitropyrene (0.06%)	4-Nitropyrene (~0.06%)
Acephenanthrylene		Two isomers (not 4- or 5-nitro- acephenanthrylene) (~0.1%)	None observed

Table 6 (continued) - 4

- 
- <sup>a</sup>Nitro-isomers listed in approximate order of decreasing yield.
- <sup>b</sup>Formed, at least partially, as a sampling artifact by reaction of  $\text{NO}_2$  and/or  $\text{NO}_2 + \text{HNO}_3$ .
- <sup>c</sup>Presence of 1-nitroacenaphthylene attributed to artifact formation during sampling.
- <sup>d</sup> $\text{NO}_3$  Radical reaction occurs by  $\text{NO}_3$  radical addition to the cyclopenta-fused ring.
- <sup>e</sup>Not expected to be formed in ambient air because of concurrent H atom abstraction from the cyclopenta-fused ring.
- <sup>f</sup>Presence of 9-nitroanthracene attributed to artifact formation during sampling.







Figure 1. Reaction scheme for the OH radical-initiated reaction of fluoranthene in the presence of  $\text{NO}_x$ .

### 3. NO<sub>3</sub> Radical Reactions

Naphthalene and the alkyl-substituted naphthalenes are observed to react in N<sub>2</sub>O<sub>5</sub> - NO<sub>3</sub> - NO<sub>2</sub> - air mixtures, in which NO<sub>3</sub> radicals are generated by the thermal decomposition of N<sub>2</sub>O<sub>5</sub>



The disappearance rates of the naphthalenes relative to those of the alkenes such as propene and trans-2-butene in these reaction mixtures as a function of the NO<sub>2</sub> concentration indicate that the PAH loss processes are kinetically equivalent to reaction with N<sub>2</sub>O<sub>5</sub> (Pitts et al., 1985a; Atkinson et al., 1987; Atkinson and Aschmann, 1987, 1988; Atkinson et al., 1990c). As discussed by Atkinson and Aschmann (1988) and Atkinson et al. (1990c), this can be due to either an elementary reaction with N<sub>2</sub>O<sub>5</sub> or to a complex reaction sequence involving initial addition of NO<sub>3</sub> to the aromatic rings to form a nitratocyclohexadienyl-type radical which then either back decomposes to reactants or reacts exclusively with NO<sub>2</sub>




with the measured rate constant  $k_{\text{obs}}$  for reaction with the NO<sub>3</sub> radical being

$$k_{\text{obs}} = k_a k_c / k_b$$

A recent experimental study (Atkinson et al., 1990c) has shown that the reaction of naphthalene in N<sub>2</sub>O<sub>5</sub> - NO<sub>3</sub> - NO<sub>2</sub> - air mixtures occurs by the initial addition of the NO<sub>3</sub> radical, rather than by an elementary reaction with N<sub>2</sub>O<sub>5</sub>, and the other PAH are expected to react in an analogous manner.

For those PAH containing substituent groups, a parallel reaction pathway involving  $\text{NO}_3$  radical reaction with the substituent group(s) can also occur (Atkinson and Aschmann, 1988; Arey et al., 1989b), in addition to  $\text{NO}_3$  radical addition to the aromatic ring. For example, for acenaphthene



and for acenaphthylene  $\text{NO}_3$  radical addition to the cyclopenta-fused  $>\text{C}=\text{C}<$  bond is the dominant reaction pathway (Atkinson and Aschmann, 1988). Table 7 gives the available rate constant data for the  $\text{NO}_3$  radical reactions which proceed by H-atom abstraction from, or  $\text{NO}_3$  radical addition to, the substituent groups, and these reaction pathways are experimentally kinetically equivalent to an  $\text{NO}_3$  radical reaction.

Table 8 gives the rate coefficients  $k_{\text{obs}}$  (where  $k_{\text{obs}} = k_a k_c / k_b$ ) for the reactions of  $\text{NO}_3$  radicals with the aromatic ring(s), and these reactions are experimentally kinetically equivalent to reaction with  $\text{N}_2\text{O}_5$ . The rate coefficients for a reaction with  $\text{N}_2\text{O}_5$ ,  $k_{\text{N}_2\text{O}_5}$ , may be readily obtained from  $k_{\text{obs}}$  since  $k_{\text{N}_2\text{O}_5} = k_{\text{obs}} / K$ , where  $K$  is the equilibrium constant for the  $\text{NO}_3 + \text{NO}_2 \rightleftharpoons \text{N}_2\text{O}_5$  reactions and is given by (Atkinson, 1990a)  $K = 1.26 \times 10^{-27} e^{11275/T} \text{ cm}^3 \text{ molecule}^{-1}$  ( $3.41 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1}$  at 298 K).

Table 7. Rate Constants k for the Gas-Phase Reactions of the NO<sub>3</sub> Radical with PAH

Aromatic	k (cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> )	at T (K)	Technique	Reference
1,4-Naphthoquinone	$< 9.5 \times 10^{-16}$	$298 \pm 2$	RR [relative to k(propene) = $9.45 \times 10^{-15}$ ] <sup>b</sup>	Atkinson et al. (1989a)
Acenaphthene	$(4.59 \pm 2.45) \times 10^{-13}$	$296 \pm 2$	RR [relative to k( <u>trans</u> -2-butene) = $3.89 \times 10^{-13}$ ] <sup>b</sup>	Atkinson and Aschmann (1988)
Acenaphthylene	$(5.45 \pm 0.47) \times 10^{-12}$	$296 \pm 2$	RR [relative to k( <u>trans</u> -2-butene) = $3.89 \times 10^{-13}$ ] <sup>b</sup>	Atkinson and Aschmann (1988)
1-Nitronaphthalene	$\leq 7.2 \times 10^{-15}$	$298 \pm 2$	RR [relative to k(propene) = $9.45 \times 10^{-15}$ ] <sup>b</sup>	Atkinson et al. (1989a)
2-Nitronaphthalene	$\leq 7.3 \times 10^{-15}$	$298 \pm 2$	RR [relative to k(propene) = $9.45 \times 10^{-15}$ ] <sup>b</sup>	Atkinson et al. (1989a)
2-Methyl-1-nitro- naphthalene	$(1.13 \pm 0.57) \times 10^{-14}$	$298 \pm 2$	RR [relative to k(propene) = $9.45 \times 10^{-15}$ ] <sup>b</sup>	Arey et al. (1990b)

Table 8. Rate Constants  $k_{\text{obs}} = k_a k_c / k_b$  for the Gas-Phase Reactions of the  $\text{NO}_3$  Radical with PAH

Aromatic	$k_{\text{obs}} = k_a k_c / k_b$ ( $\text{cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
Biphenyl	$< 5 \times 10^{-30}$	$298 \pm 2$	RR [relative to $k_{\text{obs}}$ (naphthalene) = $3.6 \times 10^{-28}$ ]	Atkinson et al. (1987)
Naphthalene	$\sim (7-10) \times 10^{-28}$	$298 \pm 2$	S-IR [relative to $K(\text{NO}_3 + \text{NO}_2 \rightleftharpoons \text{N}_2\text{O}_5)$ = $3.41 \times 10^{-11}$ ]	Pitts et al. (1985a)
	$(4.77 \pm 0.69) \times 10^{-28}$	$298 \pm 2$	S-IR [relative to $K(\text{NO}_3 + \text{NO}_2 \rightleftharpoons \text{N}_2\text{O}_5)$ = $3.41 \times 10^{-11}$ ]	Atkinson et al. (1987)
	$3.31 \times 10^{-28}$	$296 \pm 2$	RR [relative to $k(\text{propene})$ = $9.45 \times 10^{-15}$ ]	Atkinson and Aschmann (1988)
	$(1.28 \pm 0.20) \times 10^{-27}$	$272.0 \pm 0.6$	RR [relative to $k(\text{thiophene})$ = $3.93 \times 10^{-14}$ ]	Atkinson et al. (1990a)
	$(1.10 \pm 0.27) \times 10^{-27}$	$275.2 \pm 0.6$		
	$(1.32 \pm 0.25) \times 10^{-27}$	$275.2 \pm 0.6$		
	$(6.96 \pm 1.73) \times 10^{-28}$	$281.9 \pm 0.6$		
	$(5.31 \pm 1.18) \times 10^{-28}$	$289.1 \pm 0.6$		

Table 8 (continued) - 2

Aromatic	$k_{\text{obs}} = k_a k_c / k_b$ ( $\text{cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
	$(3.64 \pm 1.11) \times 10^{-28}$	$296.3 \pm 0.6$		
	$(3.66 \pm 1.26) \times 10^{-28}$	$296.9 \pm 0.6$		
Naphthalene-dg	$(4.4 \pm 0.4) \times 10^{-28}$	$298 \pm 2$	RR [relative to $k_{\text{obs}}$ (naphthalene) = $3.6 \times 10^{-28}$ ]	Atkinson et al. (1990a)
1-Methyl-naphthalene	$(8.4 \pm 1.0) \times 10^{-28}$	$298 \pm 2$	RR [relative to $k_{\text{obs}}$ (naphthalene) = $3.6 \times 10^{-28}$ ] <sup>a</sup>	Atkinson and Aschmann (1987)
	$7.0 \times 10^{-28}$	$296 \pm 2$	RR [relative to $k(\text{trans-2-butene})$ = $3.89 \times 10^{-13}$ ]	Atkinson and Aschmann (1988)
2-Methyl-naphthalene	$(1.07 \pm 0.16) \times 10^{-27}$	$298 \pm 2$	RR [relative to $k_{\text{obs}}$ (naphthalene) = $3.6 \times 10^{-28}$ ]	Atkinson and Aschmann (1987)
	$1.09 \times 10^{-27}$	$296 \pm 2$	RR [relative to $k(\text{propene})$ = $9.45 \times 10^{-15}$ ]	Atkinson and Aschmann (1988)

Table 8 (continued) - 3

Aromatic	$k_{\text{obs}} = k_a k_c / k_b$ ( $\text{cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
2,3-Dimethyl-naphthalene	$(1.48 \pm 0.44) \times 10^{-27}$	$298 \pm 2$	RR [relative to $k_{\text{obs}}$ (naphthalene) = $3.6 \times 10^{-28}$ ]	Atkinson and Aschmann (1987)
	$1.61 \times 10^{-27}$	$296 \pm 2$	RR [relative to $k$ (propene) = $9.45 \times 10^{-15}$ ]	Atkinson and Aschmann (1988)
Acenaphthene	$(1.65 \pm 0.38) \times 10^{-27}$	$296 \pm 2$	RR [relative to $k$ (trans-2-butene) = $3.89 \times 10^{-13}$ ]	Atkinson and Aschmann (1988)
1-Nitro-naphthalene	$(3.02 \pm 1.42) \times 10^{-29}$	$298 \pm 2$	RR [relative to $k$ (propene) = $9.45 \times 10^{-15}$ ]	Atkinson et al. (1989a)
2-Nitro-naphthalene	$(2.74 \pm 1.14) \times 10^{-29}$	$298 \pm 2$	RR [relative to $k$ (propene) = $9.45 \times 10^{-15}$ ]	Atkinson et al. (1989a)
1,4-Naphthoquinone	$< 8.5 \times 10^{-30}$	$298 \pm 2$	RR [relative to $k$ (propene) = $9.45 \times 10^{-15}$ ]	Atkinson et al. (1989a)

Table 8 (continued) - 4

Aromatic	$k_{\text{obs}} = k_a k_c / k_b$ ( $\text{cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$ )	at T (K)	Technique	Reference
2-Methyl-1-nitro naphthalene	$(2.74 \pm 1.80) \times 10^{-29}$	$298 \pm 2$	RR [relative to $k(\text{propene})$ $= 9.45 \times 10^{-15}$ ]	Arey et al. (1990b)
Fluoranthene	$\sim 4.7 \times 10^{-28}$	$298 \pm 2$	RR [relative to $k_{\text{obs}}(\text{naphthalene})$ $= 3.6 \times 10^{-28}$ ]	Atkinson et al. (1990a)
Pyrene	$\sim 1.4 \times 10^{-27}$	$298 \pm 2$	RR [relative to $k_{\text{obs}}(\text{naphthalene})$ $= 3.6 \times 10^{-28}$ ]	Atkinson et al. (1990a)



The reactions which involve the initial addition of the  $\text{NO}_3$  radical to the aromatic ring (termed hereafter an  $\text{N}_2\text{O}_5$  reaction for simplicity) lead to the formation of nitroarenes (Pitts et al., 1985a; Sweetman et al., 1986; Atkinson et al., 1987, 1990a; Zielinska et al., 1989a; Arey et al., 1989b), and these nitroarene yield data are given in Table 6. The reaction routes involving  $\text{NO}_3$  radical interaction with the substituent group(s) do not lead to the formation of nitroarenes, as expected from the likely subsequent chemistry (Atkinson, 1990b). The other products of these gas-phase " $\text{N}_2\text{O}_5$ " reactions of the PAH are presently not known with any certainty, although they may include hydroxynitro-PAH.

#### 4. $\text{O}_3$ Reactions

The available rate constant data for reaction of PAH with  $\text{O}_3$  are given in Table 9. Only for acenaphthylene has gas-phase reaction been observed (Atkinson and Aschmann, 1988), and reaction is also expected to occur for acephenanthrylene (Zielinska et al., 1988). Clearly, these PAH react with  $\text{O}_3$  by addition of  $\text{O}_3$  at the cyclopenta-fused ring  $>\text{C}=\text{C}<$  bond (Atkinson and Aschmann, 1988).

#### 5. Photolysis

No evidence has been observed for the gas-phase photolysis of the 2-4 ring PAH (Atkinson et al., 1984; Biermann et al., 1985; Atkinson and Aschmann, 1986, 1988). However, photolysis of 1- and 2-nitronaphthalene and 2-methyl-1-nitronaphthalene has been observed under ambient outdoor sunlight conditions (Atkinson et al., 1989a; Arey et al., 1990b). Photolysis of 1-nitronaphthalene, 2-methyl-1-nitronaphthalene and the 2-, 7- and 8-nitrofluoranthenes (the nitrofluoranthenes being totally particle-associated under atmospheric conditions) have also been observed in an indoor chamber with blacklamp irradiation (Atkinson et al., 1989a, 1990a; Arey et al., 1990b), with the photolysis rates of the 1-nitronaphthalene and 2-methyl-1-nitronaphthalene in the indoor chamber being approximately an order of magnitude higher than under ambient conditions (Atkinson et al., 1989a; Arey et al., 1990b). The photolysis rates are given in Table 10.

#### 6. Atmospheric Lifetimes of Gas-Phase PAH and PAH-Derivatives

The photolysis and reaction rate data given above can be combined with the ambient radiation flux and the ambient concentrations of OH and  $\text{NO}_3$  radicals,  $\text{NO}_2$  and  $\text{O}_3$  to allow the estimation of the lifetimes of the

Table 9. Rate Constants for the Gas-Phase Reactions of PAH and PAH-Derivatives with O<sub>3</sub>

Aromatic	$k$ (cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> )	at T (K)	Reference
Naphthalene	$< 2 \times 10^{-19}$	$294 \pm 1$	Atkinson et al. (1984)
1-Methylnaphthalene	$< 3 \times 10^{-19}$	$295 \pm 1$	Atkinson and Aschmann (1986)
2-Methylnaphthalene	$< 1.3 \times 10^{-19}$	$298 \pm 2$	Atkinson and Aschmann (1987)
2,3-Dimethylnaphthalene	$< 4 \times 10^{-19}$	$295 \pm 1$	Atkinson and Aschmann (1986)
Biphenyl	$< 4 \times 10^{-19}$	$295 \pm 1$	Atkinson and Aschmann (1986)
Acenaphthene	$< 2 \times 10^{-19}$	$294 \pm 1$	Atkinson et al. (1984)
Acenaphthylene	$< 5 \times 10^{-19}$	$296 \pm 2$	Atkinson and Aschmann (1988)
1-Nitronaphthalene	$5.5 \times 10^{-16}$	$296 \pm 2$	Atkinson and Aschmann (1988)
2-Nitronaphthalene	$< 6 \times 10^{-19}$	$298 \pm 2$	Atkinson et al. (1989a)
2-Methyl-1-nitronaphthalene	$< 6 \times 10^{-19}$	$298 \pm 2$	Atkinson et al. (1989a)
1,4-Naphthoquinone	$< 3 \times 10^{-19}$	$298 \pm 2$	Arey et al. (1990b)
	$< 2 \times 10^{-19}$	$298 \pm 2$	Atkinson et al. (1989a)

Table 10. Measured Photolysis Rates,  $k_{\text{phot}}$ , of PAH and PAH-Derivatives in a 6400 Liter Indoor Teflon Chamber with Blacklight Irradiation and Outdoors in a 1000 Liter Outdoor Teflon Chamber with Sunlight Irradiation<sup>a</sup>

Aromatic	$10^4 \times k_{\text{phot}} \text{ (s}^{-1}\text{)}$	
	Indoors	Outdoors
1-Nitronaphthalene	$15.9 \pm 1.1$	$1.37 \pm 0.10$
2-Nitronaphthalene	$\leq 2$	$1.06 \pm 0.08$
1,4-Naphthoquinone	$1.1 \pm 0.8$	$\sim 0.9$
2-Methyl-1-nitronaphthalene	$37 \pm 7$	$1.1 \pm 0.3$
3-Nitrobiphenyl	$< 0.17$	
NO <sub>2</sub>	78	43

<sup>a</sup>From Atkinson et al. (1989a) and Arey et al. (1990b).

PAH and PAH-derivatives with respect to each of these tropospheric loss processes. These lifetime data are given in Table 11. For the PAH not containing cyclopenta-fused rings, the dominant tropospheric loss process is by reaction with the OH radical, with calculated lifetimes of 1 day or less (note that OH radical reaction only occurs during daylight hours). The PAH containing cyclopenta-fused rings such as acenaphthene, acenaphthylene and acephenanthrylene are expected to also react with NO<sub>3</sub> radicals at a significant rate. (Note that NO<sub>3</sub> radical addition to the fused rings of the PAH is not significant as a tropospheric loss process for the PAH.) PAH having unsaturated cyclopenta-fused rings, such as acenaphthylene, acephenanthrylene and cyclopenta[cd]pyrene (MW 226), are expected to react with O<sub>3</sub> at a significant rate.

In contrast to O<sub>3</sub> and the OH radical, which are ubiquitous at reasonably consistent (on a day-to-day level) ambient concentrations (see, for example, Logan, 1985; Prinn et al., 1987; Arey et al., 1989a), the ambient concentrations of the NO<sub>3</sub> radical in the lower troposphere over continental areas exhibit large variations, with the mixing ratios ranging

Table 11. Calculated Atmospheric Lifetimes of PAH and PAH-Derivatives Due to Photolysis and Gas-Phase Reaction with OH and NO<sub>3</sub> Radicals and O<sub>3</sub>

PAH	Lifetime due to reaction with			
	OH <sup>a</sup>	NO <sub>3</sub> <sup>b</sup>	O <sub>3</sub> <sup>c</sup>	Photolysis <sup>d</sup>
Naphthalene	8.6 hr	100 days	>80 days	
1-Methylnaphthalene	3.5 hr	50 days	>125 days	
2-Methylnaphthalene	3.6 hr	40 days	>40 days	
2,3-Dimethylnaphthalene	2.4 hr	25 days	>40 days	
Biphenyl	2.1 days	>20 yr	>80 days	
Acenaphthene	1.8 hr	2.5 hr	>30 days	
Acenaphthylene	1.7 hr	13 min	~43 min	
Phenanthrene	6.0 hr			
Anthracene	1.4 hr			
Fluoranthene	~3.7 hr <sup>e</sup>	~85 days		
Pyrene	~3.7 hr <sup>e</sup>	~30 days		
1-Nitronaphthalene	2.9 days	3.6 yr	>28 days	1.7 hr
2-Nitronaphthalene	2.8 days	4.0 yr	>28 days	2.2 hr
1,4-Naphthoquinone	5.0 days	>100 days	>80 days	~2.6 hr
2-Methyl-1-nitro-naphthalene	>1.8 days	4.0 yr	>55 days	2.1 hr

<sup>a</sup>For a 12-hr daytime average OH radical concentration of  $1.5 \times 10^6$  molecule cm<sup>-3</sup> (Prinn et al., 1987).

<sup>b</sup>For a 12-hr average nighttime NO<sub>3</sub> radical concentration of  $2.4 \times 10^8$  molecule cm<sup>-3</sup> and an NO<sub>2</sub> concentration of  $2.4 \times 10^{12}$  molecule cm<sup>-3</sup> (Atkinson et al., 1986).

<sup>c</sup>For a 24-hr average O<sub>3</sub> concentration of  $7 \times 10^{11}$  molecule cm<sup>-3</sup> (Logan, 1985).

<sup>d</sup>For an average 12-hr daytime NO<sub>2</sub> photolysis rate of  $J_{\text{NO}_2} = 5.2 \times 10^{-3} \text{ s}^{-1}$ .

<sup>e</sup>Using estimated OH radical reaction rate constant of  $5 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  based on rate constant correlation with ionization potential (Biermann et al., 1985; Arey et al., 1990b; Atkinson et al., 1990a).

from <2 parts-per-trillion (ppt) to 430 ppt (Atkinson et al., 1986), and the ambient tropospheric concentration of the  $\text{NO}_3$  radical at any given time (during nighttime) and place must be viewed as being uncertain by at least a factor of 10. Hence to a good approximation the dominant tropospheric removal process for the PAH is by daytime reaction with the OH radical, leading to lifetimes of ~8 hours or less.

As seen from the rate constant data given in Table 5 and from the calculated lifetimes in Table 11, the presence of the nitro- substituent group in the nitroarenes leads to a marked decrease in their reactivity towards the OH radical. To date kinetic and product studies have only been carried out for three gas-phase fused-ring nitroarenes (Atkinson et al., 1989a; Arey et al., 1990b), and photolysis will be the dominant tropospheric removal process for these compounds, with calculated lifetimes of ~2 hrs.

#### 7. Evidence for Atmospheric Transformations of PAH

The recent ambient air measurement study of Arey et al. (1989a) provided clear evidence for the reactions of the volatile PAH with the OH radical, with the nighttime/daytime concentration ratios exhibiting a clear linear correlation with the OH radical reaction rate constant (Figure 2). From an estimate of the nighttime dilution rate, provided by the daytime/nighttime ratio of 3-nitrobiphenyl [a nitroarene believed to be formed only in the atmosphere from the daytime reaction of biphenyl with the OH radical in the presence of  $\text{NO}_x$  (Atkinson et al., 1987)], an average 12-hr daytime OH radical concentration of  $2.2 \times 10^6$  molecule  $\text{cm}^{-3}$  (during August) was derived, uncertain to at least a factor of 2. This estimated OH radical concentration in an urban area is essentially identical to the annually averaged global tropospheric 12-hr daytime OH radical concentration of  $1.5 \times 10^6$  molecule  $\text{cm}^{-3}$  (Prinn et al., 1987) and provides very strong evidence that the gas-phase PAH do react in the troposphere. Furthermore, as discussed in detail in the literature (Nielsen et al., 1984; Pitts et al., 1985b; Nielsen and Ramdahl, 1986; Sweetman et al., 1986; Arey et al., 1986, 1987, 1989a,b, 1990a; Ramdahl et al., 1986; Zielinska et al., 1988, 1989a,b; Atkinson et al., 1988), many of the nitroarenes observed in ambient air are only formed in the atmosphere through the gas-phase reactions of the 2-4 ring PAH. For example, nitroarene isomers observed both in OH radical-initiated reactions of the




Figure 2. Plot of the average nighttime/average daytime concentration ratio for the volatile PAH measured in Glendora, CA against their OH radical reaction rate constants (Adapted from Arey et al. 1989a).

parent PAH and in ambient air samples, but not in diesel exhaust samples include: 2-nitrofluoranthene, 2-nitropyrene, 3-nitrobiphenyl and 2-methyl-7-nitronaphthalene (Arey et al., 1989b; Zielinska et al., 1989a).

The importance of atmospheric formation of nitroarenes is clear from a comparison of the calculated and observed 3-nitrobiphenyl, 1- + 2-nitronaphthalene, 2-nitrofluoranthene and 2-nitropyrene concentrations at two sites in southern California (Arey et al., 1990b). The predicted concentrations of these nitroarenes were calculated from the reaction rate constants (Table 5) and nitroarene product formation yields (Table 6) of the OH radical-initiated reactions of biphenyl, naphthalene, fluoranthene and pyrene, respectively, using estimated OH radical concentrations and the measured ambient parent PAH concentrations, and incorporating the photolysis loss of the nitronaphthalenes (Table 10). There was excellent agreement between the calculated nitroarene concentrations and the measured ambient concentrations of the nitroarenes at both sites (Table 12). Only the nitronaphthalenes are expected to be present in direct emissions such as diesel exhaust, and the slightly higher predicted than observed concentrations for the nitronaphthalenes suggests that atmospheric formation of these species dominates over their direct emission, at least at these two sites and at the time of the measurements.

Table 12. Ambient Nitroarene Concentrations at Torrance, CA in February 1986 and Glendora, CA, in August 1989 (Arey et al., 1990b)

	<u>12-Hr Average Daytime Concentration (ng m<sup>-3</sup>)</u>			
	TORRANCE		GLENDORA	
	Measured	Calculated	Measured	Calculated
1- + 2-Nitro-naphthalene	5.9	15	4.7	8.0
3-Nitrobiphenyl	6.0	2.4	1.0	1.2
2-Nitrofluoranthene	0.3	1.3	0.27	0.41
2-Nitropyrene	0.04	0.15	0.012	0.028

#### E. Adsorbed-Phase Loss Processes of PAH and PAH-Derivatives

In addition to the physical removal of the particles (Section C) with which the  $\geq$ MW 252 PAH and some fraction of the four-ring PAH are associated, the particle-associated PAH can also be transformed by a number of chemical processes in the adsorbed phase. These include photolysis and reaction with  $O_3$ ,  $SO_2$ ,  $NO_2$  and/or  $HNO_3$  and  $N_2O_5$ , and a large number of experimental investigations of these potential reaction pathways have been carried out (see, for example, Jager and Rakovic, 1974; Lane and Katz, 1977; Pitts et al., 1978, 1980, 1985c, 1986; Katz et al., 1979; Hughes et al., 1980; Jäger and Hanus, 1980; Korfmacher et al., 1980a,b, 1981; Peters and Seifert, 1980; Blau and Güsten, 1981; Butler and Crossley, 1981; Tokiwa et al., 1981; Daisey et al., 1982; Brorström et al., 1983; Grosjean et al., 1983; Nielsen et al., 1983; Ramdahl et al., 1984; Valerio et al., 1984, 1987; Van Vaeck and Van Cauwenberghe, 1984; Wu et al., 1984; Behymer and Hites, 1985, 1988; Benson et al., 1985; Brorström-Lunden and Lindskog, 1985; Kamens et al., 1985a,b, 1986, 1988, 1989; Lindskog et al., 1985; Wu and Niki, 1985; Yokley et al., 1985, 1986; Güsten, 1986; Ménard et al., 1986; Schuetzle and Frazier, 1986; Takeda and Teranishi, 1986; Cope and Kalkwarf, 1987; Holloway et al., 1987; Arey et al., 1988; Coutant et al., 1988; Nielsen, 1988; Greenberg, 1989). Interpretation of the experimental data obtained from the laboratory studies is rendered difficult due to the diverse substrates used, and in many cases conflicting data have been obtained. As discussed below, some differences may be attributed to the availability of the PAH for reaction, i.e., whether present as an adsorbed surface layer or included in the particles, or whether the particles were "freshly" emitted or were "aged" particles.

Benzo[a]pyrene (BaP) is a carcinogenic (IRAC, 1983) PAH often measured as a surrogate for all toxic PAH in ambient air particles. As noted in Section C, BaP (MW 252) will be particle-associated in ambient air and gas-phase reactions of BaP will be unimportant. Since the adsorbed-phase reactions of BaP have been studied more than those for any other single PAH, and in addition, BaP is considered to be among the more reactive PAH (Nielsen, 1984; Greenberg and Darack, 1987), it is appropriate to discuss the adsorbed-phase reactions of BaP in some detail. As can be seen from Table 13, which gives Nielsen's reactivity classifications toward electrophilic reactions, BaP is in Class 2 of a five class



Table 13. Reactivity Scale of PAH (Including Benzene and Biphenyl) in Electrophilic Reactions (from Nielsen, 1984)

---

- Class 1: benzo[a]tetracene, dibenzo[a,h]pyrene, pentacene, tetracene
- Class 2: anthanthrene, anthracene, benzo[a]pyrene, cyclopenta[c,d]pyrene, dibenzo[a,l]pyrene, dibenzo[a,i]pyrene, dibenzo[a,c]tetracene, perylene
- Class 3: benz[a]anthracene, benzo[g]chrysene, benzo[ghi]perylene, dibenzo[a,e]pyrene, picene, pyrene
- Class 4: benzo[c]chrysene, benzo[c]phenanthrene, benzo[e]pyrene, chrysene, coronene, dibenzanthracenes, dibenzo[e,l]pyrene
- Class 5: acenaphthylene, benzofluoranthenes, fluoranthene, indeno[1,2,3-cd]fluoranthene, indeno[1,2,3-cd]pyrene, naphthalene, phenanthrene, triphenylene
- Class 6: benzene, biphenyl
- 

(Class 1 being the most reactive) scale. The PAH in Class 1 have not generally been observed in significant concentrations in diesel exhaust.

1. Reaction with O<sub>3</sub>

Benzo[a]pyrene. The results obtained concerning the stability of BaP adsorbed onto surfaces in the presence of O<sub>3</sub> are disparate. Thus, Katz and coworkers (Katz et al., 1979) and Peters and Seifert (1980) observed that BaP reacted in the presence of ambient levels of O<sub>3</sub>, and Pitts et al. (1980) identified the BaP-4,5-epoxide as a product of the reaction of O<sub>3</sub> with BaP deposited onto glass fiber filters. These studies indicated a BaP lifetime with respect to reaction with O<sub>3</sub> (at ambient concentrations, i.e., levels reached in polluted atmospheres) of the order of an hour. In contrast, Grosjean et al. (1983) observed no loss of BaP,

either deposited onto blank filters or on filters coated with ambient particles, fly ash or diesel particles, when exposed to 100 ppb of ozone for three hours. Brorström et al. (1983) observed no definitive evidence for losses of BaP (or other PAH) under high-volume sampling conditions with the addition of 200 ppb of  $O_3$ . However, it should be noted that in this latter study, no BaP was added to the ambient particles.

In contrast, the relatively recent studies of Van Vaeck and Van Cauwenberghe (1984), Lindskog et al. (1985) and Pitts et al. (1986) all obtained data showing that BaP exhibited significant losses (typically 50%) when exposed to 50-1500 ppb of  $O_3$  for time scales of 0.5-6 hr. Thus, Van Vaeck and Van Cauwenberghe (1984) observed that the percentage conversion of BaP in diesel exhaust particles, when exposed to 1.5 ppm of  $O_3$  under high-volume sampling conditions, increased slightly with the exposure time, from 62% at 0.5 hr to 87% at 4 hr. Lindskog et al. (1985) observed conversions of BaP, on soot generated in a smoke generator, at 6 hr exposure times [45% relative humidity (RH)] increasing from 3% at 0.1 ppm  $O_3$  to 80% at 1.0 ppm  $O_3$ . Pitts et al. (1986) observed an approximately 50% conversion of BaP, independent of the  $O_3$  concentration (0.05-0.29 ppm  $O_3$  at 1% RH) and of the humidity (1-50% RH at 0.20 ppm  $O_3$ ), for a 3 hr exposure time. In the study of Pitts et al. (1986), BaP was either adsorbed on filters or present in particles collected close to a freeway, and the exposures were both passive (in a chamber) and in a flow system. Another recent study found degradation of BaP on ambient particles enriched with BaP and exposed to 0.18 ppm  $O_3$ , but no degradation on ambient particles sampled with an  $O_3$  enriched (0.18 ppm) vapor stream (Coutant et al., 1988).

These data are generally consistent with the fact that BaP reacts with  $O_3$ , providing that the BaP is available for reaction. This is suggested by the often initially rapid BaP degradation followed by significantly slower BaP losses (Pitts et al., 1986; Coutant et al., 1988). Those experiments in which BaP was coated or deposited onto surfaces should then exhibit reactive losses of BaP, as generally observed (Katz et al., 1979; Peters and Seifert, 1980; Pitts et al., 1980, 1986). Fresh diesel particles (Van Vaeck and Van Cauwenberghe, 1984) and ambient particles collected near a freeway (Pitts et al., 1986) also showed BaP losses upon exposure to  $O_3$ . In contrast, BaP included in particles, as

may be expected to be the case for aged particulate matter or particles to which condensation of secondary aerosol has occurred (expected to be the case in urban atmospheres), will not be available for reaction with gaseous ozone and hence under these conditions BaP will be observed to be nonreactive (Brorström et al., 1983; Coutant et al., 1988).

Other PAH. The general conclusions drawn above concerning the reactions of BaP with ozone are also expected to apply for the other PAH, and this is supported by the available experimental data (see, for example, Brorström et al., 1983; Grosjean et al., 1983; Van Vaeck and Van Cauwenberghe, 1984; Lindskog et al., 1985; Pitts et al., 1986; Coutant et al., 1988; Greenberg, 1989). Thus, PAH included in particles will be essentially non-reactive, while those present on the particle surfaces will react with  $O_3$ , consistent with their reactivities toward  $O_3$ . The PAH which may be expected to be more reactive than BaP toward  $O_3$  include the linear PAH (not generally very abundant in diesel exhaust) and the group of PAH containing an unsaturated cyclopenta-fused ring [i.e., with an extracyclic olefinic linkage] (Greenberg, 1989). Apart from the PAH with unsaturated cyclopenta-fused rings [as noted above, acenaphthylene, a gas-phase PAH with an unsaturated cyclopenta-fused ring, reacts in the gas-phase with  $O_3$  (Atkinson and Aschmann, 1988)], the reactivities of the PAH toward electrophilic reaction with  $O_3$  are expected to follow Nielsen's reactivity scale (Table 13).

## 2. Reaction with Oxides of Nitrogen

Benzo[a]pyrene. A number of studies have been carried out to investigate the reaction of BaP when BaP-associated particles are exposed to oxides and oxyacids of nitrogen (Pitts et al., 1978, 1985; Jäger and Hanus, 1980; Butler and Crossley, 1981; Grosjean et al., 1983; Brorström et al., 1983; Ramdahl et al., 1984; Yokley et al., 1985; Lindskog et al., 1985; Brorström-Lunden and Lindskog, 1985; Kamens et al., 1986; Arey et al., 1988). Although all of these studies observed losses of BaP and, generally, formation of nitro-BaP upon exposure of BaP-containing particles to  $NO_2$  and/or  $HNO_3$ , the data reported are conflicting as to whether  $NO_2$  or  $HNO_3$  is the reactive species. In recent studies in which  $HNO_3$  was removed from the  $NO_2$  exposure gas, no reaction of adsorbed BaP with  $NO_2$  was observed (Grosjean et al., 1983; Yokley et al., 1985). In contrast, Lindskog et al. (1985) did not detect any reaction of BaP on

soot with HONO or HNO<sub>3</sub>, but did observe reaction with NO<sub>2</sub> and NO<sub>2</sub> + HNO<sub>3</sub> mixtures. In this latter study, the buffering effect of the soot may have played a role, and oxy- rather than nitro-PAH derivatives were generally observed as products.

Other PAH. Recent studies of pyrene in methylene chloride solution did not result in nitration when HNO<sub>3</sub> was present, but resulted in the formation of dinitropyrenes when N<sub>2</sub>O<sub>4</sub> was present (Ross et al., 1987). These recent results of Ross and coworkers (1987) suggest that the role of HNO<sub>3</sub> in the nitration of PAH adsorbed on diesel soot may be to react with the soot, leading to deactivation of the soot surface and thus allowing the limited amount of NO<sub>2</sub>/N<sub>2</sub>O<sub>4</sub> to nitrate the PAH. Whether under ambient conditions HNO<sub>3</sub> alone can react to nitrate PAH adsorbed on diesel soot remains in question. These PAH reactions are substrate dependent (Jäger and Hanus, 1980; Ramdahl et al., 1984; Yokley et al., 1985) and, as noted above for the O<sub>3</sub> reactions, presumably require that the PAH be accessible for reaction.

A number of studies have been carried out in which several PAH, in addition to BaP, were examined (Butler and Crossley, 1981; Tokiwa et al., 1981; Brorström et al., 1983; Ramdahl et al., 1984; Lindskog et al., 1985; Pitts et al., 1985c). In these studies in which several PAH were examined, the nitro-PAH formed were the isomers expected from electrophilic nitration and the amounts of reaction followed the reactivity ranking of the PAH toward electrophilic nitration (Nielsen, 1984; Table 13). An apparent exception is a tentative identification of 2-nitropyrene, in addition to 1-nitropyrene, from exposure of pyrene adsorbed on silica gel to NO<sub>2</sub> in N<sub>2</sub> in the presence of light (Wu and Niki, 1985).

In ambient air, NO<sub>2</sub> and HNO<sub>3</sub> will both be present. Recently, Arey et al. (1988) studied the disappearance of perdeuterated fluoranthene, pyrene, BaP, and perylene adsorbed onto ambient particles, and the formation of the perdeuterated nitro-PAH under high-volume sampling conditions during a wintertime high-NO<sub>x</sub> (maximum NO<sub>x</sub> concentrations approximately 1 ppm) pollution episode in the Los Angeles air basin. BaP-d<sub>12</sub> losses of ~40% were observed during the daylight sampling period on one of the two days, possibly mainly due to O<sub>3</sub> reaction. In contrast, no obvious overall losses of the perdeuterated fluoranthene or pyrene occurred, although

these PAH volatilized from the filters onto the downstream polyurethane foam (PUF) plugs. The data for perylene-d<sub>12</sub> loss was inconclusive.

The nitro-PAH isomers formed during the ambient air sampling were perdeuterated 3-nitroperylene, 6-nitroBaP, and 1-nitropyrene. The amounts formed followed the ranking expected from the reactivity of the parent PAH toward electrophilic nitration. The maximum nitro-PAH yield was 0.1% of 3-nitroperylene-d<sub>11</sub> and it was estimated that the 1-nitropyrene artifact, i.e., the amount formed during sample collection, was only ~0.1% of the 1-nitropyrene present on the particles. Thus, the formation of nitro-PAH under ambient conditions is likely to be unimportant, especially if most of the ambient particle-adsorbed PAH are unavailable for reaction.

### 3. Photolysis

Numerous studies have been carried out to investigate the photolysis and photooxidation of BaP and other PAH on various surfaces (Korfmacher et al., 1980; Blau and Gsten, 1981; Behymer and Hites, 1985, 1988; Kamens et al., 1985, 1986; Yokley et al., 1986; Valerio et al., 1987). All studies employing a range of substrates have concluded that photolysis rates are highly substrate dependent (Korfmacher et al., 1980; Behymer and Hites, 1985, 1988; Yokley et al., 1986), with the darker substrates leading to lower photolysis rates, presumably due to stabilization of the PAH incorporated in the particles (Behymer and Hites, 1985, 1988; Yokley et al., 1986).

Kamens et al. (1988) observed that the PAH photolysis rates (on freshly generated wood and gasoline combustion soot) depended on the temperature and ambient water concentration, with, for example, a BaP lifetime at 293 K and ~50% relative humidity at midday of ~1 hr. In contrast, for dark fly ashes, half-lives on the order of several days were observed for BaP (Behymer and Hites, 1988). The relative reactivities of the alternant PAH followed the general reactivity ranking shown in Table 13. Specifically, Behymer and Hites (1988) found for the following PAH isomer pairs that phenanthrene is more stable than anthracene, chrysene is more stable than benz[a]anthracene, benzo[e]pyrene is more stable than BaP and benzo[ghi]perylene is more stable than the linear PAH anthanthrene. Once again it can be concluded that PAH are reactive, and thus may be expected to photolyze in ambient atmospheres, but the extent to which this occurs will depend on the exact nature of the adsorbed state.

Cope and Kalkwarf (1987) and Kamens et al. (1989) have investigated the photooxidation of certain adsorbed oxygenated PAH and have found these to be generally stable to sunlight, but to decay under the influence of light and  $O_3$ .

#### 4. Ambient Air Data

Further information concerning the atmospheric reactions and lifetimes of PAH arises from ambient air data. Arey et al. (1988) have shown that under wintertime high- $NO_x$  conditions in southern California (with mid-day temperatures attaining 95 °F) losses of BaP-d<sub>12</sub> during 12-hr sampling periods were ≤40%, and nitration of BaP-d<sub>12</sub> was of negligible importance. Losses of particle-incorporated BaP would, presumably, have been lower. From an analysis of ambient air and combustion emissions data, Schuetzle and Frazier (1986) found that the benzo[e]pyrene/benzo[a]pyrene ratio (BeP/BaP) in ambient air of ~1.8 was very similar to the ratio observed in emission sources. Since BaP is much more reactive than BeP toward electrophilic reaction (Nielsen, 1984), reaction with  $O_3$  (Van Vaeck and Van Cauwenberghe, 1984) and photolysis (Behymer and Hites, 1988), this observation suggests that once the BaP and BeP are adsorbed on particles they do not undergo significant degradation in the atmosphere.

Nielsen (1988) investigated the dependence of the BaP/BeP and cyclopenta[cd]pyrene/chrysene + triphenylene ratios on the wind direction, and hence the vicinity of the source, in Denmark. Higher PAH concentrations were associated with local sources, and the BaP/BeP and cyclopenta[cd]pyrene/chrysene + triphenylene ratios were found to correlate with the BeP and chrysene + triphenylene concentrations, respectively, suggesting that BaP and cyclopenta[cd]pyrene were being degraded in the atmosphere. The author concluded that the decay rates of these two PAH were relatively fast (Nielsen, 1988). From ambient winter and summer measurements in New Jersey, Greenberg (1989) confirmed the findings of Nielsen that BaP and, especially, cyclopenta[cd]pyrene are reactive under atmospheric conditions.

Atkinson et al. (1988) measured PAH and PAH-derivatives at seven sites throughout California. Consistent with the work of Nielsen, the sites with the highest BaP/BeP ratios, Mammoth Lakes (BaP/BeP = 1.5) and Concord (BaP/BeP = 1.3), also had the highest BeP concentrations. Furthermore the BaP/BeP ratio was lowest at Glendora (BaP/BeP = 0.35) a

site where significant atmospheric reaction had occurred, as evidenced by the formation of 2-nitropyrene from pyrene (Atkinson et al., 1988). (For a complete discussion of the PAH transformations in the atmospheres sampled see Section X of Atkinson et al., 1988.) The apparent inconsistency in the data of Schuetzle and Frazier (1986) and that of Nielsen (1988) and Atkinson et al. (1988) probably reflects differences due to comparisons of averaged values from generally aged emissions (Schuetzle and Frazier, 1986) versus BaP concentrations resulting from a single emission source (such as wood smoke at Mammoth Lakes) and/or fresh emissions (Nielsen, 1988; Atkinson et al., 1988).

#### F. Atmospheric Formation Processes

There are no atmospheric formation pathways for PAH. PAH are present in essentially all combustion systems and sources of the PAH present in ambient air will include, in addition to diesel exhaust, emissions from gasoline-fueled automobiles, industrial processes, domestic heating systems, waste incineration facilities, tobacco smoking, agricultural burns and several natural sources, including forest fires and volcanic eruptions (Nikolaou et al., 1984).

In contrast to the PAH, nitro-PAH and other PAH derivatives (including hydroxy-PAH) are formed in the atmosphere from the gas- and adsorbed-phase reactions of the parent PAH, as well as being directly emitted from combustion sources. As discussed in Section D, the reactions of the gas-phase PAH appear to dominate in the formation of nitro-PAH (and other PAH derivatives) in the atmosphere. In ambient air studies conducted in California, the majority of the nitro-PAH observed are formed from the gas-phase atmospheric reactions of the parent PAH (Arey et al., 1987, 1989a, 1990a; Atkinson et al., 1988; Zielinska et al., 1989b). It should be noted that a large fraction of the atmospheric transformation products of the gas-phase PAH remain unidentified.

## G. References

- Arey, J., Zielinska, B., Atkinson, R., Winer, A. M., Ramdahl, T. and Pitts, J. N., Jr. (1986): The formation of nitro-PAH from the gas-phase reactions of fluoranthene and pyrene with the OH radical in the presence of  $\text{NO}_x$ . *Atmos. Environ.*, 20, 2339-2345.
- Arey, J., Zielinska, B., Atkinson, R., and Winer A. M. (1987): Polycyclic aromatic hydrocarbon and nitroarene concentrations in ambient air during a winter-time high- $\text{NO}_x$  episode in the Los Angeles basin. *Atmos. Environ.* 21, 1437-1444.
- Arey, J., Zielinska, B., Atkinson, R., and Winer, A. M. (1988): Formation of nitroarenes during ambient high-volume sampling. *Environ. Sci. Technol.*, 22, 457-462.
- Arey, J., Atkinson, R., Zielinska, B. and McElroy, P. A. (1989a): Diurnal concentrations of volatile polycyclic aromatic hydrocarbons and nitroarenes during a photochemical air pollution episode in Glendora, California. *Environ. Sci. Technol.*, 23, 321-327.
- Arey, J., Zielinska, B., Atkinson, R. and Aschmann, S. M. (1989b) Nitroarene products from the gas-phase reactions of volatile polycyclic aromatic hydrocarbons with the OH radical and  $\text{N}_2\text{O}_5$ . *Int. J. Chem. Kinet.*, 21, 775-799.
- Arey, J., Zielinska, B., Atkinson, R. and McElroy, P. A. (1990a): Ambient air sampling at seven locations in California for polycyclic aromatic compounds and mutagenicity. Part II. Concentrations of selected polycyclic aromatic hydrocarbons and nitroarenes. *Environ. Sci. Technol.*, to be submitted.
- Arey, J., Atkinson, R., Aschmann, S. M. and Schuetzle, D. (1990b): Experimental investigation of the atmospheric chemistry of 2-methyl-1-nitronaphthalene and a comparison of predicted nitroarene concentrations with ambient air data. *Polycyclic Aromat. Compounds*, 1, 33-50.
- Atkinson, R. (1986): Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. *Chem. Rev.*, 86, 69-201.
- Atkinson, R. (1988): Atmospheric Transformations of Automotive Emissions. In: Air Pollution, the Automobile, and Public Health, A. Y. Watson, R. R. Bates and D. Kennedy (Eds.), National Academy Press, Washington, DC, pp. 99-132.
- Atkinson, R. (1989): Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. *J. Phys. Chem. Ref. Data*, Monograph. 1, 1-246.
- Atkinson, R. (1990a): Kinetics and mechanisms of the gas-phase reactions of the  $\text{NO}_3$  radical with organic compounds. *J. Phys. Chem. Ref. Data*, in press.



- Atkinson, R. (1990b): Gas-phase tropospheric chemistry of organic compounds: a review. *Atmos. Environ.*, 24A, 1-41.
- Atkinson, R. and Aschmann, S. M. (1985): Rate constants for the gas-phase reaction of hydroxyl radicals with biphenyl and the monochlorobiphenyls at  $295 \pm 1$  K. *Environ. Sci. Technol.*, 19, 462-464.
- Atkinson, R. and Aschmann, S. M. (1986): Kinetics of the reactions of naphthalene, 2-methylnaphthalene, and 2,3-dimethylnaphthalene with OH radicals and with  $O_3$  at  $295 \pm 1$  K. *Int. J. Chem. Kinet.*, 18, 569-573.
- Atkinson, R. and Aschmann, S. M. (1987): Kinetics of the gas-phase reactions of alkylnaphthalenes with  $O_3$ ,  $N_2O_5$  and OH radicals at  $298 \pm 2$  K. *Atmos. Environ.*, 21, 2323-2326.
- Atkinson, R. and Aschmann, S. M. (1988): Kinetics of the reactions of acenaphthene and acenaphthylene and structurally-related aromatic compounds with OH and  $NO_3$  radicals,  $N_2O_5$  and  $O_3$  at  $296 \pm 2$  K. *Int. J. Chem. Kinet.*, 20, 513-539.
- Atkinson, R., Aschmann, S. M. and Pitts, J. N., Jr. (1984): Kinetics of the reactions of naphthalene and biphenyl with OH radicals and with  $O_3$  at  $294 \pm 1$  K. *Environ. Sci. Technol.*, 18, 110-113.
- Atkinson, R., Winer, A. M. and Pitts, J. N., Jr. (1986): Estimation of night-time  $N_2O_5$  concentrations from ambient  $NO_2$  and  $NO_3$  radical concentrations and the role of  $N_2O_5$  in night-time chemistry. *Atmos. Environ.*, 20, 331-339.
- Atkinson, R., Arey, J., Zielinska, B. and Aschmann, S. M. (1987): Kinetics and products of the gas-phase reactions of OH radicals and  $N_2O_5$  with naphthalene and biphenyl. *Environ. Sci. Technol.*, 21, 1014-1022.
- Atkinson, R., Arey, J., Winer, A. M., Zielinska, B., Dinoff, T. M., Harger, W. P. and McElroy, P. A. (1988): A Survey of Ambient Concentrations of Selected Polycyclic Aromatic Hydrocarbons (PAH) at Various Locations in California. Final Report, California Air Resources Board Contract No. A5-185-32, May.
- Atkinson, R., Aschmann, S. M., Arey, J., Zielinska, B. and Schuetzle, D. (1989a): Gas-phase atmospheric chemistry of 1- and 2-nitronaphthalene and 1,4-naphthoquinone. *Atmos. Environ.*, 23, 2679-2690.
- Atkinson, R., Aschmann, S. M., Arey, J. and Carter, W. P. L. (1989b): Formation of ring-retaining products from the OH radical-initiated reactions of benzene and toluene. *Int. J. Chem. Kinet.*, 21, 801-827.
- Atkinson, R., Arey, J., Zielinska, B. and Aschmann, S. M. (1990a): Kinetics and nitro-products of the gas-phase OH and  $NO_3$  radical-initiated reactions of naphthalene- $d_8$ , fluoranthene- $d_{10}$  and pyrene. *Int. J. Chem. Kinet.*, in press.

- Atkinson, R., Aschmann, S. M. and Arey, J. (1990b): Formation of ring-retaining products from the OH radical-initiated reactions of o-, m- and p-xylene. *Int. J. Chem. Kinet.*, submitted for publication.
- Atkinson, R., Tuazon, E. C. and Arey, J. (1990c): Reactions of naphthalene in  $\text{N}_2\text{O}_5$  -  $\text{NO}_3$  -  $\text{NO}_2$  - air mixtures. *Int. J. Chem. Kinet.*, in press.
- Barbella, R., Bertoli, C., Ciajolo, A. and D'Anna, A. (1988): Soot and unburnt liquid hydrocarbon emissions from diesel engines. *Combust. Sci. and Tech.*, 59, 183-198.
- Barbella, R., Ciajolo, A., D'Anna, A. and Bertoli, C. (1989): Effect of fuel aromaticity on diesel emissions. *Combust. and Flame*, 77, 267-277.
- Behymer, T. D. and Hites, R. A. (1985): Photolysis of polycyclic aromatic hydrocarbons adsorbed on simulated atmospheric particulates. *Environ. Sci. Technol.*, 19, 1004-1006.
- Behymer, T. D. and Hites, R. A. (1988): Photolysis of polycyclic aromatic hydrocarbons adsorbed on fly ash. *Environ. Sci. Technol.*, 22, 1311-1319.
- Benson, J. M., Brooks, A. L., Cheng, Y. S., Henderson, T. R. and White, J. E. (1985): Environmental transformation of 1-nitropyrene on glass surfaces. *Atmos. Environ.*, 19, 1169-1174.
- Bidleman, T. F. (1988): Atmospheric processes. *Environ. Sci. Technol.*, 22, 361-367.
- Bidleman, T. F. and W. T. Foreman (1987): Vapor-particle partitioning of semivolatile organic compounds. In: Sources and Fates of Aquatic Pollutants, R. A. Hites and S. J. Eisenreich (Eds.), *Adv. Chem. Series*, 216, pp. 27-56.
- Biermann, H. W., MacLeod, H., Atkinson, R., Winer, A. M. and Pitts, J. N., Jr. (1985): Kinetics of the gas-phase reactions of the hydroxyl radical with naphthalene, phenanthrene, and anthracene. *Environ. Sci. Technol.*, 19, 244-248.
- Blau, L. and Gusten, H. (1981): Quantum yields of the photodecomposition of polynuclear aromatic hydrocarbons adsorbed on silica gel. In: Aromatic Hydrocarbons, M. Cooke, A. J. Dennis and G. L. Fisher (Eds.), Battelle Press, Columbus, OH, pp. 133-144.
- Bockhorn, H., Fetting, F., Wannemacher, G. and Wenz, H. W. (1982): Optical studies of soot particle growth in hydrocarbon oxygen flames. 19th Int. Symp. on Combustion, 1982; The Combustion Institute, Pittsburgh, PA, pp. 1413-1420.
- Brorström, E., Grennfelt, P. and Lindskog, A. (1983): The effect of nitrogen dioxide and ozone on the decomposition of particle-associated polycyclic aromatic hydrocarbons during sampling from the atmosphere. *Atmos. Environ.*, 17, 601-605.

- Brorström-Lundén, E. and Lindskog, A. (1985): Degradation of polycyclic aromatic hydrocarbons during simulated stack gas sampling. *Environ. Sci. Technol.*, 19, 313-316.
- Butler, J. D. and Crossley, P. (1981): Reactivity of polycyclic aromatic hydrocarbons adsorbed on soot particles. *Atmos. Environ.*, 15, 91-94.
- Ciajolo, A., Barbella, R., Mattiello, M., and D'Alessio, A. (1982): Axial and radial measurements of soot and PAH in a light oil flame. 19th Int. Symp. on Combustion, 1982; The Combustion Institute, Pittsburgh, PA, pp. 1369-1377.
- Cope, V. W. and Kalkwarf, D. R. (1987): Photooxidation of selected polycyclic aromatic hydrocarbons and pyrenequinones coated on glass surfaces. *Environ. Sci. Technol.*, 21, 643-648.
- Coutant, R. W., Brown, L., Chuang, J. C., Riggin, R. M. and Lewis, R. G. (1988): Phase distribution and artifact formation in ambient air sampling for polynuclear aromatic hydrocarbons. *Atmos. Environ.*, 22, 403-409.
- Daisey, J. M., Lewandowski, C. G. and Zorz, M. (1982): A photoreactor for investigations of the degradation of particle-bound polycyclic aromatic hydrocarbons under simulated atmospheric conditions. *Environ. Sci. Technol.*, 16, 857-861.
- Dolan, D. F., Kittelson, D. B. and Pui, D. Y. H. (1980): Diesel exhaust particle size distribution measurement techniques. SAE Technical Paper 800187, SAE Trans.
- Eisenreich, S. J., Looney B. B. and Thornton, J. D. (1981): Airborne organic contamination in the Great Lakes ecosystem. *Environ. Sci. Technol.*, 15, 30-38.
- Flessel, P., Guirguis, G., Cheng, J., Chang, K., Hahn, E., Chan, R., Ondo, J., Fenske, R. Twiss, S., Vance, W. and Wesolowski, J. (1984): Monitoring of Mutagens and Carcinogens in Community Air. Final Report, California Air Resources Board Contract No. A1-029-32, May.
- Frenklach, M. (1989): On the driving force of soot formation. 22nd Int. Symp. on Combustion, 1988; The Combustion Institute, Pittsburgh, PA, pp. 1075-1082.
- Frenklach, M., Clary, D. W., Gardiner, W. C., Jr. and Stein, S. E. (1988): Effect of fuel structure on pathways to soot. 21st Int. Symp. on Combustion, 1986; The Combustion Institute, Pittsburgh, PA, pp. 1067-1076.
- Gibson, T. L. (1982): Nitroderivatives of polynuclear aromatic hydrocarbons in airborne and source particulate matter. *Atmos. Environ.*, 16, 2037-2040.
- Gibson, T. L. (1983): Sources of direct-acting nitroarene mutagens in airborne particulate matter. *Mutat. Res.*, 122, 115-121.

- Globlicki, P. J. and Begeman, C. R. (1979): Particle size variation in diesel car exhaust. SAE Technical Paper 790421, SAE Trans.
- Graedel, T. E., and Weschler, C. J. (1981): Chemistry within aqueous atmospheric aerosols and raindrops. J. Geophys. Res., 19, 505-539.
- Greenberg, A (1989): Phenomenological study of benzo[a]pyrene and cyclopenteno[cd]pyrene decay in ambient air using winter/summer comparisons. Atmos. Environ., 23, 2797-2799.
- Greenberg, A. and Darack, F. B. (1987): Atmospheric Reactions and Reactivity Indices of Polycyclic Aromatic Hydrocarbons. In Molecular Structures and Energetics, J. L. Liebman and A. Greenberg (Eds), VCH Publishers, Inc., New York, NY, pp. 1-47.
- Grosjean, D., Fung, K. and Harrison, J. (1983): Interactions of polycyclic aromatic hydrocarbons with atmospheric pollutants. Environ. Sci. Technol., 17, 673-679.
- Güsten, H. (1986): Photocatalytic degradation of atmospheric pollutants on the surface of metal oxides. In: Chemistry of Multiphase Atmospheric Systems, W. Jaeschke (Ed.), Springer-Verlag, Berlin, pp. 567-592.
- Hayano, S., Jang-Ho, L., Furuya, K., Kikuchi, T., Someya, T., Oikawa, C., Iida, Y., Matsushita, H., Kinouchi, T., Manabe, Y. and Ohnishi, Y. (1985): Formation of hazardous substances and mutagenicity of PAH produced during the combustion process in a diesel engine. Atmos. Environ., 19, 1009-1015.
- Henderson, T. R., Sun, J. D., Li, A. P., Hanson, R. L., Bechtold, W. E., Harvey, T. M., Shabanowitz, J. and Hunt D. F. (1984): GC/MS and MS/MS studies of diesel exhaust mutagenicity and emissions from chemically defined fuels. Environ. Sci. Technol., 18, 428-434.
- Holloway, M. P., Biaglow, M. C., McCoy, E. C., Anders, M., Rosenkranz, H. S. and Howard, P. C. (1987): Photochemical instability of 1-nitropyrene, 3-nitrofluoranthene, 1,8-dinitropyrene and their parent polycyclic aromatic hydrocarbons. Mutat. Res., 187, 199-207.
- Hughes, M. M., Natusch, D. F. S., Taylor, R. R. and Zeller, M. V. (1980): Chemical Transformations of Particulate Polycyclic Organic Matter. In Polynuclear Aromatic Hydrocarbons: Chemistry and Biological Effects, A. Bjorseth and A. J. Dennis (Eds), Battelle Press, Columbus, OH, pp. 1-8.
- IARC Monographs (1983): IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Polynuclear Aromatic Compounds, Part 1, Chemical, Environmental and Experimental Data, Vol. 32, International Agency for Research on Cancer, France, pp. 211-224.
- Jäger, J. and Rakovic, M. (1974): Sulphur-dioxide-induced qualitative changes in polycyclic aromatic hydrocarbons adsorbed on solid carriers. J. Hyg. Epidemio. Microbio. Immunol., 2, 137-143.

- Jäger, J. and Hanus, V. (1980): Reaction of solid carrier-adsorbed polycyclic aromatic hydrocarbons with gaseous low-concentrated nitrogen dioxide. *J. Hyg. Epidemio. Microbio. Immunol.*, 24, 1-12.
- Jensen, T. E. and Hites, R. A. (1983): Aromatic diesel emissions as a function of engine conditions. *Anal. Chem.*, 55, 594-599.
- Johnston, J. H. (1988) Automotive Emissions. In: Air Pollution, the Automobile, and Public Health, A. Y. Watson, R. R. Bates and D. Kennedy (Eds.), National Academy Press, Washington, DC, pp. 39-75.
- Jonsson, A., Persson, K. A. and Grigoriadis, V. (1985): Measurements of some low molecular-weight oxygenated, aromatic, and chlorinated hydrocarbons in ambient air and in vehicle emissions. *Environ. Int.*, 11, 383-392.
- Kamens, R. M., Perry, J. M., saucy, D. A., Bell, D. A., Newton, D. L. and Brand, B. (1985a): Factors which influence polycyclic aromatic hydrocarbon decomposition on wood smoke particles. *Environ. Int.*, 11, 131-136.
- Kamens, R., Bell, D., Dietrich, A., Perry, J., Goodman, R., Claxton, L. and Tejada, S. (1985b): Mutagenic transformations of dilute wood smoke systems in the presence of ozone and nitrogen dioxide. Analysis of selected high-pressure liquid chromatography fractions from wood smoke particle extracts. *Environ. Sci. Technol.*, 19, 63-69.
- Kamens, R. M., Fulcher, J. N. and Zhishi, G. (1986): Effects of temperature on wood soot PAH decay in atmospheres with sunlight and low NO<sub>x</sub>. *Atmos. Environ.*, 20, 1579-1587.
- Kamens, R. M., Zhishi, G., Fulcher, J. N. and Bell, D. A. (1988): Influence of humidity, sunlight, and temperature on the daytime decay of polyaromatic hydrocarbons on atmospheric soot particles. *Environ. Sci. Technol.*, 22, 103-108.
- Kamens, R. M., Karam, H., Guo, J., Perry, J. M. and Stockburger, L. (1989): The behavior of oxygenated polycyclic aromatic hydrocarbons on atmospheric soot particles. *Environ. Sci. Technol.*, 23, 801-806.
- Katz, M., Chan, C., Tosine, H. and Sakuma, T. (1979): Relative rates of photochemical and biological (in vitro) oxidation of polynuclear aromatic hydrocarbons. In: Polynuclear Aromatic Hydrocarbons, P. W. Jones and P. Leber (Eds.), Ann Arbor Science Publishers, Ann Arbor, MI, pp. 171-189.
- King, C. M. (1988): Metabolism and biological effects of nitropyrene and related compounds. Report No. 16, Health Effects Institute, Cambridge, MA, February.
- Kittelson, D. B., Du, C.-J. and Zweidinger, R. B. (1985): Measurements of polycyclic aromatic compounds in the cylinders of an operating diesel engine. EPA-600/D-85/012, January.

- Klöpffer, W., Frank, R., Kohl, E.-G. and Haag, F. (1986): Quantitative erfassung der photochemischen transformationsprozesse in der troposphäre. *Chemiker-Zeitung*, 110, 57-61.
- Korfmacher, W. A., Natusch, D. F. S., Taylor, D. R., Mamantov, G. and Wehry, E. L. (1980a): Oxidative transformations of polycyclic aromatic hydrocarbons adsorbed on coal fly ash. *Science*, 207, 763-765.
- Korfmacher, W. A., Wehry, E. L., Mamantov, G. and Natusch, D. F. S. (1980b): Resistance to photochemical decomposition of polycyclic aromatic hydrocarbons vapor-adsorbed on coal fly ash. *Environ. Sci. Technol.*, 14, 1094-1099.
- Korfmacher, W. A., Mamantov, G., Wehry, E. L., Natusch, D. F. S. and Mauney, T. (1981): Nonphotochemical decomposition of fluorene vapor-adsorbed on coal fly ash. *Environ. Sci. Technol.*, 15, 1370-1375.
- Kroto, H. (1988): Space, stars, C<sub>60</sub>, and soot. *Science*, 242, 1139-1142.
- Lane, D. A. and Katz, M. (1977): The photomodification of benzo[a]pyrene, benzo[b]fluoranthene and benzo[k]fluoranthene under simulated conditions. *Adv. Environ. Sci. Technol.*, 8, 137-154.
- Lee, P. S., Gorski, R. A., Johnston, J. T. and Soderholm, S. C. (1989): Generation of diesel particles coated with polycyclic aromatic compounds - evidence suggesting that dinitropyrene formation is a collection artifact. *J. Aerosol Sci.*, 20, 627-637.
- Leuenberger, C., Ligocki, M. P., and Pankow, J. F. (1985): Trace organic compounds in rain. 4. Identities, concentrations and scavenging mechanisms for phenols in urban air and rain. *Environ. Sci. Technol.* 19, 1053-1058.
- Ligocki, M. P., Leuenberger, C., and Pankow, J. F. (1985a): Trace organic compounds in rain - III. Particle scavenging of neutral organic compounds. *Atmos. Environ.* 19, 1619-1626.
- Ligocki, M. P., Leuenberger, C. and Pankow, J. F. (1985b): Trace organic compounds in rain - II. Gas scavenging of neutral organic compounds. *Atmos. Environ.*, 19, 1609-1617.
- Lindskog, A., Brorström-Lundén, E. and Sjödin, Å. (1985): Transformation of reactive PAH on particles by exposure to oxidized nitrogen compounds and ozone. *Environ. Internat.*, 11, 125-130.
- Logan, J. A. (1985): Tropospheric ozone: seasonal behavior, trends, and anthropogenic influence. *J. Geophys. Res.*, 90, 10463-10482.
- Lorenz, K. and Zellner, R. (1983): Kinetics of the reactions of OH-radicals with benzene, benzene-d<sub>6</sub> and naphthalene. *Ber. Bunsenges. Phys. Chem.*, 87, 629-636.

- Lorenz, K. and Zellner, R. (1984): presented at 8th Int. Symp. on Gas Kinetics, Nottingham, U. K., July 15-20; private communication to R. Atkinson, 1985.
- Mackay, D., Bobra, A., Chan, D. W. and Shiu, W. Y. (1982): Vapor pressure correlations for low-volatility chemicals. *Environ. Sci. Technol.*, 16, 645-649.
- Macknick, A. B. and Prausnitz, J. M. (1979): Vapor pressures of high-molecular-weight hydrocarbons. *J. Chem. Eng. Data*, 24, 175-178.
- Menard, H., Lalancette, J. M., Lefebvre, Y. and Belanger, D. (1986): Ozone degradation of polycyclic aromatic hydrocarbons adsorbed on asbestos. *Anal. Lett.*, 19, 307-314.
- Nakagawa, R., Kitamori, S., Horikawa, K., Nakashima, K. and Tokiwa, H. (1983): Identification of dinitropyrenes in diesel-exhaust particles. Their probable presence as the major mutagens. *Mutat. Res.*, 124, 201-211.
- National Research Council (1982): Diesel Cars: Benefits, Risks, and Public Policy. Diesel Impacts Study Committee, National Research Council, National Academy Press, Washington, D. C., 142 pp.
- National Research Council (1983): Polycyclic Aromatic Hydrocarbons: Evaluation of Sources and Effects. The Committee on Pyrene and Selected Analogues, National Research Council, National Academy Press, Washington, D.C.
- Nielsen, T. (1984): Reactivity of polycyclic aromatic hydrocarbons toward nitrating species. *Environ. Sci. Technol.*, 18, 157-163.
- Nielsen, T. (1988): The decay of benzo[a]pyrene and cyclopenteno[cd]-pyrene in the atmosphere. *Atmos. Environ.*, 22, 2249-2254.
- Nielsen, T. and Ramdahl, T. (1986): Discussion on "Determination of 2-nitrofluoranthene and 2-nitropyrene in ambient particulate matter: evidence for atmospheric reactions". *Atmos. Environ.*, 20, 1507.
- Nielsen, T., Ramdahl, T. and Bjorseth, A. (1983): The fate of airborne polycyclic organic matter. *Environ. Health Perspec.*, 47, 103-114.
- Nielsen, T., Seitz, B. and Ramdahl, T. (1984): Occurrence of nitro-PAH in the atmosphere in a rural area. *Atmos. Environ.*, 18, 2159-2165.
- Nikolaou, K., Masclet, P. and Mouvier, G. (1984): Sources and chemical reactivity of polynuclear aromatic hydrocarbons in the atmosphere - a critical review. *Sci. Total Environ.*, 32, 103-132.
- Obuchi, A, Ohi, A., Aoyama, H. and Ohuchi, H. (1987): Evaluation of gaseous and particle emission characteristics of a single cylinder diesel engine. *Combustion and Flame*, 70, 215-224.

- Pankow, J. F. (1989): Overview of the gas phase retention volume behavior of organic compounds on polyurethane foam. *Atmos. Environ.*, 23, 1107-1111.
- Paputa-Peck, M. C., Marano, R. S., Schuetzle, D., Riley, T. L., Hampton, C. V., Prater, T. J., Skewes, L. M., Jensen, T. E., Ruehle, P. H., Bosch, L. C. and Duncan, W. P. (1983): Determination of nitrated polynuclear aromatic hydrocarbons in particulate extracts by capillary column gas chromatography with nitrogen selective detection. *Anal. Chem.*, 55, 1946-1954.
- Pederson, T. C. and Siak, J.-S. (1981): The role of nitro aromatic compounds in the direct-acting mutagenicity of diesel exhaust particle extracts. *J. Appl. Toxicol.*, 1, 54-60.
- Peters, J. and Seifert, B. (1980): Losses of benzo(a)pyrene under the conditions of high-volume sampling. *Atmos. Environ.*, 14, 117-119.
- Pierson, W. R., Gorse, R. A., Jr., Szkariat, A. C., Brachaczek, W. W., Japar, S. M., Lee, F. S.-C., Zweidinger, R. B. and Claxton, L. D. (1983): Mutagenicity and chemical characteristics of carbonaceous particulate matter from vehicles on the road. *Environ. Sci. Technol.*, 17, 31-44.
- Pitts, J. N., Jr., Van Cauwenberghe, K. A., Grosjean, D., Schmid, J. P., Fitz, D. R., Belser, W. L., Jr., Knudson, G. B. and Hynds, P. M. (1978): Atmospheric reactions of polycyclic aromatic hydrocarbons: facile formation of mutagenic nitro derivatives. *Science*, 202, 515-519.
- Pitts, J. N., Jr., Lokensgard, D. M., Ripley, P. S., Van Cauwenberghe, K. A., Van Vaeck, L., Shaffer, S. D., Thill, A. J. and Belser, W. L., Jr. (1980): "Atmospheric" epoxidation of benzo(a)pyrene by ozone: formation of the metabolite benzo(a)pyrene-4,5-oxide. *Science*, 210, 1347-1349.
- Pitts, J. N., Jr., Atkinson, R., Sweetman, J. A. and Zielinska, B. (1985a): The gas-phase reaction of naphthalene with  $N_2O_5$  to form nitronaphthalenes. *Atmos. Environ.*, 19, 701-705.
- Pitts, J. N. Jr., Sweetman, J. A., Zielinska, B., Winer, A. M. and Atkinson, R. (1985b): Determination of 2-nitrofluoranthene and 2-nitropyrene in ambient particulate organic matter: Evidence for atmospheric reactions. *Atmos. Environ.*, 19, 1601-1608.
- Pitts, J. N., Jr., Sweetman, J. A., Zielinska, B., Atkinson, R., Winer, A. M. and Harger, W. P. (1985c): Formation of nitroarenes from the reaction of polycyclic aromatic hydrocarbons with dinitrogen pentoxide. *Environ. Sci. Technol.*, 19, 1115-1121.
- Pitts, J. N., Jr., Paur, H.-R., Zielinska, B., Arey, J., Winer, A. M., Ramdahl, T. and Mejia, V. (1986): Factors influencing the reactivity of polycyclic aromatic hydrocarbons adsorbed on filters and ambient POM with ozone. *Chemosphere*, 15, 675-685.



- Prado, G., Garo, A., Ko, A. and Sarofim, A. (1985): Polycyclic aromatic hydrocarbons: formation and destruction in a laminar diffusion flame. 20th Int. Symp. on Combustion 1984; The Combustion Institute, Pittsburgh, PA, pp. 989-996.
- Prinn, R., Cunnold, D., Rasmussen, R., Simmonds, P., Alyea, F., Crawford, A., Fraser, P. and Rosen, R. (1987): Atmospheric trends in methylchloroform and the global average for the hydroxyl radical. *Science*, 238, 945-950.
- Ramdahl, T., Bjorseth, A., Lokensgard, D. M. and Pitts, J. N., Jr. (1984): Nitration of polycyclic aromatic hydrocarbons adsorbed to different carriers in a fluidized bed reactor. *Chemosphere*, 13, 527-534.
- Ramdahl, T., Zielinska, B., Arey, J., Atkinson, R., Winer, A. M. and Pitts, J. N., Jr. (1986): Ubiquitous occurrence of 2-nitrofluoranthene and 2-nitropyrene in air. *Nature*, 321, 425-427.
- Robbat, A., Jr., Corso, N. P., Doherty, P. J. and Wolf, M. H. (1986): Gas chromatographic chemiluminescent detection and evaluation of predictive models for identifying nitrated polycyclic aromatic hydrocarbons in a diesel fuel particulate extract. *Anal. Chem.*, 58, 2078-2084.
- Rosenkranz, H. S. and Mermelstein, R. (1983): Mutagenicity and genotoxicity of nitroarenes. All nitro-containing chemicals were not created equal. *Mutat. Res.*, 114, 217-267.
- Ross, D. S., Hum, G. P. and Schmitt, R. J. (1987): Diesel exhaust and pyrene nitration. *Environ. Sci. Technol.*, 21, 1130-1131.
- Ruehle, P. H., Bosch, L. C. and Duncan, W. P. (1985): Synthesis of nitrated polycyclic aromatic hydrocarbons. In Nitrated Polycyclic Aromatic Hydrocarbons, C. M. White (Ed), Dr. Alfred Huethig Verlag, Heidelberg, pp. 169-235.
- Schuetzle, D. (1983): Sampling of vehicle emissions for chemical analysis and biological testing. *Environ. Health Perspect.*, 47, 65-80.
- Schuetzle, D. and Perez, J. M. (1983): Factors influencing the emissions of nitrated-polynuclear aromatic hydrocarbons (nitro-PAH) from diesel engines. *J. Air Pollut. Control Assoc.*, 33, 751-755.
- Schuetzle, D. and Frazier, J. A. (1986): Factors influencing the emission of vapor and particulate phase components from diesel engines. In: Carcinogenic and Mutagenic Effects of Diesel Engine Exhaust, N. Ishinishi, A. Koizumi, R. O. McClellan and W. Stöber (Eds.), Elsevier Science Publishers B. V., pp. 41-63.
- Schuetzle, D., Lee, F. S.-C., Prater, T. J. and Tejada, S. B. (1981): The identification of polynuclear aromatic hydrocarbon (PAH) derivatives in mutagenic fractions of diesel particulate extracts. *Intern. J. Environ. Anal. Chem.*, 9, 93-144.

- Schuetzle, D., Riley, T. L., Prater, T. J., Harvey, T. M. and Hunt, D. F. (1982): Analysis of nitrated polycyclic aromatic hydrocarbons in diesel particulates. *Anal. Chem.*, 54, 265-271.
- Sonnefeld, W. J., Zoller, W. H. and May, W. E. (1983): Dynamic coupled-column liquid chromatographic determination of ambient temperature vapor pressures of polynuclear aromatic hydrocarbons. *Anal. Chem.*, 55, 275-280.
- Sweetman, J. A., Zielinska, B., Atkinson, R., Ramdahl, T., Winer, A. M. and Pitts, J. N., Jr. (1986): A possible formation pathway for the 2-nitrofluoranthene observed in ambient particulate organic matter. *Atmos. Environ.*, 20, 235-238.
- Takeda, N. and Teranishi, K. (1986): Effects of environmental conditions on polycyclic aromatic hydrocarbons and mutagenicity of airborne particulate matter. *Bull. Environ. Contam. Toxicol.*, 36, 685-692.
- Tokiwa, H. and Ohnishi, Y. (1986): Mutagenicity and carcinogenicity of nitroarenes and their sources in the environment. *CRC Critical Reviews in Toxicology*, 17, 23-60.
- Tokiwa, H., Nakagawa, R., Morita, K. and Ohnishi, Y. (1981): Mutagenicity of nitro derivatives induced by exposure of aromatic compounds to nitrogen dioxide. *Mutat. Res.*, 85, 195-205.
- Toqan, M., Farmayan, W. R., Beér, J. M., Howard, J. B. and Teare, J. D. (1985): PAH and soot formation in fuel-rich turbulent coal liquid and natural gas diffusion flames. 20th Int. Symp. on Combustion, 1984, The Combustion Institute, Pittsburgh, PA, pp. 1075-1081.
- Van Vaeck, L. and Van Cauwenberghe, K. (1984): Conversion of polycyclic aromatic hydrocarbons on diesel particulate matter upon exposure to ppm levels of ozone. *Atmos. Environ.*, 18, 323-328.
- Valerio, F., Bottino, P., Ugolini, D., Cimberle, M. R., Tozzi, G. A. and Frigerio, A. (1984): Chemical and photochemical degradation of polycyclic aromatic hydrocarbons in the atmosphere. *Sci. Total Environ.*, 40, 169-188.
- Valerio, F., Antolini, E. and Lazzarotto, A. (1987): A model to evaluate half-lives of PAHs naturally occurring on airborne particulate. *Intern. J. Environ. Anal. Chem.*, 28, 185-196.
- Williams, R. L. (1982): Diesel particulate emissions. In: Toxicological Effects of Emissions from Diesel Engines, J. Lewtas (Ed.), Elsevier Science Publishing Co., Inc., New York, NY, pp. 15-32.
- Williams, P. T., Bartle, K. D. and Andrews, G. E. (1986): The relation between polycyclic aromatic compounds in diesel fuels and exhaust particulates. *Fuel*, 65, 1150-1158.

- Williams, D. J., Milne, J. W., Quigley, S. M., Roberts, D. B. and Kimberlee, M. C. (1989): Particulate emissions from "in-use" motor vehicles-II. Diesel vehicles. *Atmos. Environ.*, 23, 2647-2661.
- Wu, C.-H. and Niki, H. (1985): Fluorescence spectroscopic study of kinetics of gas-surface reactions between nitrogen dioxide and adsorbed pyrene. *Environ. Sci. Technol.*, 19, 1089-1094.
- Wu, C.-H., Salmeen, I. and Niki, H. (1984): Fluorescence spectroscopic study of reactions between gaseous ozone and surface-adsorbed polycyclic aromatic hydrocarbons. *Environ. Sci. Technol.*, 18, 603-607.
- Xu, X. B., Nachtman, J. P., Jin, Z. L., Rappaport, S. M. and Burlingame, A. L. (1982): Isolation and identification of mutagenic nitro-PAH in diesel-exhaust particulates. *Anal. Chimica Acta*, 136, 163-174.
- Yamasaki, H., Kuwata, K. and Kuge, Y. (1984): Determination of vapor pressure of polycyclic aromatic hydrocarbons in the supercooled liquid phase and their adsorption on airborne particulate matter. *J. Chem. Soc. Japan*, 8, 1324-1329.
- Yokley, R. A., Garrison, A. A., Mamantov, G. and Wehry, E. L. (1985): The effect of nitrogen dioxide on the photochemical and nonphotochemical degradation of pyrene and benzo[a]pyrene adsorbed on coal fly ash. *Chemosphere*, 14, 1771-1778.
- Yokley, R. A., Garrison, A. A., Wehry, E. L. and Mamantov, G. (1986): Photochemical transformation of pyrene and benzo[a]pyrene vapor-deposited on eight coal stack ashes. *Environ. Sci. Technol.*, 20, 86-90.
- Zetzsch, C. (1982): 15th Informal Conference on Photochemistry, Stanford, California, June 27-July 1.
- Zielinska, B., Arey, J., Atkinson, R. and McElroy, P. A. (1988): Nitration of acephenanthrylene under simulated atmospheric conditions and in solution and the presence of nitroacephenanthrylene(s) in ambient air. *Environ. Sci. Technol.*, 22, 1044-1048.
- Zielinska, B., Arey, J., Atkinson, R. and McElroy, P. A. (1989a): Formation of methylnitronaphthalenes from the gas-phase reactions of 1- and 2-methylnaphthalene with OH radicals and N<sub>2</sub>O<sub>5</sub> and their occurrence in ambient air. *Environ. Sci. Technol.*, 23, 723-729.
- Zielinska, B., Arey, J., Atkinson, R. and Winer, A. M. (1989b): The nitroarenes of molecular weight 247 in ambient particulate samples collected in Southern California. *Atmos. Environ.*, 23, 223-229.

\*00001329\*



ASSET